

A New Combination in *Chonemorpha* (Apocynaceae)

David J. Middleton

Department of Botany, Trinity College, Dublin 2, Ireland

ABSTRACT. The new combination *Chonemorpha verrucosa* is made, and a key to the four species of *Chonemorpha* in Thailand is given.

***Chonemorpha verrucosa* (Blume) D. J. Middleton**, comb. nov. Basionym: *Tabernaemontana verrucosa* Blume, Bijdr. 1029. 1826. *Rhynchodia verrucosa* (Blume) Woodson, Sunyatsenia 3: 102. 1936. TYPE: *Blume s.n.* (holotype, L; photograph in K).

Chonemorpha was first described by G. Don in 1837, although many of the 13 species he described have since been referred to other genera (see Chatterjee, 1947). The oldest name now referable to *Chonemorpha* is *C. fragrans* (Moon) Alston based on a drawing and description by Rheede (1689), which he called "Belutta-kaka-kodi," and validated as *Echites fragrans* by Moon (1824) who referred to Rheede. This species encompasses many of the subsequently published names. *Chonemorpha* is a conserved name against *Belutta-kak* Adanson, 1763.

Rhynchodia was first validly published in Benth & Hooker (1876), although de Candolle (1844) and Miquel (1857) had previously published this genus under the names *Rhyncospermum* and *Cercocoma*, respectively. These are illegitimate later homonyms of *Rhyncospermum* Reinwardt (1828) and *Cercocoma* Wallich ex G. Don (1837). A number of species have been described over the large range of the genus, but it appears that they are all referable to one variable species.

Pichon (1950) placed both *Chonemorpha* and *Rhynchodia* in his subtribe Chonemorphinae of tribe Parsonsieae, while Lý (1986) described a new tribe, Anodendreae, containing both genera. Lý further suggested that *Chonemorpha* evolved from *Rhynchodia*. These two genera are extremely similar in leaf type, calyx and corolla structure, and stamen, disk, and gynoecium type. Because the only aspect separating the two genera is the smaller flowers in *Rhynchodia*, I believe they should be united.

Chonemorpha appears to be most closely related to *Epigynum* and *Trachelospermum*.

Kerr (1939) included two species, under the names *C. grandieriana* Pierre ex Spire and *C. macrantha*

Pitard, in his list of the species in Thailand. These are synonyms of *C. fragrans* (Moon) Alston and *C. griffithii* J. D. Hooker, respectively. A key to the four species of *Chonemorpha* found in Thailand follows:

KEY TO THE SPECIES OF *CHONEMORPHA* IN THAILAND

- 1a. Calyx 3–4.5 mm long; corolla tube 4.3–7 mm long *C. verrucosa* (Blume) D. J. Middleton
- 1b. Calyx 1–3.9 cm long; corolla tube 2.8–6.2 cm long.
 - 2a. Sepals connate for about half their length, sparsely puberulent or glabrous; corolla tube 4.7–6.2 cm long . . . *C. fragrans* (Moon) Alston
 - 2b. Sepals connate for most of their length or almost free, densely puberulent or tomentose; corolla tube 2.8–3.9 cm long.
 - 3a. Sepals almost free to base
. *C. griffithii* J. D. Hooker
 - 3b. Sepals connate for most of their length
. *C. megacalyx* Pierre ex Spire

Acknowledgments. I thank the Royal Society, London, and the Royal Irish Academy, Dublin, for financial support for my work on Thai Apocynaceae, and John Parnell for comments on the manuscript.

Literature Cited

- Benth, G. & J. D. Hooker. 1876. Apocynaceae. In: Genera Plantarum 2: 681–728. Reeve, London.
- Candolle, A. L. P. de. 1844. Apocynaceae. In: Prodromus 8: 317–489. Treuttel & Würtz, Paris.
- Chatterjee, D. 1947. The genus *Chonemorpha* G. Don (Apocynaceae). Kew Bull. 1947: 47–52.
- Don, G. 1837. Apocynaceae. In: A General History of the Dichlamydeous Plants 4: 69–105. Rivington, London.
- Kerr, A. F. G. 1939. Apocynaceae. In: Flora Siamensis Enumeratio 2: 422–476. Siam Society, Bangkok.
- Lý, T. D. 1986. Die Familie Apocynaceae Juss. in Vietnam. Feddes Repert. 97: 235–273, 405–466, 607–689.
- Miquel, F. A. W. 1857. Apocynaceae. In: Flora van nederlandsch Indië 2: 384–459. Van de Post, Amsterdam.
- Moon, A. 1824. Catalogue of the Indigenous and Exotic Plants Growing in Ceylon. Wesleyan Mission, Colombo.
- Pichon, M. 1950. Classification des Apocynacées: XXV, Echitoidées. Mémoires du Muséum Nationale d'Histoire Naturelle, sér. B, Bot. 1: 1–174.
- Rheede, H. A. 1689. Hortus Indicus Malabaricus 9: 7, t. 5 & 6.



Middleton, David J. 1993. "A New Combination in Chonemorpha (Apocynaceae)." *Novon a journal of botanical nomenclature from the Missouri Botanical Garden* 3, 455–455. <https://doi.org/10.2307/3391393>.

View This Item Online: <https://www.biodiversitylibrary.org/item/14663>

DOI: <https://doi.org/10.2307/3391393>

Permalink: <https://www.biodiversitylibrary.org/partpdf/122243>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.