

THE GRASS FLORA OF COLUMBIA, MISSOURI

LISLE JEFFREY AND FRANCIS DROUET

DURING the spring and summer of 1933 we have spent considerable time collecting and studying the grasses in this region. The following list is presented as the result of the first year's work and should be appended to Rickett's *Flora of Columbia* (Univ. Mo. Stud. 6(1). 1931), together with a few additions made previously in RHODORA 35: 361–362. Reference is made below to the earlier Columbia *Flora* of Daniels (Univ. Mo. Stud. Sci. Ser. 1(2). 1907) and to Bush's revision of the grasses of the Daniels herbarium (Amer. Midl. Nat. 12: 343 ff. 1931). Names of species preceded by an asterisk (*) in the list below are reported as collected for the first time in the flora. We wish to thank Mr. B. F. Bush, Dr. W. E. Maneval, and Dr. J. A. Steyermark for their cooperation during the course of the work.

AGROSTIS STOLONIFERA L. *A. alba* of Amer. Auth. and of the Columbia *Floras*, not L. Including *A. alba* var. *vulgaris* of Daniels, Bush, and Rickett. Most of the specimens can be distinguished as var. MAJOR (Gaud.) Farwell. See Ann. Rep. Nat. Mus. Canada, Dept. Mines Bull. 50 (1926): 122. 1928.

AGROSTIS PERENNANS (Walt.) Tuckerm. Including *A. Schweinitzii* Trin. Several new stations have been found for the typical form.

ARISTIDA BASIRAMEA Engelm. On a freshly cut hillside near Rocky Fork Creek northwest of Columbia, Drouet 967, Aug. 16, 1933. Not Daniels' *A. basiramea*.

*ARISTIDA LONGESPICA Poir. *A. gracilis* Ell. Roadside eight miles north of Columbia, Jeffrey 246, Sept. 6; loess ridges on the Roby Farm south of Rocheport, Jeffrey 253, Sept. 8; along railroad north of Hallsville, Jeffrey 300, Sept. 16; Richland Road, Jeffrey 325, Sept. 27; Highway C near Harrisburg, Jeffrey 331, Sept. 22, 1933. See Contrib. U. S. Nat. Herb. 22(pt. 7): 538, 539. 1924.

*CYNODON DACTYLON (L.) Pers. Established in parkings and waste ground in Columbia, where it has escaped from lawns. One specimen is from West Broadway near East Parkway, Jeffrey 316, Sept. 22, 1933. First observed here by Dr. Steyermark.

ELYMUS CANADENSIS L. *E. robustus* Scribn. & J. G. Sm. The typical form and f. GLAUCIFOLIUS (Muhl.) Fernald (var. *glaucofolius* Gray) have been collected abundantly during the past year. See RHODORA 35: 191. 1933.

ELYMUS VIRGINICUS L. var. TYPICUS (Scribn.) Fernald and the f. HIRSUTIGLUMIS (Scribn.) Fernald (var. *hirsutiglumis* Hitchc.). Both forms are well represented in the collections of the past season. See RHODORA 35: 198.

ELYMUS VIRGINICUS L. var. GLABRIFLORUS (Vasey) Bush. *E. glabriflorus* Scribn. & Ball. *E. glabrifolius* of Rickett's *Flora*. Abun-



Jeffrey, L and Drouet, Francis. 1934. "The grass flora of Columbia, Missouri." *Rhodora* 36, 415–7.

View This Item Online: <https://www.biodiversitylibrary.org/item/14506>

Permalink: <https://www.biodiversitylibrary.org/partpdf/122446>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.