

- chromosomiques nouveaux chez diverses dicotylédones et monocotylédones d'Afrique occidentale. Bull. Jard. Bot. Bruxelles **28**: 315-329.
- MONTGOMERY, F. H. AND S. YANG. 1960. Cytological studies in the genus *Erigeron*. Canad. Jour. Bot. **38**: 381-386.
- SATÓ, D. 1960. The protokaryotype and phylogeny in plants. Sci. Pap. Coll. Gen. Educ., Univ. Tokyo **10**: 303-327.
- SOLBRIG, O. T. 1961. Synopsis of the genus *Xanthocephalum*. Rhodora **63**: 151-164.
- TORRES, A. M. 1961. Hybridization studies in *Zinnia*. Am. Jour. Bot. **48**: 549. (Abstract)
- TURNER, B. L. 1960a. Documented chromosome numbers of plants. Madroño **15**: 219-221.
- _____. 1960b. Meiotic chromosome numbers in Texas species of the genus *Coreopsis* (Compositae-Heliantheae). Southw. Nat. **5**: 12-15.
- _____. AND W. L. ELLISON. 1960. Chromosome numbers in the Compositae. I. Tex. Jour. Sci. **12**: 146-151.
- _____. AND H. S. IRWIN. 1960. Chromosome numbers in the Compositae. II. Meiotic counts for fourteen species of Brazilian Compositae. Rhodora **62**: 122-126.
- _____, J. BEAMAN AND H. F. L. ROCK. 1961a. Chromosome numbers in the Compositae. V. Mexican and Guatemalan species. Rhodora **63**: 121-129.
- _____, W. L. ELLISON AND R. M. KING. 1961b. Chromosome numbers in the Compositae. IV. North American species, with phyletic interpretations. Am. Jour. Bot. **48**: 216-223.
- _____. and M. C. JOHNSTON. 1961. Chromosome numbers in the Compositae. III. Certain Mexican species. Brittonia **13**: 64-69.

CHROMOSOME NUMBERS IN MEXICAN AND GUATEMALAN COMPOSITAE¹

J. H. BEAMAN AND B. L. TURNER

The chromosome numbers reported here were obtained from bud material collected by Beaman in the summer of 1960. The counts were made by Turner (except in *Seigesbeckia* which Beaman examined) from pollen mother cell squashes as outlined by Turner and Ellison (1960). The voucher specimens were determined by Beaman, except

¹Supported by National Science Foundation grants G-9025 and G-9045.

Helianthella quinquenervis which was determined by Dr. C. B. Heiser, Jr., and they are deposited in the Michigan State University Herbarium. A nearly complete set is also in the Herbarium of the University of Texas. All material studied is listed in Table 1.

DISCUSSION

EUPATORIEAE—*Piqueria pilosa* ($n=12$). Turner, Powell, and King (1962) reported two other collections of this species as $n=12$.

ASTEREAE—*Erigeron coronarius* ($n=18$). Diploid counts of $n=9$ were obtained by De Jong and Longpre (unpublished) in two Durango collections of this species. *E. coronarius* as presently understood is polymorphic, and further study may indicate that the variations are correlated with chromosomal races.

Xanthocephalum linearifolium ($n=4$). The count in this collection was noted by Solbrig (1961) who reported counts for three species of *Xanthocephalum*.

HELIANTHEAE—*Bidens triplinervia* var. *macrantha* ($n=12$). Tetraploid and hexaploid populations of this widely distributed and variable taxon occur on the Guatemalan volcanoes of Pleistocene or recent age (Beaman, De Jong, and Stoutamire, 1962). This diploid material comes from the older, non-volcanic Sierra de los Cuchumatanes.

Jaegeria petiolaris ($n=9$). Turner, Powell and King (1962) have found two other species (*J. pedunculata* Hook. & Arn. and *J. hirta* Less.) with $n=18$. The present report establishes the tetraploid nature of the other counts.

Calea sp. ? ($n=16$). Turner et al. (1961 (1962) have reported approximate counts of $n=16$, 17, and 18 for several other species, and they have obtained a definitive count for only one species, *C. trichotoma* with $n=18$. The generic determination of our material may possibly be in error; if it belongs to the genus *Calea*, it is apparently undescribed.

Siegesbeckia nudicaulis ($n=15$), *S. repens* ($n=15$), *S. triangularis* ($n=15$). The first two of these species are very restricted endemics, while the latter is more widespread and may be conspecific with *S. orientalis* (*sens. lat.*) in which Diers (1961) found $n=15$.

HELENIEAE—*Microspermum debile* ($n=12$). This is the first count reported in this small and little collected genus.

SENECIONEAE—*Werneria nubigena* ($n=50$). Counts in 13 species of *Werneria*, including a count of $2n=212 \pm 8$ in *W. nubigena*, were reported by Diers (1961). All but one species he investigated had somatic numbers of 100 or higher.

SUMMARY


Meiotic chromosome counts are reported for 25 species of

Mexican and Guatemalan Compositae. These include the first counts published for 16 taxa. The first report is given for the genus *Microspermum* ($n=12$). A count of $n=9$ was found for *Jaegeria petiolaris*, thus establishing the tetraploid nature of previous chromosome reports of $n=18$ for this genus. Numbers consistent with established basic numbers or previously reported counts were found in *Piqueria*, *Erigeron*, *Grindelia*, *Heterotheca*, *Xanthocephalum*, *Helianthella*, *Sanvitalia*, *Siegesbeckia*, *Viguiera*, *Florestina*, *Helenium*, *Cacalia*, *Senecio*, *Cirsium*, and *Hieracium*. Numbers differing with previously reported counts or basic numbers were obtained in *Bidens triplinervia* var. *macrantha* ($n=12$), *Calea* sp. ? ($n=16$), and *Werneria nubigena* ($n=50$).

— DEPARTMENT OF BOTANY AND PLANT PATHOLOGY, MICHIGAN STATE UNIVERSITY, EAST LANSING; PLANT RESEARCH INSTITUTE, THE UNIVERSITY OF TEXAS, AUSTIN.




1



2



3



4

Figs. 1-4. Meiotic chromosomes of selected species, approximately $\times 2000$. Fig. 1. *Jaegeria petiolaris* ($n=9$). Fig. 2. *Calea* sp. ? ($n=16$). Fig. 3. *Microspermum debile* ($n=12$). Fig. 4. *Werneria nubigena* ($n=50$).

LITERATURE CITED

- BEAMAN, J. H., D. C. D. DE JONG, AND W. P. STOUTAMIRE. 1962. Chromosome studies in the alpine and subalpine floras of Mexico and Guatemala. Amer. Jour. Bot. 49: 41-50.
- DIERS, L. 1961. Der Anteil an Polyploiden in den Vegetationsgürteln der Westkordillere Perus. Zeitsch. für Botanik 49: 437-488.
- SOLBRIG, O. T. 1961. Synopsis of the genus *Xanthocephalum* (Compositae). Rhodora 63: 151-164.
- TURNER, B. L., AND W. L. ELLISON. 1960. Chromosome numbers in the Compositae. I. Meiotic chromosome counts for 25 species of Texas Compositae including 6 new generic reports. Texas Jour. Sci 12: 146-151.
- TURNER, B. L., W. L. ELLISON, AND R. M. KING. 1961. Chromosome numbers in the Compositae. IV. North American species, with phyletic interpretations. Amer. Jour. Bot. 48: 216-223.
- TURNER, B. L., AND M. C. JOHNSTON. 1961. Chromosome numbers in the Compositae—III. Certain Mexican species. Brittonia 13: 64-69.
- TURNER, B. L., M. POWELL, AND R. M. KING. 1962. In press. Chromosome numbers in the Compositae. VI. Additional Mexican and Guatemalan species. Rhodora 64: 251-271.

TABLE 1. Summary of collections studied.

Taxon	Locality	n	chromosome number
EUPATORIEAE			
<i>Piqueria pilosa</i> H.B.K.	State of Mexico. Nevado de Tolhuca, northwest side of mtn., ca. 3715 m. alt. Beaman 3471.	12	
ASTERAEAE			
<i>Erigeron coronarius</i> Greene	State of Mexico. 5 kilometers north of Atlacomulco, ca. 2500 m. alt. Beaman 3370.	18	
<i>E. scaposus</i> DC.	State of Mexico. 6 kilometers north of Atlacomulco; ca. 2500 m. alt. Beaman 3358a.	18	
<i>Grindelia inuloides</i> Willd.	Nuevo Leon. Cerro Potosí, summit of mtn., ca. 3650 m. alt. Beaman 3346.	6	
<i>Heterotheca inuloides</i> Cass. var. <i>inuloides</i>	State of Mexico. 3 kilometers southeast of Amecameca; ca. 2500 m. alt. Beaman 3474.	9	
<i>Xanthocephalum linearifolium</i> (DC.) Greenman	Distrito Federal, Mexico. At La Cima Station between Mexico and Cuernavaca, 3035 m. alt. Beaman 3653.	4	

HELIANTHEAE

<i>Bidens triplinervia</i> var. <i>macrantha</i> (Wedd.) Sherff	Guatemala. Dept. of Huehuetenango. Between Tojiah and Chemal at Km. 319.5 on Ruta Nacional 9 N, ca. 3380 m. alt. <i>Beaman</i> 3743.	12
<i>Helianthella quinquenervis</i> (Hook.) A. Gray	Nuevo Leon. Cerro Potosí, near summit of mtn. on northeast side, ca. 3600 m. alt. <i>Beaman</i> 3339.	15
<i>Jaegeria petiolaris</i> Robins.	Michoacan. In large llano ca. 4 kms. southeast of Cerro San Andres, ca. 10 kms. (straight line distance) north of Ciudad Hidalgo, ca. 2930 m. alt. <i>Beaman</i> 4245.	9 (Fig. 1)
<i>Calea</i> sp. ?	Guatemala. Dept. of Huehuetenango. Between Tojiah and San Juan Ixcoy at Km. 323.5 on Ruta Nacional 9 N, ca. 3200 m. alt. <i>Beaman</i> 3956.	16 (Fig. 2)
<i>Sanvitalia procumbens</i> Lam.	Puebla. 1.5 miles west of Chachapa, ca. 2300 m. alt. <i>Beaman</i> 3615.	8
<i>Siegesbeckia nudicaulis</i> Standl. & Steyermark	Guatemala. Dept. of Huehuetenango, between Tojiah and San Juan Ixcoy at Km. 324.5 on Ruta Nacional 9 N, ca. 3140 m. alt. <i>Beaman</i> 3844.	15
<i>S. repens</i> Robins. & Greenm.	Oaxaca. Llano de las Flores, on the Oaxaca-Valle Nacional highway 20 kilometers east of Ixtlan, ca. 2870 m. alt. <i>Beaman</i> 3699.	15
<i>S. triangularis</i> Cav.	Guatemala. Dept. of Huehuetenango, between Tojiah and San Juan Ixcoy at Km. 323.5 on Ruta Nacional 9 N, ca. 3200 m. alt. <i>Beaman</i> 3849.	15
<i>Viguiera hemsleyana</i> Blake	Oaxaca. On the Oaxaca-Valle Nacional highway, 8 miles east of Ixtlan, ca. 3600 m. alt. <i>Beaman</i> 3662.	ca 34

HELENIEAE

<i>Florestina pedata</i> (Cav.) Cass.	Puebla. 1.5 miles west of Chachapa, ca. 2300 m. alt. <i>Beaman</i> 3616.	10
<i>Helenium hoopesii</i> A. Gray	Nuevo Leon. Cerro Potosí, sum-	15

Microspermum debile
Benth.

mit of mtn., ca. 3650 m. alt. Beaman 3347.

Oaxaca. On the Oaxaca-Valle 12 (Fig. 3) Nacional highway, on Cerro Pelon, 25.5 miles east of Ixtlan, ca. 2950 m. alt. Beaman 3663.

SENECIONEAE

Cacalia peltata H.B.K.

Puebla. 5.3 miles southwest of 30 San Salvador el Seco; ca. 2500 m. alt. Beaman 3619.

Senecio conzattii Greenm.

Oaxaca. Llano de las Flores, on 20 the Oaxaca-Valle Nacional highway 20 kms. east of Ixtlan, ca. 2870 m. alt. Beaman 3700.

S. iodanthus Greenm.

Michoacan. Summit of Cerro San 20 Andres, ca. 12 kms. (straight line distance) north of Ciudad Hidalgo, 3589 m. alt. Beaman 4241.

Werneria nubigena H.B.K.

Guatemala. Dept. of Huehuetenango. Between Tojiah and Chemal at Km. 319.5 on Ruta Nacional 9 N, ca. 3380 m. alt. Beaman 3741.

CYNAREAE

Cirsium skutchii Blake

Guatemala. Dept. of Huehuetenango. Between Tojiah and Chemal at Km. 319.5 on Ruta Nacional 9 N, ca. 3380 m. alt. Beaman 3742.

CICHORIEAE

Hieracium mexicanum Less. State of Mexico. Iztaccihuatl, 9 south side of mtn. at La Joya, ca. 3990 m. alt. Beaman 3501.

H. selerianum Zahn

Guatemala. Dept. of Huehuetenango. Between Tojiah and Chemal at Km. 317.5 on Ruta Nacional 9 N, ca. 3400 m. alt. Beaman 3802.



BHL

Biodiversity Heritage Library

Beaman, John H. and Turner, B. L. 1962. "Chromosome numbers in Mexican and Guatemalan Compositae." *Rhodora* 64, 271–276.

View This Item Online: <https://www.biodiversitylibrary.org/item/14535>

Permalink: <https://www.biodiversitylibrary.org/partpdf/122816>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.