

## RANGE EXTENSIONS OF VASCULAR PLANTS FROM THE SEWARD PENINSULA, NORTHWEST ALASKA

BY SYLVIA KELSO<sup>1</sup>

As a barrier, filter, or avenue of migration, the concept of a North American-Asiatic land connection linking Siberia to Alaska through the Bering and Chukchi seas has been the subject of much interest since it was first speculated upon by Wallace in 1876. Enough data have accumulated now from the disciplines of biology, geology, and anthropology, among others, to accept unequivocally that such a connection did exist through not only one but several periods in the past, the most recent ending some 12,000 years ago (c.f. Hopkins, 1979). The vast area under the influence of the land bridge, known as Beringia, stretches from the Kolyma River in Siberia to the MacKenzie Delta in Canada (Yurtsev, 1972). Today, many biotic distribution patterns in this region have their origins in aspects of paleogeography (c.f. Kurten, 1966; Krassilov, 1976; Sher, 1976). As more becomes known about contemporary patterns of biotic distribution, more can be explained about the geography of times past. Concurrently, as more is learned about historical changes, more can be understood about the patterns of today.

The location of Alaska's Seward Peninsula in the heart of the North American half of Beringia (Figure 1) has long stirred the interest of phytogeographers in the floristics of that region. From the early 19th century and the work of the botanist Chamisso who voyaged through the Bering Strait with the explorer Otto von Kotzebue, to the present, numerous collections have been made along the coast of the peninsula (c.f. Flett, 1901; Porsild, 1938; Hultén, 1968; Young, 1971; Racine, 1979) that was easily accessible by sea or more recently, by air. Unfortunately, however, due to its size and general inaccessibility from lack of a road system, the interior of the Seward Peninsula, which may be a critical area for our understanding of the American sector of Beringia, was neglected and remained generally blank except for sporadic collections from a few isolated points. Road systems now extend from Nome to the northwest, north, and east. This has made much of the interior more

---

<sup>1</sup>Present address: University of Alaska Museum  
Fairbanks, Alaska 99701




Figure 1. Seward Peninsula, Alaska.



accessible and blank areas can begin to be filled. During the field seasons of 1981 and 1982, the author had the opportunity to make collections in previously unexplored portions of the Seward Peninsula. This paper notes important floristic discoveries for the region.

The Seward Peninsula is generally mountainous and includes the York, Kigluaik, Darby, and Bendeleben Ranges (Figure 1). These offer rugged alpine terrain in contrast to the rolling lowlands and the opportunity for a strong connection to the mountain chains of Northeast Asia and to the Brooks and Alaska Ranges in Alaska. Much of the area was subject only to local alpine glaciation or none at all during the Pleistocene (Sainsbury, 1967). The Kigluaik Mts. have only recently emerged from glaciation and contain a relict glacier in the vicinity of Mt. Osborne (A.B. Till, U.S. Geologic Survey, pers. comm.). In contrast to a periglacial environment, anomalously warm habitats created by geothermal activity can be seen at Serpentine and Pilgrim Hot Springs.

Botanical exploration of the interior of the Seward Peninsula is only just now beginning. As the alpine regions become better known, puzzling floristic gaps may well be filled. The following taxa have recently been discovered on the Seward Peninsula and constitute noteworthy range extensions for the area (c.f. Hultén, 1968). Specimens cited here are deposited at the University of Alaska Museum Herbarium, Fairbanks (ALA).

#### GRAMINEAE

##### ***Alopecurus aequalis* Sobol.**

BENDELEBEN QUAD.<sup>1</sup>: Mile 60, Kougarak Rd, Central Seward Peninsula. In the margins of an ephemeral pond, 70 msm.  
*Kelso 82-219* 15 August 1982

This is a circumpolar taxon that was believed to have a notable gap in its distribution between the Chukchi and Seward Peninsulas (Figure 2). Kozhevnikov (1981) and Yurtsev et al. (1979) have now reported it from the interior of the Chukchi Peninsula and this discovery on the Alaska side of the Bering Strait shows the gap does not exist.

---

<sup>1</sup>Note: The term "Quad" refers to the U.S. Geologic Survey topographic map at the scale of 1:250,000. These are used in lieu of county divisions which do not exist in Alaska.




Figure 2. Ranges of *Alopecurus aequalis* (solid line, squares) and *Agrostis scabra* (dotted line, circles)




Figure 3. Ranges of *Carex bicolor* (solid line, squares) and *Crepis nana* (dotted line, circles)



***Agrostis scabra* Willd.**

BENDELEBEN QUAD.: Pilgrim Hot Springs, Central Seward Peninsula, 70 msm. *Kelso* 82-133 20 July 1982

This is a boreal taxon common in open or disturbed areas throughout the coniferous forest region of North America east to Newfoundland, with isolated occurrences in Chukotka (Figure 2). This location at Pilgrim Hot Springs (c.f. *C.H. Racine* #439 from Serpentine Hot Springs at ALA) extends the range northwestward.

## CYPERACEAE

***Carex bicolor* All.**

TELLER QUAD.: Cape Prince of Wales. Shore of Lopp Lagoon, in sandy gravel near the reindeer corral, 0 msm. *Kelso* 82-151 26 July 1982

*Carex bicolor* is an arctic-alpine taxon found in Alaska at higher elevations in the Brooks and Alaska Ranges with outliers in Chukotka (Figure 3). This discovery at Cape Prince of Wales is the first on the Seward Peninsula and extends the American range considerably to the westward. The sea level habitat exemplifies the strong alpine connections common on the Seward Peninsula where habitat and severe climatic conditions create alpine-type environments at very low elevations.

## COMPOSITAE

***Crepis nana* Richards.**

BENDELEBEN QUAD.: Kougarok Rd., Mile 59, in riparian gravel of the Pilgrim River, 70 msm. *Kelso* 82-201 12 August 1982

*Crepis nana* is an arctic-alpine plant known from Asia and North America. It has a wide range from Labrador to Central Siberia, but the distribution has been mapped as sporadic. On the Seward Peninsula it is known from Port Clarence on the coast and this occurrence in the interior is not surprising (Figure 3).

## PRIMULACEAE

***Androsace alaskana* Cov. and Standl.**

BENDELEBEN QUAD.: Kougarok Rd, Mile 51. On limestone outcrop, in sandy crevices, 75 msm. *Kelso* 82-223 17 August 1982

This is a North American taxon of relatively narrow distribution throughout the Alaska Peninsula to the Wrangell-St. Elias Moun-




Figure 4. Ranges of *Aphragmus eschscholtzianus* (solid line, squares) and *Androsace alaskana* (dotted line, circles)

tains. Its presence on the Seward Peninsula, widely disjunct from the rest of the range, is quite remarkable (Figure 4).

#### CRUCIFERAE

##### ***Aphragmus eschscholtzianus* Andrz.**

TELLER QUAD.: Cape Prince of Wales. On wet solifluction soil, steep north-facing limestone slope, 80 msm. *Kelso* 81-352 27 July 1981

*Aphragmus eschscholtzianus* has a very similar distribution pattern to that of *Androsace alaskana*, reaching from the Aleutians into the Wrangell-St. Elias region. Similarly, this discovery is a disjunct outlier on the Seward Peninsula; the plant has also been found once in the Kigluaik Mts. and is now known from the western tip of the peninsula as well (Figure 4).

#### REFERENCES

- FLETT, J. B. 1901. Notes on the flora around Nome City. *Plant World* (4): 67-68.
- HOPKINS, D. M. 1979. Landscape and climate of Beringia during late Pleistocene and Holocene times. pp. 15-41. *In*: W. S. Laughlin and A. B. Harper (eds.) *The First Americans: Origins, Affinities and Adaptations*. Gustav Fischer, N.Y.
- HULTÉN, E. 1968. *Flora of Alaska and Neighboring Territories*. Stanford University Press, Stanford, California.
- KOZHEVNIKOV, Y. 1981. *Catalogus plantarum vascularum terrae tschuktschorum*. *Novosti Systematiki Vysshikh Rastenii* 18: 229-248.
- KRASSILOV, V. A. 1976. The role of Bering land connections in the formation of the Cenozoic floras of east Asia and North America. *In*: *Beringia in Cenozoic*. "Report of the All-Union Symposium on the Bering Land Bridge, Khabarovsk, 1973. pp. 129-134.
- KURTEN, B. 1966. Pleistocene mammals and the Bering Bridge. *Comm. Biol. Soc. Fennica* 29: 1-7.
- PORSILD, A. E. 1938. Flora of Little Diomed Island in Bering Strait. *Transactions of the Royal Society of Canada ser. 3, sect. 5*(32): 21-38.
- RACINE, C. H. 1979. Floristic and vegetational studies. *In*: H. R. Melchior (ed.) *Biological Survey of the Bering Land Bridge National Monument*. Revised final report to the National Park Service, Alaska Cooperative Park Studies, University of Alaska, Fairbanks.
- SAINSBURY, C. L. 1967. Upper Pleistocene features in the Bering Strait area. U.S. Geologic Survey Professional Paper, 575-D: 203-213.
- SHER, A. V. 1976. The role of Beringian land in the development of holarctic mammalian faunas in the late Cenozoic. pp. 227-241. *In*: "Beringia in Cenozoic." Report of the All-Union Symposium on the Bering Land Bridge, Khabarovsk, 1973.



- WALLACE, H. R. 1876. The geographical distribution of animals. Harper, N.Y.
- YOUNG, S. B. 1971. The vascular flora of St. Lawrence Island with special reference to floristic zonation in the arctic region. *Contr. Gray Herb.* **201**: 11-115.
- YURTSEV, B. A. 1972. Phytogeography of northeastern Asia and the problem of transsberingian floristic interrelations. pp. 19-54. *In*: A. Graham (ed.) *Floristics and paleofloristics of Asia and eastern North America*. Elsevier Publishing Co., Amsterdam.
- YURTSEV, B. A., V. V. PETROVSKY, A. A. KOROBKOV, T. M. KOROLEVE, & V. Y. KAAZHIVIN. 1979. A review of the geographic distribution of vascular plants of the Chukotka tundra. *Bull. Mosk. O-Va Ispyt. Prirody. Otd. Biol.* **84**(5): 111-120; **84**(6): 74-83.

DEPT. OF ENVIRONMENTAL, POPULATION AND ORGANISMIC BIOLOGY  
UNIVERSITY OF COLORADO, BOULDER, COLORADO

AND

UNIVERSITY OF COLORADO MUSEUM HERBARIUM  
BOULDER, COLORADO 80309



Kelso, Sylvia. 1983. "RANGE EXTENSIONS OF VASCULAR PLANTS FROM THE SEWARD PENINSULA, NORTHWEST ALASKA." *Rhodora* 85, 371–379.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/24161>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/123183>

**Holding Institution**

Missouri Botanical Garden, Peter H. Raven Library

**Sponsored by**

Missouri Botanical Garden

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.