

CURLYGRASS IN NEW YORK STATE. — During a recent collecting trip through the eastern tip of Long Island, my son, Andrew R. Moldenke, called me to a small cranberry bog near Montauk, Suffolk County, to show me a large stand of curly-grass (*Schizaea pusilla*). He collected material of it under his collection no. 38, dated July 23, 1960.

Fernald, in the 8th edition of "Gray's Manual of Botany" (1950), gives the distribution of this plant as "Nfld. and N. S.; Pine Barrens of N. J.". Gleason, in the "New Britton and Brown's Illustrated Flora" (1952 & 1958), records it from Newfoundland, Nova Scotia, Goose County, Ontario, and New Jersey.

It would appear, therefore, that this New York station is worthy of note. The plant was growing in association with cranberry, three species of sundew (*Drosera intermedia*, *D. rotundifolia*, and *D. filiformis*), and marsh St. Johnswort (*Triadenum virginicum*). — HAROLD N. MOLDENKE.



Moldenke, Harold N. 1960. "Curlygrass in New York State." *Rhodora* 62, 294–294.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/14533>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/124232>

**Holding Institution**

Missouri Botanical Garden, Peter H. Raven Library

**Sponsored by**

Missouri Botanical Garden

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.