Gyldenstolpe, N. 1924. Zoological results of the Swedish expedition to central Africa 1921. Vertebrata. I. Birds. Kungl. Svenska Vetenskapsakad. Handl. (3), vol. 1, No. 3.

Jackson, Sir F. 1938. The birds of Kenya Colony and the Uganda Protectorate. Vol. 3. London: Gurney and Jackson.

Neumann, O. 1908. [*Estrilda atricapilla graueri*, subsp. n.]. *Bull. Erit. Orn. Cl.* 21: 55. Prigogine, A. In press. Les populations de *Estrilda atricapilla* (Verreaux) de l'Afrique

Prigogine, A. In press. Les populations de *Estrilda atricapilla* (Verreaux) de l'Afrique centrale et description d'une nouvelle race. *Rev. Zool. afr.*

Reichenow, A. 1902. Neue afrikanische Arten. Orn. Monatsber. 10: 184-185.

— 1904/5. Die Vogel Afrikas. Vol. 3. Neudamm: Neumann.

— 1912. Die Vogelfauna des Mittelafrikanischen Seengebietes. Wiss. Ergebn. Deutsch. Zentral-Afrika-Exped. 1907–1908. Vol. 3. Leipzig.

Sclater, W. L. 1930. Systema avium Aethiopicarum. Part 2. London.

White, C. M. N. 1963. A revised check list of African flycatchers, etc. Lusaka: Govt. Printer.

New Subspecies of Caprimulgus sericocaudatus from the Amazon River Basin

by Robert W. Dickerman

Received 13th November 1974

The Silky-tailed Nightjar *Caprimulgus sericocaudatus* is an exceedingly rare species in North American ornithological collections. It is represented by only six specimens from Misiones Province of Argentina; two from south-eastern Brazil (Sao Paulo and Curitiba); and the two specimens, on which the species was described, from the Rivioli (Massena) Collection, labeled only South America. De Schauensee (1970) also included Paraguay (Capitan Meza, alto Parana) in the range of the species, but I have not been able to locate (nor could he provide me with) the reference to this locality. The population of the Amazon basin is represented by a specimen from Santarem, Brazil and one from Yarinacocha in Amazonian Peru.

Recently Dr. Hannalore E. Hinsch made a collection of birds at the Instituto Veterinario de Investigaciones Tropicales y de Alturas 59 km east of Pucallpa, Loreto, Peru, while carrying out a survey for arthropod-borne viruses. The birds were shipped frozen to New York where they were prepared and identified by the author to provide voucher specimens for study. Included was a single specimen of this species.

Upon comparison of this individual with the three specimens available in the American Museum of Natural History, differences were noted and thus the above enumerated material was assembled (except for the two specimens of the type series).

The characters noted proved consistent, and comparisons of representative birds with the type series in the Academy of Natural Sciences of Philadelphia immediately indicated that the name *sericocaudatus* was not based upon birds from the Amazon basin, and that population awaited description. However, first it was necessary to restrict the type locality of the nominate form. Stone (1899) compiled a list of the types in the Academy and cited the bird upon which the description was based (No. 21,905) as the type and it bears a type label although it was not so designated in the original description. Cory (1919) gave the range as "southeastern Brazil, Bahia, southwards". Mr. R. M. de Schaeunsee (in conversation) informed me that most early collections from southeastern Brazil were exported from Bahia or Rio de Janiero. The species was not recorded from the state of Bahia by Olivero M de O. Pinto (1935). As first revisor of the species, I suggest restricting the type locality of *Caprimulgus (Antrostomus) sericocaudatus* (Cassin) 1849 to the vicinity of Rio de Janiero, Brazil, that being a probably source of the type material closest to the range of the species as indicated by presently available specimens.

It should be noted that Cory, and de Schauensee (*loc. cit*) were in error in citing the publication date for the species as 1848. The original description bears the date 30 October 1849 and was correctly cited by Stone (*loc. cit.*) and Peters (1940).

The population of the Amazon River Basin may now be described as:

Caprimulgus sericocaudatus mengeli subsp. nov.

Diagnosis: Smaller (see Table) and darker, more sooty, than the nominate form.

	TABLE	
Measurements	of Caprimulgus	sericocaudatus
	and a start and a second and a	

	wing		tail	
	males	females	males	females
sericocaudatus	177-182	167-184	144-148	136-147
	179.4	175.8	146.4	141.4
	(n=5)	(n=5)	(n=5)	(n=5)
mengeli	157 & 163	156	124 & 127	116*
Tail apparently retaine	d from first basic	olumage.		

Type: American Museum of Natural History, no. 811079, adult male, collected by Hannalore E. Hinsch, on 15th November 1972 at Instituto Veterinario de Investigaciones Tropicales y de Alturas, 59 km east of Pucallpa, Loreto Department, Peru. Original field number HEH-A-506.

Range: Amazon River basin.

Material: Besides the type, a 3 from Yarinacocha, Peru, and a 9 from Santarem, Brazil.

Name: With great satisfaction I name this form for Dr. Robert M. Mengel, student of Caprimulgids and friend.

Notes: De Schauensee (1970) has described the female plumage of the species. The "young" bird mentioned in the original description is in reality another adult male with a slight difference in wing formula. The only female of *mengeli* is the Santerem specimen, which appears to be an immature. Its rectrices are narrower and more rounded and are only diffusely tipped with buff as compared to those of the other available apparently adult females.

ACKNOWLEDGEMENTS

I am grateful to the curators of the Los Angeles County Museum, Carnegie Museum, Field Museum of Natural History and the Academy of Natural Sciences of Philadelphia for loaning material or for allowing me to examine specimens in their care.

References:

Cory, C. B. 1919. Catalogue of birds of the Americas, Part II, No. 2. Field Mus. Nat. Hist. Pub. 197, Zool. Series XIII.

De Schauensee, R. M. 1970. A guide to the birds of South America. Edinburgh: Oliver & Boyd. Pinto, Oliver M. de O. 1935. Aves. da Bahia. Rev. Mus. Paul. (XIX), 19: 1-326.

- Peters, James L. 1940. Check-list of birds of the world 4: 1-291. Cambridge, Mass.: Harvard Univ. Press.
- Stone, W. 1899. A study of the type specimens of birds in the collection of the Academy of Natural Sciences of Philadelphia, with a brief history of the collection. *Proc. Acad. Nat. Sci. Phila.* 51:5 91.

<sup>Cassin, J. 1849. Descriptions of new species of the family caprimulgidae, specimens of which are in the collections of the Academy of Natural Sciences of Philadelphia. Proc. Acad. Nat. Sci. Phila. 4: 236-239.
Cory, C. B. 1919. Catalogue of birds of the Americas, Part II, No. 2. Field Mus. Nat. Hist. Pub.</sup>



Dickerman, Robert William. 1975. "New subspecies of Caprimulgus sericocaudatus from the Amazon River Basin." *Bulletin of the British Ornithologists' Club* 95, 18–19.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/126880</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/144044</u>

Holding Institution Natural History Museum Library, London

Sponsored by Natural History Museum Library, London

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: British Ornithologists' Club License: <u>http://creativecommons.org/licenses/by-nc-sa/4.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.