## Some Revision in Antechinus (Marsupialia)-2

By N. A. WAKEFIELD\* and R. M. WARNEKE\*\*

## SUMMARY

Antechinus stuartii is recognized as a species distinct from A. flavipes, and the taxonomy of the two is reviewed. It is shown that, in recent literature, the name A. flavipes flavipes has been misapplied to A. stuartii. A neotype is designated for A. stuartii.

Plastic and cranial characters of major populations of these species and of *A. godmani* are examined and compared statistically. The distribution of each is mapped.

A. flavipes flavipes, of temperate eastern Australia, and A. f. leucogaster, of south-western Australia, are animals of open dry sclerophyll forests, but two populations of the species in tropical Queensland are restricted to rainforest habitats.

A. stuartii stuartii occurs in dense dry sclerophyll and wet sclerophyll forests of the Eastern Highlands and coastal areas of south-eastern Australia. This habitat is essentially adjacent to that of A. flavipes flavipes, but the two species are sympatric in some places. The tropical race "adustus" belongs, not to A. flavipes but to A. stuartii.

An appendix shows a revised distribution of Antechinus minimus maritimus. Taxonomy is revised as follows:

#### ANTECHINUS FLAVIPES

#### **Antechinus flavipes flavipes**

- (*Phascogale flavipes* Waterhouse, 1837; north of Hunter River, New South Wales.)
- South-eastern Queensland, New South Wales, Victoria, south-eastern South Australia.
- (Subjective synonym: *Phascogale rufo*gaster Gray, 1841; South Australia.)

#### Antechinus flavipes leucogaster

- (*Phascogale leucogaster* Gray, 1841; Canning River, Western Australia.) South-western Western Australia.
- Antechinus flavipes, populations of uncertain taxonomy.

North-east Queensland; Cape York Peninsula.

March, 1967

## ANTECHINUS STUARTII

#### Antechinus stuartii stuartii

- (Antechinus stuartii Macleay, 1841; Manly, New South Wales. Neotype: Waterfall, National Park, New South Wales.)
- South-eastern Queensland, New South Wales, Victoria.
- (Subjective synonyms: Antechinus unicolor Gould, 1854; Sydney, New South Wales. Antechinus flavipes burrelli Le Souef and Burrell, 1926; Ebor, north-eastern New South Wales.)

#### Antechinus stuartii adustus

(*Phascogale flavipes adusta* Thomas, 1923; Ravenshoe, north-eastern Queensland.)

North-eastern Queensland.

## I. MATERIALS AND METHODS

#### (a) Sources

The sources of specimens and data are as follows:

American Museum of Natural History. (AMNH)

Australian Museum Sydney. (AM)

- British Museum (Natural History). (BM)
- Department of Zoology and Comparative Physiology, Monash University.
- Division of Wildlife Research, C.S.I.R.O., Canberra.
- Fisheries and Wildlife Department, Melbourne. (FWD)
- Museum of Comparative Zoology, Harvard.

<sup>\*</sup> Department of Zoology and Comparative Physiology, Monash University, Clayton, Victoria.

toria. \*\* Fisheries and Wildlife Department, Victoria.

- National Museum of Victoria. (NMV)
- Queensland Institute of Medical Research.
- Queensland Museum, Brisbane.

Smithsonian Institution, United States National Museum.

South Australian Museum, Adelaide. Western Australian Museum, Perth.

Although Antechinus flavipes and A. stuartii are widely distributed in eastern Australia and are abundant in areas of favourable habitat, both species are poorly represented in most museum collections. Of a total of about 250 specimens of A. flavipes and 400 specimens of A. stuartii examined, almost half are in the Fisheries and Wildlife Department collection.

In the course of the study, a programme of field observation and trapping was carried out, and, unless otherwise acknowledged, the details of habitat, set out in Sections III and VIII, are based on observations made by us in the course of that field work.

The other species dealt with in this paper, *A. godmani*, is known only from a small area in Queensland, and of the 37 specimens examined by us only 4 were available in Australian museums.

## (b) Characters examined and techniques of measurement

The morphological features investigated and the techniques of measurement used were discussed in the previous paper of this series (Wakefield and Warneke, 1963). They are, in brief, as follows:

## Characters examined

Size of body and appendages.

Size and morphology of the skull and teeth.

Characters of the pelage.

Characters of the feet.

External changes associated with breeding.

## Techniques of measurement

Measurements of the body were taken using a measuring board, vernier calipers and a steel tape. The tail was measured from the cloaca; the pes from the heel to the tip of the longest toe, excluding the claw; and the ear from the tragoid notch to the tip.

While these techniques were used for all specimens in the Fisheries and Wildlife Department collection, they may not have been precisely the same as those used for measurements recorded with specimens from other institutions.

Figure 1 shows the seven measurements taken of the skull and dentition. The instruments used were either a HELIOS dial-reading or vernier calipers calibrated to 0.1 mm., and the measurements were taken with the aid of a binocular microscope at 6 to 10 magnifications. All cranial measurements recorded in this paper (except Thomas's of A. godmani) were made by this technique, and are recorded to the nearest 0.05 mm.

Specimens with milk teeth (deciduous fourth premolars) or in which the subsequent permanent premolars were not fully erupted were excluded from the tables of measurements.

### (c) Statistical analysis

The "t" test, as outlined in Karmel (1959), was used for the comparisons, and the values of P were obtained from the statistical tables of Fisher and Yates (1957). Results are expressed in terms of the highest probability level exceeded (e.g. P < 0.02) of the range of probability levels 0.10, 0.05, 0.02, 0.01 and 0.001.

## II. TAXONOMY OF ANTECHINUS FLAVIPES

The species was described by Waterhouse (1838) as *Phascogale flavipes*,

Vict. Nat.—Vol. 84







and the type, from "north of Hunter River, N.S.W.", is in the BM., No. 55.12.24.75 (skin and skull).

In recent literature—for example, Iredale and Troughton (1934) and Tate (1947: 127)—what we now regard as Antechinus stuartii (Sections VII-XI) has been identified as A. flavipes flavipes. But measurements and other details of the type specimen (communicated to us by R. E. Hill\*), and Waterhouse's original description, identify the nominate form of A. flavipes according to our treatment of it in Sections III-VI.

Gray (1841) described a specimen from "Adelaide, South Australia" (BM. No. 41.1251; Skin and skull) as *Phascogale rufogaster*. Based on this, the trinomial *Antechinus flavipes rufogaster* has been used (Iredale and Troughton, *loc. cit.*; *et al.*) for a South Australian population of *A*. *flavipes*.

Gould (1863) did not confuse A. flavipes with A. stuartii. Of A. flavipes he stated: "I have observed it to be very abundant both in New South Wales and in South Australia", and remarked that "specimens from both

\* Mammal Department, British Museum (Natural History).

countries presented little or no difference in size or colour". Our studies confirm this observation (Sections IV and V); the epithets *rufogaster* and *flavipes* apply to the one geographical race (Figure 2), and we regard *rufogaster* as a subjective synonym of *A*. *flavipes flavipes*.

Gray (*loc. cit.*) described another specimen, from "Canning River, Western Australia" (BM. No. 41.1244), as *Phascogale leucogaster*. This represents a population that is currently recognized (Iredale and Troughton, *loc. cit.*) as *A. flavipes leucogaster*. It is geographically isolated (Figure 2), and our comparisons (Sections III and IV) indicate that this subspecific status is justified.

While there has been sufficient material available to provide a reasonably clear picture of *A. flavipes* in the temperate region, the relatively few specimens available from northern Australia have allowed no such clarification of the status of its tropical populations. The population of North-east Queensland probably warrants separate subspecific status, and that of Cape York certainly does. Correct taxonomic procedure in the latter case will not be clear until ample comparative material



Fig. 2. Distribution of Antechinus flavipes.

Vict. Nat.—Vol. 84



Fig. 3. Distribution of Antechinus stuartii, and (inset) of A. godmani.

March, 1967

is available of the Arnhem Land animal which Thomas (1904) named *Phascogale bella*.

(The names Antechinus unicolor, A. flavipes burrelli and A. flavipes adustus are excluded from the synonymy of A. flavipes; they belong to that of A. stuartii (Section VII).)

## III. DISTRIBUTION AND HABITAT OF ANTECHINUS FLAVIPES

The main population, which comprises the nominate race, *A.f. flavipes*, is found in south-eastern Queensland, eastern New South Wales, Victoria and south-eastern South Australia (Figure 2). It is mainly on the inland side of the Great Dividing Range but is nearcoastal at both the northern and southwestern extremities of its distribution.

The habitat is fairly open forest, mainly in areas of rainfall between 15 and 25 inches per annum. According to the classification of Wood and Williams (1960), the areas occupied are partly dry sclerophyll forest and partly woodland.

Krefft (1866) reported A. flavipes as abundant in 1857 in the woodlands of Black Box (Eucalyptus bicolor) and Red Gum (E. camaldulensis) of the Murray River flood plains, from Echuca to Mildura. Its distribution along that river evidently linked the Adelaide-Mount Lofty Range population with that of Victoria and New South Wales.

At least in Victoria, most trees with which the species is associated are fibrous-barked eucalypts, either "box" species or "stringy-barks". Food, comprising insects mainly, is obtained from the trunks and large limbs of these trees and about logs. The animals are almost wholly nocturnal and crepuscular, but individuals are occasionally seen abroad in full daylight.

Homes, for daytime refuge, are nor-

mally hollow logs and hollow trunks of trees. Specimens caught at Glenlofty (western Victoria) were living in the trunk of a Red Stringybark (Eucalyptus macrorrhyncha), with Yellow Box (E. melliodora) and Longleaf Box (E. goniocalyx) in association as well as scattered logs and a sparse ground cover of saw-sedge (Gahnia radula) and tussock grass (Poa) (Figure 4). In the lower Glenelg district (south-western Victoria) the habitat was Brown Stringybark (E. baxteri) and Messmate (E. obligua), with a few logs and an abundance of brackers (Pteridium esculentum) and tussock grass. In the Grampians (western Victoria) the species commonly occupies crevices in sandstone cliffs, and along the lower Glenelg River it occasionally inhabits holes in limestone outcrops.

The Western Australian race, A.f. leucogaster, is recorded mainly from forests of Jarrah (Eucalyptus marginata) and Tuart (E. gomphocephala), in the Perth-Blackwood-Albany area.

The habitats of both the North-east Queensland population and that of Cape York Peninsula are tropical rainforests. The former is recorded from Mount Finnegan, Speewah, Russell River and Innisfail; the latter is from McIlwraith Range (Rocky Scrub), Tozer Range, Iron Range and upper Nesbit River. The habitat of each of these localities has been described by Brass, (1953).

## IV. PLASTIC CHARACTERS OF ANTECHINUS FLAVIPES

Both Jones (1923) and Finlayson (1958), using South Australian material and thus avoiding confusion with *A. stuartii*, have given accurate accounts of *A. f. flavipes* (under the synonym, however, of *A. f. rufogaster*). Taken together, their notes provide a very detailed description of the external morphology of the form.

Vict. Nat.-Vol. 84

A. flavipes is a lithe, compactly built animal of shrew-like rather than murine form. The tail is shorter than the head-body length and is not incrassated. The ears, thin in substance, are comparatively large and conspicuous when fully erect but may be folded against the head when the animal is alarmed. The pes, like those of squirrels (Sciuridae), are capable of a considerable degree of rotation, which, together with the prominent and strongly ridged pads on the manus and pes, are features of other members of the Dasyuridae with pronounced scansorial habits.

In the past, much emphasis has been placed on plastic characters in the description of subspecies. The variation observed and measured in each of the geographically isolated populations of *A. flavipes*, as shown in Figure 2, is given below.

#### (a) Body measurements

Details of body measurements, and of numbers of specimens from which these were taken, are set out in Tables 1-4. The figures for *A*. *f*. *flavipes* do not include data of any Queensland specimens, but apply only to material from New South Wales, Victoria and South Australia.

In these tables, the average figures for tail, ear and pes lengths have been expressed as percentages of head-body length. These percentages indicate that body proportions are essentially the same for each of the eastern Australian series, as well as for the two sexes within each series.

For most of the specimens from North-east Queensland and Cape York (i.e. those in the AMNH), the ear had been measured from the crown, instead of the tragoid notch. In several instances both measurements were recorded,



Fig. 4. Habitat of Antechinus flavipes and A. stuartii, Glenlofty, Victoria.

Body measurements of Antechinus flavipes flavipes (south-eastern Australia) Measurements in millimetres

		Range	Mean±S.E.	Standard deviation
Head-body	23 ♂♂ 16 ♀♀	$\begin{array}{rrr} 101 & -135 \\ 94 & -120 \end{array}$	$\frac{116 \cdot 2 \pm 2 \cdot 46}{107 \cdot 9 \pm 2 \cdot 14}$	$ \begin{array}{r} 11 \cdot 80 \\ 8 \cdot 57 \end{array} $
Tail	23 ♂♂ 18 ♀♀	$\begin{array}{rrr} 70 & -112 \\ 78 & -96 \end{array}$	$\begin{array}{c} 95 \cdot 0 \pm 2 \cdot 44 \ *(82 \%) \\ 86 \cdot 9 \pm 1 \cdot 09 \ (81 \%) \end{array}$	$\begin{array}{c} 11\cdot 72\\ 4\cdot 65\end{array}$
Ear	23 ♂♂ 18 ♀♀	$14 \cdot 5 - 21$ 13 - 20	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c}1\cdot 86\\1\cdot 72\end{array}$
Pes	25 ♂♂ 18 ♀♀	$     \begin{array}{r}       18 & -22 \\       17 & -20     \end{array} $	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c} 1 \cdot 04 \\ 1 \cdot 02 \end{array}$

\* Per cent of head-body length.

#### TABLE 2

Body measurements of Antechinus flavipes leucogaster (south-western Australia) Measurements in millimetres

		Range	Mean±S.E.	Standard deviation
Head-body	4 ♂♂ 9 ♀♀	$\begin{array}{rrr} 100 & -108 \\ 86 & -104 \end{array}$	$105 \cdot 3 \pm 1 \cdot 87$ 94 · 9 ± 1 · 77	$\begin{array}{r} 3\cdot 74 \\ 5\cdot 30 \end{array}$
Tail	4 ♂♂ 9 ♀♀	$73 - 94 \\ 65 - 80$	$85 \cdot 3 \pm 4 \cdot 44 * (81\%)$ $73 \cdot 1 \pm 1 \cdot 49 (77\%)$	8·87 4·46
Ear	4 ♂♂ 9 ♀♀	17 - 18 15 - 18	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0·50 1·97
Pes	4 ♂♂ 9 ♀♀	$   \begin{array}{r}     18 & -20 \\     16 & -18   \end{array} $	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0·96 0·67

\* Per cent of head-body length.

and these, together with "notch" measurements for the remainder of the series, indicate that the difference between the two is of the order of 5 mm. For the sake of uniformity "crown" measurements were converted to "notch" by the addition of this length.

Sexual dismorphism of the order of about 8 per cent is apparent in all linear measurements save that of ear length, but sufficient data for statistical evaluation were available only in the case of A. f. flavipes and the Cape York form of the species. In both forms the difference in pes length was statistically significant, with P < 0.001in both cases. In head-body and tail length, Cape York males were significantly larger than females (P < 0.001) but in the tests of A. f. flavipes the highest probability levels exceeded were only 0.02 and 0.01 respectively.

From the tables it can be seen that the Cape York form is approximately

		Range	Mean±S.E.	Standard deviation	
Head-body	5 33 8 99	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\frac{131 \cdot 0 \pm 4 \cdot 03}{119 \cdot 4 \pm 3 \cdot 79}$	9·03 10·72	
Tail	5 ở ở 8 qq	$\begin{array}{rrr} 100 & -114 \\ 83 & -105 \end{array}$	$\begin{array}{c} 107 \cdot 2 \pm 2 \cdot 39 \ *(82 \%) \\ 93 \cdot 5 \pm 3 \cdot 13 \ (78 \%) \end{array}$	5·36 8·86	
†Ear	5 33 7 99	$     \begin{array}{r}       19 & - & 21 \\       16 & - & 19     \end{array} $	$\begin{array}{cccc} 19 \cdot 4 \pm 0 \cdot 40 & (15\%) \\ 17 \cdot 3 \pm 0 \cdot 60 & (15\%) \end{array}$	0·90 1·60	
Pes	5 33 8 99	$21 - 24 \\ 19 - 23$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1·33 1·27	

## Body measurements of *Antechinus flavipes* from North-east Queensland Measurements in millimetres

\* Per cent of head-body length.

 $\dagger$  "Crown" measurement +5 mm.

## TABLE 4

Body measurements of Antechinus flavipes from Cape York Peninsula, Queensland Measurements in millimetres

		Range	Mean±S.E.	Standard deviation
Head-body	40 ♂♂ 27 ♀♀	$\begin{array}{rrrr} 118 & -154 \\ 108 & -140 \end{array}$	$140 \cdot 0 \pm 1 \cdot 03 \\ 122 \cdot 4 \pm 1 \cdot 52$	6·57 7·92
Tail	40 ♂♂ 26 ♀♀	$\begin{array}{rrr} 101 & -133 \\ 91 & -110 \end{array}$	$\begin{array}{c} 118 \cdot 8 \pm 1 \cdot 00 \ *(85 \%) \\ 100 \cdot 7 \pm 0 \cdot 87 \ (82 \%) \end{array}$	6·33 4·46
†Ear	39 ♂♂ 25 ♀♀	$     \begin{array}{r}       16 - 20 \\       16 - 19     \end{array} $	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0·97 0·87
Pes	40 ♂♂ 25 ♀♀	$23 \cdot 5 - 27$ 21 - 24	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\begin{array}{c} 0\cdot 93 \\ 0\cdot 92 \end{array}$

\* Per cent of head-body length.

 $\dagger$  "Crown" measurement +5 mm.

20 per cent longer in head-body, tail and pes length than is A. f. flavipes. The data on the North-east Queensland form are less reliable, but they indicate that it is intermediate in size between the Cape York form and A. f. flavipes. As the figures for A. f. leucogaster show it to be about 10 per cent small xthan A. f. flavipes, it is apparent that a well-defined clinal gradient in body size exists, with the largest form at

March, 1967

the northern extremity and the smallest form at the south-western extremity of the species' arc-like range.

## (b) Manus and pes

The normal appearance of the palmar and plantar surfaces of the feet of A. flavipes is shown in Figure 5. In all specimens examined the first interdigital (pollical) and inner metacarpal pads on the manus were well

	Fusion on both feet		Fusion on one foot		Fusion on neither foot	
	N	%	N	%	N	%
flavipes (manus) stuartii " godmani "		Ξ	3	3	21 98 26	100 97 100
flavipes (pes) stuartii " godmani "	1 78 11	5 77 42	1 13 4	5 13 16	19 10 11	90 10 42

Fusion of first interdigital pad with inner metacarpal pad on manus and with inner metatarsal pad on pes in Antechinus flavipes, A. stuartii and A. godmani

separated, but in Jones (1923: 96) an illustration of the palm of A. *f. flavipes* shows the inner metacarpal lacking or fused with the first interdigital. This is apparently a rare occurrence.

In Figure 5 the inner metatarsal pad of the pes is shown distinctly separate from the first interdigital pad (hallucal). However in some specimens these pads were in contact or, more rarely, fused (see Table 5). This latter condition was noted on the pes of two specimens from Victoria.

In A. flavipes the granulations on the palm and sole are not pigmented as in A. swainsonii (see Wakefield and Warneke, 1963: 202) and are therefore less obvious. However in most specimens of A. flavipes we have seen, several conspicuously large granules were noted between the outer metatarsal pad and the heel, as can be seen in Figure 5. In most cases there were two, in others none, one or three.

## (c) Pelage

A. flavipes flavipes. The main pile is dense and moderately soft, but a profusion of long, coarse guard hairs imparts a rather crisp texture to the coat. There is marked antero-posterior differentiation in dorsal colour: the head and foreparts are dark grizzled grey and the lower back and rump are suffused with buff or reddish fawn. This toning is a composite effect due to an intermingling of several discrete colours in the fur. The dominant colour of the main pile is confined to a broad band between a basal zone of deep slate grey and an extreme terminal zone of dark brown to black; it is white on the head and shoulders and a rich ochraceous buff on the rump. A grizzled effect is produced by the interspersed guard hairs which are wholly black or tipped with light brown.

The flanks, ventrum and inner side of the limbs are in marked contrast, as guard hairs are lacking and the colour in the fur (ochraceous buff to golden tan) extends from the basal zone of slate grey right to the tip of each hair. The inguinal and gular areas are less brightly coloured.

Orbital crescents of light buff, and a patch of buff fur at the base and to the back of each ear, are very conspicuous. The ears are sparsely clothed with short adpressed buff hairs, some at the lower part of the conch being dark terminally. The dorsal surfaces of manus and pes are buff to ochraceous buff, undiluted by darker elements.

The tail is clothed with short ad-

pressed hairs which tend to mask rings of small epidermal scales underneath. Above it is buff, grizzled with black, becoming wholly black at the apex where the hairs are noticeably longer. Below it is deep buff, darkening towards the apex.

These notes on the pelage of A. *flavipes flavipes* were compiled after

examination of a large series of skins from South Australia, Victoria, New South Wales and south-eastern Queensland. For the most part these specimens agreed well with descriptions of both the type, from north-eastern New South Wales (Waterhouse, 1838), and the South Australian population (Jones, 1923: 95-6; and Fin-



Fig. 5. Foot characters of Antechinus flavipes. Left—Left pes. Lower right—Left manus. Upper right—Foreclaws.

layson, 1958: 145). However, in certain localities variants do occur, but only to the extent of a reduced intensity of the colour of the lower back, flanks and ventrum, to a pale buff. This was noted in single specimens from various localities and in whole series from certain Queensland localities: Imbil (26°28' S., 152°41' E.), Barney View (28°15' S., 152°47' E.), and Frazer Island (at about 25° S.) off the coast.



Fig. 6. Skull and mandible of an adult male Antechinus flavipes flavipes from Frazer Island, S.E. Queensland. (Queensl. Mus., No. J.11258). Fig. 7. Skull and mandible of an adult female

Antechinus flavipes leucogaster from Kalamunda, W. Aust. (FWD, No. D.642).

Vict. Nat.-Vol. 84

A. flavipes leucogaster. In the Western Australian animal the bright ochraceous tonings of the typical form are replaced by a subdued and sombre ochraceous brown. However, the physical components of the pelage and the pattern of colours are essentially the same in all respects save one; the ochraceous brown of the flanks does not extend across the ventrum. In ventral aspects, white-tipped fur extends from the gular to the in-



Photos: J. Cooper, FWD.

Fig. 8. Skull and mandible of the neotype of Antechinus stuartii stuartii, an adult male from Waterfall, National Park, N.S.W. (AM, No. M.5294).

March, 1967

Fig. 9. Skull and mandible of an adult male Antechinus godmani from Evelyn, N. Queensland (Queensl. Mus., No. J.3672).

		Range	Mean±S.E.	Standard Deviation	
Basalar length (B.L.)	45 ♂♂ 26 ♀♀	$\begin{array}{r} 25 \cdot 75 - 30 \cdot 80 \\ 23 \cdot 45 - 28 \cdot 20 \end{array}$	$\frac{28 \cdot 09 \pm 0 \cdot 21}{25 \cdot 80  0 \cdot 17}$	1 · 408 0 · 866	
Zygomatic breadth (Z.B.)	48 ♂♂ 28 ♀♀	$\begin{array}{c} 16 \cdot 15 - 20 \cdot 40 \\ 15 \cdot 75 - 18 \cdot 35 \end{array}$	${}^{18\cdot 34 \pm 0\cdot 15}_{16\cdot 81 \ 0\cdot 12}$	$\begin{array}{c}1\cdot032\\0\cdot620\end{array}$	
Postorbital constriction (P.C.)	49 ♂♂ 30 ♀♀	$5 \cdot 90 - 7 \cdot 05$ $5 \cdot 95 - 7 \cdot 20$	$     \begin{array}{r}       6 \cdot 52 \pm 0 \cdot 04 \\       6 \cdot 36 & 0 \cdot 05     \end{array} $	$0.273 \\ 0.291$	
Palatalar length (P.L.)	50 ♂♂ 30 ♀♀	$\begin{array}{r} 13 \cdot 80 - 16 \cdot 80 \\ 12 \cdot 75 - 15 \cdot 30 \end{array}$	${}^{15 \cdot 29 \pm 0 \cdot 11}_{14 \cdot 24 \ 0 \cdot 09}$	0·761 0·510	
Anterior palatine foramen	50 ♂♂ 29 ♀♀	$2 \cdot 25 - 3 \cdot 70$ $2 \cdot 25 - 3 \cdot 05$	${}^{2\cdot 89 \pm 0\cdot 04}_{2\cdot 71 \ 0\cdot 05}$	$\begin{array}{c} 0\cdot 294\\ 0\cdot 250\end{array}$	
Breadth at M <sup>3</sup> (B.M.)	50 ♂♂ 30 ♀♀	$9 \cdot 45 - 11 \cdot 10$ $9 \cdot 10 - 10 \cdot 60$	${}^{10\cdot27}_{9\cdot80}{}^{\pm0\cdot06}_{0\cdot06}$	$\begin{array}{c} 0\cdot 422\\ 0\cdot 334\end{array}$	
Length, $M^{1-3}$	51 ♂♂ 34 ♀♀	$\begin{array}{rrrr} 5 \cdot 40 - & 6 \cdot 50 \\ 5 \cdot 30 - & 6 \cdot 20 \end{array}$	$5 \cdot 87 \pm 0 \cdot 03 \\ 5 \cdot 70  0 \cdot 05$	$\begin{array}{c} 0\cdot 210\\ 0\cdot 304\end{array}$	
P.C./B.L., as per cent	42 ♂♂ 26 ♀♀	$\begin{array}{rrr} 19 \cdot 6 & -25 \cdot 6 \\ 22 \cdot 4 & -26 \cdot 6 \end{array}$	$\begin{array}{c} 23 \cdot 2 \\ 25 \cdot 4 \end{array} \begin{array}{c} \pm 0 \cdot 21 \\ 0 \cdot 32 \end{array}$	$\begin{array}{c}1\cdot 33\\1\cdot 63\end{array}$	
Z.B./B.L., as per cent	43 ♂♂ 25 ♀♀	$62 \cdot 1 - 68 \cdot 7$ $61 \cdot 9 - 67 \cdot 4$	$\begin{array}{c} 65 \cdot 3 \\ 64 \cdot 9 \\ 0 \cdot 35 \end{array} \pm \begin{array}{c} 0 \cdot 25 \\ 0 \cdot 35 \end{array}$	$\begin{array}{c}1\cdot 60\\1\cdot 78\end{array}$	
P.L./B.L., as per cent	43 ♂♂ 26 ♀♀	$52 \cdot 4 - 56 \cdot 7 52 \cdot 6 - 57 \cdot 7$	$54 \cdot 4 \pm 0 \cdot 14 55 \cdot 0  0 \cdot 23$	0·92 1·19	
B.M./B.L., as per cent	44 ♂♂ 26 ♀♀	$34 \cdot 0 -39 \cdot 0$ $34 \cdot 8 -40 \cdot 0$	$36.7 \pm 0.17 \\ 38.0  0.23$	1 · 11 1 · 16	
P.C./Z.B., as per cent	48 ♂♂ 28 ♀♀	$\begin{array}{r} 30 \cdot 8 & -39 \cdot 2 \\ 34 \cdot 8 & -41 \cdot 2 \end{array}$	$\begin{array}{c} 35 \cdot 6 \\ 37 \cdot 8 \\ 0 \cdot 30 \end{array} \pm \begin{array}{c} 0 \cdot 30 \\ 0 \cdot 30 \end{array}$	$\begin{array}{c} 2\cdot 06\\ 1\cdot 58\end{array}$	

## Cranial measurements of Antechinus flavipes flavipes Measurements in millimetres

guinal region and onto the inner side of the limbs. The whiteness is variably modified by the basal colour of slate grey showing through.

North-east Queensland series. The dorsal colour pattern of specimens from this region is the same as that of A. f. flavipes, with strong anteroposterior differentiation and overtones of black due to the guard hairs. The head and shoulders are dark, lightly tinged with brownish buff, and the lower back and rump are a warm rust brown. The latter colour is especially bright on the flanks where the guard hairs are lacking, but on the ventrum it is much reduced in intensity and is

further diluted by a grey undertone due to a basal zone of slate grey.

Orbital crescents and ear patches of brown or brownish buff are inconspicuous in some specimens. The dorsal surface of both manus and pes is brownish buff. The tail is darker above as in the typical form but with darker tones of brown corresponding with those of the body fur, and is dark brown to black at the apex.

Cape York series. The pelage of specimens from this region is quite distinct. The pile is of shorter length and is more coarse than that of A. f. flavipes even though the guard hairs are less strongly developed, and be-

-		Range	Mean±S.E.	Standard deviation
Basalar length (B.L.)	8 ởở 10 ⊊♀	$\begin{array}{r} 25 \cdot 50 - 27 \cdot 70 \\ 23 \cdot 40 - 26 \cdot 55 \end{array}$	$\frac{26 \cdot 56 \pm 0 \cdot 30}{24 \cdot 79  0 \cdot 28}$	0·846 0·886
Zygomatic breadth (Z.B.)	9 ♂♂ 10 ♀♀	$16 \cdot 00 - 19 \cdot 50 \\ 14 \cdot 70 - 17 \cdot 40$	${}^{17\cdot40\pm0\cdot33}_{15\cdot830\cdot25}$	0·984 0·794
Postorbital constriction (P.C.)	10 ởở 13 qq	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$6 \cdot 32 \pm 0 \cdot 09 \\ 6 \cdot 15  0 \cdot 04$	0·270 0·143
Palatalar length (P.L.)	11 ♂♂ 12 ♀♀	$13 \cdot 75 - 15 \cdot 00 \\ 13 \cdot 00 - 14 \cdot 55$	${}^{14\cdot40\pm0\cdot13}_{13\cdot58}_{0\cdot11}$	0·414 0·374
Anterior palatine foramen	13 ♂♂ 16 ♀♀	$2 \cdot 50 - 3 \cdot 10$ $2 \cdot 20 - 3 \cdot 25$	$2 \cdot 82 \pm 0 \cdot 05$ $2 \cdot 65  0 \cdot 07$	0·195 0·286
Breadth at M <sup>3</sup> (B.M.)	11 ♂♂ 12 ♀♀	$9 \cdot 30 - 10 \cdot 20$ $8 \cdot 75 - 9 \cdot 65$	$9 \cdot 68 \pm 0 \cdot 08 \\ 9 \cdot 20  0 \cdot 08$	0·276 0·284
Length, M <sup>1-3</sup>	11 33 13 99	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$5 \cdot 43 \pm 0 \cdot 05 \\ 5 \cdot 42  0 \cdot 06$	0·180 0·215
P.C./B.L., as per cent	8 ♂♂ 10 ♀♀	$\begin{array}{r} 22 \cdot 9 & -25 \cdot 2 \\ 23 \cdot 7 & -26 \cdot 2 \end{array}$	$\begin{array}{ccc} 23 \cdot 8 & \pm 0 \cdot 29 \\ 24 \cdot 8 & 0 \cdot 23 \end{array}$	0·82 0·74
Z.B./B.L., as per cent	8 33 9 99	$\begin{array}{rrr} 62 \cdot 7 & -70 \cdot 4 \\ 60 \cdot 5 & -65 \cdot 5 \end{array}$		$\begin{array}{c}2\cdot 33\\1\cdot 86\end{array}$
P.L./B.L., as per cent	8 ♂♂ 10 ♀♀	$53 \cdot 5 - 55 \cdot 4$ $53 \cdot 8 - 55 \cdot 5$	$54 \cdot 3 \pm 0 \cdot 24 \\ 54 \cdot 9  0 \cdot 16$	0·67 0·51
B.M./B.L., as per cent	8 ♂♂ 10 ♀♀	$35 \cdot 4 - 37 \cdot 6$ $35 \cdot 8 - 39 \cdot 1$	${36\cdot 5 \atop 37\cdot 0} {\pm 0\cdot 31 \atop 0\cdot 36}$	0·86 1·29
P.C./Z.B., as per cent	9 ♂♂ 10 ♀♀	$33 \cdot 3 - 39 \cdot 3$ $36 \cdot 2 - 42 \cdot 3$	$36.6 \pm 0.64 \\ 38.8 \pm 0.63$	$1.91 \\ 2.00$

Cranial measurements of *Antechinus flavipes leucogaster* Measurements in millimetres

cause the guard hairs are not prominent the fur over the head and shoulders has not the cold tones which are characteristic of the southern and already western forms described; differentiation antero-posterior is slight and indistinct. The general appearance is of slightly grizzled pale brown over the whole dorsum, scarcely warmer with a tinge of rust brown on the rump, and pale brown on the flanks, ventrum and inner side of the limbs. The basal zone of the main pile is everywhere slate grey.

Orbital crescents are absent and ear patches are not differentiated, other than by the absence of darker elements which are present in the ad-

March, 1967

jacent fur. The dorsal surface of both manus and pes is fawn. The colours of the upper and under surfaces of the tail correspond with the adjacent body fur, darkening to brown or occasionally black at the apex.

## V. SKULL AND DENTITION OF ANTECHINUS FLAVIPES

For its size the skull is quite massively constructed; it is broad across the zygomatic region, is of moderate depth, and has a short conical rostrum (see Figures 6 and 7). Features of particular significance are the relatively narrow postorbital constriction (narrower in relation to zygomatic breadth in mature animals than in the young), absence of postorbital processes, and short anterior palatine foramina. The alisphenoid bulla is inflated and rounded in form while the "mastoid" portion, which is almost entirely enclosed by the periotic, is scarcely so.

The teeth are robust and powerful.  $I^1$  is at least twice the bulk of  $I^2$ , is cylindrical or very slightly flattened laterally, markedly proödont and separated from  $I^2$  by a wide diastema. 12-4 crowded together with are their triangular crowns overlapping;  $I^2 > I^3 > I^4$ . The upper canine is a broad tapered peg, slightly curved.  $P^4 > P^3 > P^1$ , with  $P^4$  at least twice the bulk of P<sup>1</sup>. There is no metaconic projection at the rear of M<sup>4</sup>. The lower incisors are semi-recumbent;  $I_1 > I_2 > I_3$ . The lower canine is short, with a distinct posterior basal ledge.  $P_3 > P_1 > P_4$ .

The deciduous premolar  $dP^4$ , which is lost early in life, is a tiny lowcrowned molariform tooth with three roots. By comparison,  $dP_4$  is even smaller, being less than half its bulk. It is similar to the permanent premolars in form but is proportionately broader at the base and has a much lower crown. It has two roots which may be either divergent or contiguous.

A summary of seven cranial measurements and five skull proportions of each of the four geographically isolated populations is set out in Tables 6-9. Using the "t" test, the three variants have been compared statistically with the typical form.

A. f. leucogaster. According to Tate (1947: 128) this is a "thoroughly good race with larger bullae and smaller teeth than any of the east Australian races". A cursory examination of the average values of the skull proportions in Table 7 is sufficient to show that A. f. leucogaster is virtually identical to A. f. flavipes in the form of the skull, but in all the linear measurements taken it is of smaller size. Most of these linear differences from the typical form are not of high statistical significance, but in the case of breadth at M<sup>3</sup> both male and female A. f. leucogaster are quite distinctly smaller than the typical form (P<0.001 for both tests).

In all morphological features the dentition of the two forms is essentially the same.

North-east Queensland series. Again, the skull proportions of this form differ to only a minor degree from those of the typical form. From the few specimens measured, males average about the same size and females slightly larger in skull size than their southern counterparts. There is not, however, a skull measurement common to both sexes which sets the North-east Queensland form apart from A. f. flavipes.

Cape York series. Skulls from this region were the largest seen; the average dimensions of the female series exceeding, in most measurements, those of the series of A. f. flavipes males. The statistical significance of the lineal differences between these two forms were all of high order. On the other hand the two forms are closely similar in skull proportions. The greatest difference, in the proportion Z.B./B.L. (Table 9), does suggest that the Cape York form is narrower in the zygomatic region, but statistically this difference is not significant.

Sexual dismorphism. The most obvious sexual differences in the skull were those of length and width. In A. f. flavipes, A. f. leucogaster and in the Cape York form, differences in

Vict. Nat.—Vol. 84



Comparison of Antechinus stuartii and Antechinus flavipes Above: A. stuartii, J, Glenlofty, Victoria (FWD, No. DY.417). Below: A. flavipes, J, near Nelson, Victoria (FWD, No. DY.409).

(Note: In the lower picture the normal grey component of face and back coloration has been rendered bluish by the photography.)

basalar length, palatalar length and zygomatic breadth were found to be of high statistical significance (P<0.001), but in the small series from North-east Queensland no appreciable sexual dimorphism in the skull was demonstrated.

The length of the anterior palatine foramen was a remarkably constant feature between the sexes of each of the four populations. Postorbital constriction, molar row and breadth at  $M^3$  are not influenced by sex to any marked degree, except in A. f. flavipes and A. f. leucogaster where males are significantly broader at  $M^3$ .

When A. flavipes is considered in terms of its distribution along the Eastern Highlands and west across southern Australia, several anatomical trends are apparent. The most obvious is skull size (length and breadth), the largest occurring in Cape York Peninsula and the smallest in south-western Western Australia. A second and related trend can be seen in the length of the tooth row (see also Tate, 1947: 126-7). The average alveolar length of  $M^{1-3}$  of the South Australian series examined by us was  $5.95 \pm 0.04$ mm.

Tate (1947: 126), in discussing the *A. flavipes* group (in which he included *A. stuartii* and *A. godmani*), described an ascending gradient in relative bulla size from the Papuan region through north Queensland to south and finally south-western Australia. This slight trend was apparent in the complex of the four populations of *A. flavipes* examined in this paper (see Figures 6 and 7).

## VI. BREEDING CONDITION OF ANTECHINUS FLAVIPES

*Females.* In the non-breeding condition the pouch area is scarcely differentiated, its site on the lower abdomen being indicated by a small patch of rather coarse hairs of a uniform light fawn colour. Nipple counts made on females in this condition are difficult and unreliable as the nipples are quite minute and do not project from the skin.

As breeding is restricted to a short period each year and the populations are low at that time, relatively few females showing signs of breeding activity are represented in museum collections and, unfortunately, not all of them are dated. However, those with reliable data indicate that in Victoria the young are born in early August (NMV. No. R.12784), while in north Queensland they are born between and early September mid-August (AMNH series). No data on the Western Australian population were available.

In Victoria, lactation continues through to mid-November at least, by which time the young are no longer carried by the mother (FWD. No. D.233). In this specimen the mammary area was large and conspicuous, being devoid of hair and stained reddish brown. Its lateral margins were marked by raised ridges of skin and anteriorly it was bordered by a conspicuous fringe of long reddish hairs. Six pairs of elongated nipples were arranged symmetrically, parallel to the lateral margins.

In A. flavipes the nipple number is variable. In the Western Australian form satisfactory counts were obtained from only two specimens, both of which had 10. In the typical form the number is 10 or 12, and either may be noted in females from the same locality. Only two reliable counts from North-east Queensland were available, both of which were 8. Counts made on Cape York females were of either 10 or 8 nipples, but most of the latter were recorded with a query.

		Range	Mean±S.E.	Standard Deviation
Basalar length (B.L.)	9 33 9 44	$\begin{array}{r} 25 \cdot 95 - 31 \cdot 35 \\ 26 \cdot 15 - 28 \cdot 70 \end{array}$	$\frac{28 \cdot 29 \pm 0 \cdot 50}{26 \cdot 42  0 \cdot 45}$	1·490 1·340
Zygomatic breadth (Z.B.)	9 33 9 9	$\frac{17 \cdot 00 - 19 \cdot 60}{16 \cdot 70 - 19 \cdot 00}$	${}^{18\cdot37\pm0\cdot28}_{17\cdot60}_{0\cdot22}$	0.835 0.658
Postorbital constriction (P.C.)	9 ♂♂ 10 ♀♀	$\begin{array}{r} 6\cdot 50-\ 7\cdot 30\\ 6\cdot 00-\ 7\cdot 20\end{array}$	$6 \cdot 80 \pm 0 \cdot 08 \\ 6 \cdot 73  0 \cdot 12$	0·250 0·369
Palatalar length (P.L.)	9 33 9 99	$\begin{array}{c} 13 \cdot 90 - 17 \cdot 05 \\ 14 \cdot 35 - 15 \cdot 70 \end{array}$	${ \begin{array}{c} 15 \cdot 25 \pm 0 \cdot 33 \\ 14 \cdot 79 & 0 \cdot 17 \end{array} } $	0·984 0·517
Anterior palatine foråmen	9 33 10 ⊊⊊	$\begin{array}{r} 2 \cdot 40 - 3 \cdot 20 \\ 2 \cdot 40 - 3 \cdot 00 \end{array}$	$2 \cdot 82 \pm 0 \cdot 09 \\ 2 \cdot 71  0 \cdot 06$	0·254 0·187
Breadth at M <sup>3</sup> (B.M.)	9 33 9 9 9	9·50-11·25 9·85-10·90	${}^{10\cdot19\pm0\cdot20}_{10\cdot40\ 0\cdot12}$	0·587 0·359
Length, M <sup>1-3</sup>	9 ♂♂ 10 ⊊♀	$\begin{array}{r} 5 \cdot 40 - \ 6 \cdot 20 \\ 5 \cdot 60 - \ 6 \cdot 50 \end{array}$	$5.86 \pm 0.11$ 5.98  0.09	0·315 0·279
P.C./B.L., as per cent	9 ♂♂ 10 ♀♀	$\begin{array}{r} 22 \cdot 0 & -25 \cdot 8 \\ 22 \cdot 4 & -27 \cdot 0 \end{array}$	$\begin{array}{c} 24 \cdot 1 \\ 24 \cdot 5 \\ 0 \cdot 66 \end{array} \pm \begin{array}{c} 0 \cdot 37 \\ 0 \cdot 66 \end{array}$	1 · 12 1 · 15
Z.B./B.L., as per cent	9 ♂♂ 10 ♀♀	$ \begin{array}{r}     61 \cdot 6 & -70 \cdot 4 \\     62 \cdot 0 & -66 \cdot 2 \end{array} $	$\begin{array}{c} 65 \cdot 0 \\ 64 \cdot 4 \\ 0 \cdot 49 \end{array} \pm \begin{array}{c} 0 \cdot 85 \\ 0 \cdot 49 \end{array}$	$\begin{array}{c} 2\cdot 54\\ 1\cdot 56\end{array}$
P.L./B.L., as per cent	9 ♂♂ 10 ♀♀	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$53 \cdot 3 \pm 0 \cdot 41 \\ 54 \cdot 8  0 \cdot 23$	$\begin{array}{c} 1\cdot 22\\ 0\cdot 74\end{array}$
B.M./B.L., as per cent	9 ♂♂ 10 ♀♀	$34 \cdot 1 - 37 \cdot 2$ $36 \cdot 0 - 39 \cdot 3$	$36 \cdot 2 \pm 0 \cdot 33 \\ 37 \cdot 9  0 \cdot 31$	0·98 1·00
P.C./Z.B., as per cent	9 ♂♂ 10 ♀♀	$35 \cdot 2 - 39 \cdot 4$ $35 \cdot 9 - 40 \cdot 7$	$\begin{array}{c} 37 \cdot 1 \\ 38 \cdot 1 \end{array} \begin{array}{c} \pm 0 \cdot 53 \\ 0 \cdot 43 \end{array}$	$\begin{array}{c} 1\cdot 60\\ 1\cdot 48\end{array}$

Cranial measurements of Antechinus flavipes from North-east Queensland Measurements in millimetres

Males. In other Antechinus dealt with by us (Wakefield and Warneke, 1963: 208), the only external sign of sexual maturity in males was the size of the testes, considerable enlargement occurring some weeks before pouch development in the females. Our observations on live Victorian specimens indicate that this is also the normal pattern of development in male A. flavipes.

## VII. TAXONOMY OF ANTECHINUS STUARTII

Macleay (1841) published both the genus *Antechinus* and the specific epithet *stuartii* as novelties to describe

a male specimen from Spring Cove (i.e. Manly), New South Wales. In the original diagnosis the number of upper incisors was given as 6, but this was later corrected by a note (Macleay, 1842) stating that the dental formula was the same as that of *Phascogale*.

The original description was made from the notes of "J. Stuart, Esq.", and it was stated that the specimen had been lost. However, Macleay's later note indicated that he had examined a (the?) skeleton. As recent endeavours to locate this skeleton in the Macleay Museum, Sydney, and elsewhere, have been unsuccessful,

		Range	Mean±S.E.	Standard Deviation
Basalar length (B.L.)	39 ♂♂ 20 ♀♀	$\begin{array}{r} 29 \cdot 00 - 32 \cdot 65 \\ 26 \cdot 75 - 29 \cdot 50 \end{array}$	$30 \cdot 38 \pm 0 \cdot 17$ 28 \cdot 14 \cdot 0 \cdot 18	1 · 048 0 · 787
Zygomatic breadth (Z.B.)	38 ♂♂ 21 ♀♀	$\begin{array}{c} 17 \cdot 80 - 20 \cdot 50 \\ 16 \cdot 60 - 18 \cdot 85 \end{array}$	${}^{19\cdot17\pm0\cdot10}_{17\cdot67}_{0\cdot15}$	0 · 590 0 · 704
Postorbital constriction (P.C.)	41 ♂♂ 26 ♀♀	$\begin{array}{c} 6\cdot 35 - \ 7\cdot 35 \\ 6\cdot 25 - \ 7\cdot 20 \end{array}$	${}^{6\cdot 90\pm 0\cdot 03}_{6\cdot 72\ 0\cdot 04}$	$\begin{array}{c} 0\cdot 217\\ 0\cdot 212\end{array}$
Palatalar length (P.L.)	40 ♂♂ 27 ♀♀	$\begin{array}{c} 15 \cdot 60 - 17 \cdot 85 \\ 14 \cdot 90 - 16 \cdot 50 \end{array}$	${}^{16\cdot68\pm0\cdot09}_{15\cdot52}_{0\cdot06}$	$\begin{array}{c} 0\cdot 569\\ 0\cdot 326\end{array}$
Anterior palatine foramen	41 ♂♂ 27 ♀♀	$2 \cdot 45 - 3 \cdot 55$ $2 \cdot 40 - 3 \cdot 55$	$3.00 \pm 0.05 \\ 2.90  0.05$	0·298 0·266
Breadth at M <sup>3</sup> (B.M.)	40 ♂♂ 27 ♀♀	$10 \cdot 25 - 11 \cdot 70$ $9 \cdot 75 - 11 \cdot 15$	${}^{10\cdot91\pm0\cdot05}_{10\cdot550\cdot06}$	0·290 0·317
Length, M <sup>1-3</sup>	42 ♂♂ 28 ♀♀	$5 \cdot 85 - 6 \cdot 75$ $5 \cdot 80 - 6 \cdot 30$	${}^{6\cdot13\pm0\cdot03}_{6\cdot04}_{0\cdot03}$	0·182 0·135
P.C./B.L., as per cent	39 ♂♂ 20 ♀♀	$\begin{array}{rrr} 20 \cdot 0 & -24 \cdot 5 \\ 22 \cdot 5 & -26 \cdot 1 \end{array}$	$\begin{array}{ccc} 22 \cdot 5 & \pm 0 \cdot 16 \\ 24 \cdot 0 & 0 \cdot 20 \end{array}$	0·98 0·89
Z.B./B.L., as per cent	37 ♂♂ 17 ♀♀	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	${62 \cdot 8 \pm \atop 62 \cdot 9 } {0 \cdot 23 \atop 0 \cdot 36 }$	$= 1 \cdot 37$ $1 \cdot 48$
P.L./B.L., as per cent	39 ♂♂ 20 ♀♀	$\begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{ccc} 54 \cdot 5 & \pm 0 \cdot 14 \\ 55 \cdot 0 & 0 \cdot 21 \end{array}$	0.89 0.96
B.M./B.L., as per cent	38 ởở 20 ♀♀	$33 \cdot 2 - 38 \cdot 0$ $36 \cdot 4 - 38 \cdot 9$	$\begin{array}{ccc} 35 \cdot 6 & \pm 0 \cdot 19 \\ 37 \cdot 7 & 0 \cdot 17 \end{array}$	$\begin{array}{c}1\cdot17\\0\cdot78\end{array}$
P.C./Z.B., as per cent	38 ♂♂ 21 ♀♀	$32 \cdot 4 - 38 \cdot 5$ $34 \cdot 3 - 41 \cdot 4$	$35 \cdot 9 \pm 0 \cdot 23 \\ 38 \cdot 3  0 \cdot 41$	$\begin{array}{c}1\cdot 44\\1\cdot 88\end{array}$
	and the second s			

Cranial measurements of Antechinus flavipes from Cape York Peninsula, Queensland Measurements in millimetres

we conclude that the type specimen, is indeed lost.

As a neotype we have selected a male specimen in the Australian Museum, No. M5294 (spirit specimen and skull), collected at Waterfall, National Park, New South Wales, on August 7, 1932. Its measurements in mm. are as follows: head-body 101, tail 94, ear 17 and pes 18; of the skull: basalar length 26.30, zygomatic breadth 16.90, postorbital constriction 6.95, palatalar length 14.70, anterior palatine foramen 2.75, breadth at M<sup>3</sup> 9.50, and length of M<sup>1-3</sup> 5.50.

Our reasons for accepting that Macleay's *Antechinus stuartii* is, in fact, the species we treat as such in this paper, and not any other eastern New South Wales phascogalinid, are as follows:

- 1. The large size (total length  $9\frac{1}{2}$ inches) excludes members of the *murina-leucopus* group of *Sminthopsis*, and size together with the comparatively long tail ("cauda fere corporio longitudinem aequante") also excludes *S. crassicaudata*.
- 2. The pelage colour ("fulvus abdomine artubusque subtus albescentibus"), and the coastal habitat, exclude *Antechinus flavipes*.
- 3. The comparatively long tail excludes *Antechinus swainsonii*.

The tail features ("cauda teres 4 pilosa gracilis") excludes Phascogale tapoatafa.

Gould (1854) described Antechinus unicolor from New South Wales (BM. No. 54,11.19.2). The plate is an accurate portrayal of the almost concolorous Antechinus stuartii. One of Gould's specimens, a female, like Macleay's original, is of greater size than any example of A. stuartii measured by us (see Table 10).

Le Souef and Burrell (1926) described Antechinus flavipes burrelli from Ebor, north-eastern New South Wales. The type specimens (AM. No. M.2593), and others from that locality, are of A. stuartii, and they do not demonstrate any feature which would justify subspecific status. Ebor is within the geographical range of what we regard as the nominate form of A. stuartii (see Figure 3); hence "burrelli" is treated as a subjective synonym of A. stuartii stuartii.

Iredale and Troughton (loc. cit.) misapplied the trinomial Antechinus flavipes flavipes to the central-eastern New South Wales population which is, in fact, the nominate form of A. stuartii. This error has been perpetuated in many subsequent publications dealing with that population (Troughton, 1941; Tate, 1947; Horner and Taylor, 1959; Marlow, 1961; etc.).

Thomas (1923) established the trinomial Phascogale flavipes adusta for specimens of Antechinus from Ravenshoe in the Atherton area of northeastern Queensland. Measurements of these, communicated to us by R. E. Hill, including the type specimen (BM. No. 22.12.18.54), and the original diagnosis, show that "adusta" belongs, not to A. flavipes, but to A. stuartii. Thomas himself later realized that two species were involved, designating one "flavipes" and the other "unicolor" on his specimen labels; "adusta" he placed with "unicolor" (W. D. L. Ride\*, in litt., 12.x.1965).

Not having examined sufficient specimens of this tropical population of A. stuartii, we are unable to assess its status. The average body and cranial measurements of 5 males and of 2 females that are given in Table 12 at least demonstrate that it is comparable in size and form to A. stuartii stuartii (see Table 10). The population appears to be geographically isolated (see Figure 3) and can be distinguished as A. stuartii adustus, pending more knowledge of its morphology and distribution.

The distribution of A. stuartii overlaps that of A. flavipes in several areas. The two have been shown to be truly sympatric by the trapping of typical specimens of each at (a) Glenlofty, western Victoria (37°04' S., 143°12' E.), and (b) "Stoney Creek", midway between Murrumbateman and Gundaroo, south-eastern New South Wales  $(35^{\circ}00' \text{ S.}, 149^{\circ}09' \text{ E.})$ , the latter by P. Woolley, A.N.U., Canberra. This evidence of reproductive isolation supports our action in recognizing the two forms as distinct species.

## VIII. DISTRIBUTION AND HABITAT OF ANTECHINUS STUARTII

The nominate race, A. s. stuartii, extends from the extreme south-east of Queensland, through eastern New South Wales and eastern and central Victoria, to the extreme south-west of Victoria. In this area it occurs in the highlands about the Great Dividing Range and in the coastal tracts between the highlands and the sea.

Essentially, its area of distribution is on the coastal side of that of A. f. flavipes; the ranges of the two are complementary, with little overlap. On the other hand, the general range of

88

<sup>\*</sup> Director, Western Australian Museum. Vict. Nat.-Vol. 84

A. s. stuartii is practically the same as that of A. swainsonii mimetes (see Wakefield and Warneke, 1963, Figure 2), and these two are commonly sympatric.

A. s. stuartii occupies a comparatively wide variety of habitats within the general categories of wet sclerophyll forest and dry sclerophyll forest. Rainfall on these habitats ranges from about 25 to over 60 inches per annum, and the elevations are from sea level to at least 4500 feet. Our observations suggest that this species is the most abundant marsupial in south-eastern Australia.

At Loch Valley the rainfall is about 56 inches per annum, and the natural vegetation is a forest of White Mountain Ash (*Eucalyptus regnans*), with a mid-storey of Silver Wattle (*Acacia dealbata*) and some Myrtle Beech (*Nothofagus cunninghamii*), and a dense under-storey of shrubbery and ferns. In a similar habitat, in the Marysville area of east-central Victoria, several specimens were caught in traps placed about 20 feet above the ground on limbs of Silver Wattle trees.

Specimens of A. stuartii were found in a fairly open forest of Snow Gum (Eucalyptus pauciflora) at 4500 feet elevation, in the upper Buchan River area (eastern Victoria); they were identified as A. flavipes (Wakefield, 1960). In the same district, at about 3000 feet near Mount Seldomseen, examples were trapped in a dense **Broad-leaf** forest of Peppermint (Eucalyptus dives), Messmate (E. obliqua) and Silver Wattle, with numerous logs and much low shrubbery.

In coastal dune scrubs at Mallacoota (eastern Victoria) the main trees of the habitat are Gum Myrtle (Angophora intermedia), banksias (Banksia integrifolia, B. marginata), Bracelet Honey-myrtle (Melaleuca armillaris), with bracken and several shrub species of the heath family (*Epacrida-ceae*). In the same district, specimens were caught in runways of bush rats (*Rattus fuscipes assimilis*) in an area almost devoid of trees, under a dense ground cover of Spear Grasstree (*Xanthorrhoea hastilis*) and numerous sedge species (*Cyperaceae*) and grasses.

Mount Clay near Portland At (south-western Victoria) the habitat is fairly open forest of stringybark eucalypts (E. baxteri, E. obligua) with fallen logs and a dense ground cover of bracken, Dwarf She-oak (Casuarina), flat-pea (Platylobium), Flame Heath (Astroloma conostephioides) and speargrass (Stipa muelleri). The population of A. stuartii in this general area was cited by Finlayson (1958) as the "Heathmere variant" of A. flavipes. In the Gorae Forest, west of Mount Clay, A. stuartii has been caught within 10 miles of the lower Glenelg habitat of A. flavipes (see Section III), but the ranges of the two species have not been found to overlap in this district.

In the Ballarat district (western Victoria), with a rainfall of 27 inches per annum, there is a comparatively pale variant of A. stuartii in forests of Narrow-leaf Peppermint (E. radiata) and Messmate, with a light ground cover of tussock-grass (Poa), some bracken and scattered clumps of Small Grasstree (Xanthorrhoea minor). Homes are made in hollow limbs and trunks of the peppermint trees, and many nests have been found in stacks of firewood that have been left in the bush for two or three years to dry. The nests are of leaves and bark.

The Glenlofty habitat, where A. stuartii and A. flavipes are sympatric, is described in Section III. A single specimen of A. stuartii was trapped there amongst tussock-grass and sawsedge (Gahnia radula) which provided a 50 per cent ground cover on a nar-

Body	measurements	of	Antechinus	stuartii	stuartii	(total	series)
	Me	asu	irements in	millime	tres		

		Range	Mean±S.E.	Standard Deviation
Head-body	123 ♂♂ 63 ♀♀	$77 -121 \\ 74 -120$	$97.4 \pm 0.70$ 91.2 $1.18$	7·80 9·29
Tail	119 ♂♂ 67 ♀♀	$\begin{array}{rrr} 73 & -109 \\ 65 & -99 \end{array}$	$\begin{array}{c}94 \cdot 1 \pm 0 \cdot 73 \ * (97 \%) \\84 \cdot 3 \ 0 \cdot 91 \ (92 \%)\end{array}$	7·99 7·50
Ear	109 ♂♂ 73 ♀♀	$13 \cdot 5 - 19$ $13 \cdot 5 - 19$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 · 28 1 · 51
Pes	126 ♂♂ 78 ♀♀	$\begin{array}{rrrr} 15 \cdot 5 - & 19 \cdot 5 \\ 14 & - & 20 \end{array}$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0·89 1·06

\* Per cent of head-body length.

T	4 4
ARIE	
IADLL	11

Body measurements of Antechinus stuartii stuartii (Loch Valley series) Measurements in millimetres

		Range Mean±S.E.		Standard Deviation	
Head-body	58 ♂♂ 13 ♀♀	$77 -107 \\ 76 - 93$	$95 \cdot 3 \pm 0 \cdot 85 \\ 84 \cdot 9  1 \cdot 45$	$\begin{array}{r} 6\cdot 49 \\ 5\cdot 23 \end{array}$	
Tail	56 ♂♂ 13 ♀♀	$     \begin{array}{r}       80 & -108 \\       78 & -99     \end{array} $	98·0±0·75 *(103%) 87·5 1·73 (103%)	$5 \cdot 64 \\ 6 \cdot 23$	
Ear	58 ♂♂ 13 ♀♀	$ \begin{array}{r} 13.5 - 17 \\ 14 - 16.5 \end{array} $	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0·87 0·75	
Pes	58 ♂♂ 13 ♀♀	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	0.98 0.82	

\* Per cent of head-body length.

row strip of flat ground along a dry water course. It had a pale pelage typical of the Ballarat variant (see Plate). Specimens of A. flavipes were caught 65 yards to the south, 260 vards to the east, and about half a mile to the west and south respectively of the locus of the A. stuartii specimen, some in traps placed about logs within the margin of the sedge-grass flat. In addition to the eucalypts previously the Candlebark mentioned. Gum (E. rubida) occurred at that spot (Figure 4).

The Grampians population of A. stuartii is within the geographical range of A. flavipes, and both species are distributed generally in these mountains. A specimen of A. stuartii and one of A. flavipes were trapped at the same spot, about 2 miles west of Mount Frederick, but at different times.

Troughton (1941) gave details of A. stuartii inhabiting crevices in the sandstones of the Sydney area, though he identified it as A. flavipes, and he commented on the agility of the ani-

Vict. Nat.—Vol. 84

90

	33	<u></u>		33	99
Head-body	109 · 2	100.0	Anterior palatine foramen	2.96	2.80
Tail	94.0	87.0	Breadth at M <sup>3</sup> (B.M.)	8.91	8.90
Ear	No	data	Length, M <sup>1-3</sup>	5.24	5.30
Pes	21.8	20.5	P.C./B.L., as per cent	26.3	26.8
Basalar length (B.L.)	25.25	24 · 50	Z.B./B.L., as per cent	61.9	62.4
Zygomatic breadth (Z.B.)	15.78	15.25	P.L./B.L., as per cent	54 · 5	55.3
Postorbital constriction (P.C.)	6.55	6.55	B.M./B.L., as per cent	35.3	36.3
Palatalar length (P.L.)	13.79	13.50	P.C./Z.B., as per cent	41.0	43.0

Antechinus stuartii adustus: average body and cranial measurements of 5 males and of 2 females Measurements in millimetres

mals in running about on the walls and ceilings of caves. In Victoria, evidence of cliff occupancy by *A. stuartii* was provided by several specimens trapped at Murrindal (eastern Victoria) and one at Glen Aire on the Otway Peninsula (south-western Victoria), in both cases at limestone cliffs.

Wakefield (1954) commented on the climbing ability and speed of movement of *A. stuartii* (again as *A. flavipes*), and recorded several observations of individuals abroad in full daylight. The species is normally nocturnal and, like *A. flavipes*, its usual food is insects, obtained partly about tree trunks and limbs and partly about logs and amongst ground litter.

A. s. adustus is known only from Ravenshoe and near Townsville, in north-eastern Queensland, and it is presumably an animal of the tropical rainforests.

## IX. PLASTIC CHARACTERS OF ANTECHINUS STUARTII

The general comments on A. flavipes at the beginning of Section IV apply equally well to A. stuartii, as the external differences between them,

March, 1967

except in pelage colour, are not very obvious. In the following notes attention has been drawn to the major points of distinction by which A. stuartii can be recognized.

#### (a) Body measurements

The body measurements of a series of 123 males and 63 females are summarized in Table 10. Specimens from all parts of the species range are represented but the bulk of the series is from Loch Valley (38°00' S., 145°33' E.) in eastern Victoria. Details of the latter are included separately (Table 11) to show the variation in a relatively large sample from one locality.

It is apparent from these data that the body proportions of both sexes are similar and that males are larger than females in head-body, tail and pes length. This sexual dimorphism is of high statistical significance (P<0.001 in each case). If these same data are compared with those of *A*. *f. flavipes* (Table 1) it is seen that *A. stuartii* is a much smaller animal with a proportionately longer tail.

		Range	Mean±S.E.	Standard Deviation
Basalar length (B.L.)	122 ♂♂ 63 ♀♀	$\begin{array}{r} 21 \cdot 40 - 27 \cdot 70 \\ 21 \cdot 35 - 26 \cdot 20 \end{array}$	$\frac{24 \cdot 33 \pm 0 \cdot 11}{23 \cdot 35  0 \cdot 13}$	1·185 1·062
Zygomatic breadth (Z.B.)	120 ♂♂ 59 ♀♀	$\frac{14 \cdot 20 - 17 \cdot 40}{13 \cdot 50 - 16 \cdot 40}$	${ \begin{array}{c} 15\cdot 61 \pm 0\cdot 06 \\ 14\cdot 97 & 0\cdot 10 \end{array} } \\$	0.662 0.751
Postorbital constriction (P.C.)	134 33 70 99	$5 \cdot 90 - 7 \cdot 45$ $5 \cdot 85 - 7 \cdot 00$	$6 \cdot 68 \pm 0 \cdot 02 \\ 6 \cdot 38  0 \cdot 03$	$\begin{array}{c} 0\cdot 231\\ 0\cdot 234\end{array}$
Palatalar length (P.L.)	130 33 67 99	$11 \cdot 50 - 15 \cdot 65$ $11 \cdot 65 - 14 \cdot 60$	${}^{13\cdot 39\pm 0\cdot 06}_{12\cdot 94 0\cdot 08}$	0.630 0.688
Anterior palatine foramen	133 33 66 99	$2 \cdot 15 - 4 \cdot 50$ $2 \cdot 25 - 4 \cdot 60$	$2.71 \pm 0.04$ 2.68  0.05	$0.425 \\ 0.432$
Breadth at M <sup>3</sup> (B.M.)	130 ♂♂ 70 ♀♀	7·90- 9·95 7·70-10·00	$     \begin{array}{r}       8 \cdot 70 \pm 0 \cdot 04 \\       8 \cdot 64 & 0 \cdot 06     \end{array} $	0·440 0·505
Length, M <sup>1-3</sup>	128 ♂♂ 70 ♀♀	$\begin{array}{r} 4 \cdot 50 - 5 \cdot 70 \\ 4 \cdot 50 - 5 \cdot 60 \end{array}$	$5.07 \pm 0.03$ 5.09  0.03	0·289 0·292
P.C./B.L., as per cent	121 33 63 99	$\begin{array}{rrrr} 23 \cdot 2 & -30 \cdot 7 \\ 24 \cdot 5 & -30 \cdot 4 \end{array}$	$\begin{array}{c} 27 \cdot 5 \\ 27 \cdot 4 \\ 0 \cdot 16 \end{array} \pm \begin{array}{c} 0 \cdot 12 \\ 0 \cdot 16 \end{array}$	$\begin{array}{c}1\cdot 28\\1\cdot 28\end{array}$
Z.B./B.L., as per cent	115 33 57 99	$     \begin{array}{r}       60 \cdot 3 & -68 \cdot 7 \\       60 \cdot 0 & -69 \cdot 2     \end{array} $	${64\cdot 1 \atop 64\cdot 2 } {\pm 0\cdot 14 \atop 0\cdot 28}$	$\begin{array}{c}1\cdot 55\\2\cdot 13\end{array}$
P.L./B.L., as per cent	121 33 62 99	$51 \cdot 1 - 57 \cdot 5$ $53 \cdot 3 - 57 \cdot 2$	$\begin{array}{ccc} 54 \cdot 6 & \pm 0 \cdot 09 \\ 55 \cdot 3 & 0 \cdot 12 \end{array}$	0·98 0·93
B.M./B.L., as per cent	117 ♂♂ 63 ♀♀	$32 \cdot 9 - 38 \cdot 5$ $34 \cdot 6 - 39 \cdot 3$	$\begin{array}{c} 35 \cdot 7 \\ 37 \cdot 0 \\ \end{array} \begin{array}{c} \pm 0 \cdot 09 \\ 0 \cdot 15 \end{array}$	0·96 1·20
P.C./Z.B., as per cent	119 ♂♂ 58 ♀♀	$\begin{array}{r} 37 \cdot 6 \ -46 \cdot 2 \\ 38 \cdot 1 \ -47 \cdot 6 \end{array}$	$\begin{array}{c} 42 \cdot 9 \\ 42 \cdot 7 \\ 0 \cdot 24 \end{array} \pm \begin{array}{c} 0 \cdot 16 \\ 0 \cdot 24 \end{array}$	1 · 79 1 · 84

## Cranial measurements of Antechinus stuartii stuartii (total series) Measurements in millimetres

#### (b) Manus and pes

The feet of a series of 78 specimens from Victoria and New South Wales were examined. The appearance of the palmar and plantar surfaces are similar to those of *A. flavipes* (Figure 5). The chief difference between the feet of these two species lies in the frequency of fusion of the first interdigital and inner metacarpal (-tarsal) pads. Fusion occurs more frequently in *A. stuartii*, as shown by Table 5.

As in *A. flavipes*, several discrete and conspicuously large granules are normally present between the outer metatarsal pad and the heel. Another of similar size is often to be found near the outer margin of the third interdigital pad. In some 10 per cent of cases this latter structure is strongly striated, forming a super-numerary pad on the same transverse line as the interdigital pads but not exceeding half their size. Super-numerary pads were not observed on the comparatively few specimens available of *A. flavipes*.

The strong development of striated pads in A. flavipes and A. stuartii appears to be an adaptation to their arboreal habits and is in marked contrast to the condition seen in A. swainsonii and A. minimus, both of which are terrestrial in habit (Wakefield and Warneke, 1963). The foot pads of both A. flavipes and A. stuartii are large and prominent, and only in the

Vict. Nat.—Vol. 84

			the second s	
		Range	Mean±S.E.	Standard Deviation
Basalar length (B.L.)	52 ♂♂ 12 ♀♀	$\begin{array}{r} 21 \cdot 40 - 24 \cdot 55 \\ 21 \cdot 35 - 23 \cdot 05 \end{array}$	$ \begin{array}{r} 23 \cdot 55 \pm 0 \cdot 08 \\ 22 \cdot 03  0 \cdot 12 \end{array} $	0·585 0·432
Zygomatic breadth (Z.B.)	51 ♂♂ 11 ♀♀	$14 \cdot 45 - 16 \cdot 10$ $13 \cdot 50 - 14 \cdot 85$	${}^{15\cdot25\pm0\cdot06}_{14\cdot270\cdot14}$	$\begin{array}{c} 0\cdot 423 \\ 0\cdot 465 \end{array}$
Postorbital constriction (P.C.)	59 ♂♂ 12 ♀♀	$6 \cdot 30 - 7 \cdot 10$ $6 \cdot 00 - 6 \cdot 70$	$6 \cdot 68 \pm 0 \cdot 03 \\ 6 \cdot 36  0 \cdot 06$	0·194 0·208
Palatalar length (P.L.)	57 ♂♂ 12 ♀♀	$11 \cdot 50 - 13 \cdot 65$ $11 \cdot 65 - 12 \cdot 50$	${}^{12\cdot84}_{12\cdot08}{}^{\pm0\cdot05}_{0\cdot09}$	$\begin{array}{c} 0\cdot 380\\ 0\cdot 295\end{array}$
Anterior palatine foramen	58 ♂♂ 12 ♀♀	$2 \cdot 15 - 3 \cdot 20$ $2 \cdot 25 - 2 \cdot 90$	${}^{2\cdot 63}_{2\cdot 52} {}^{\pm 0\cdot 03}_{0\cdot 05}$	0·205 0·182
Breadth at M <sup>3</sup>	55 ♂♂ 12 ♀♀	$\begin{array}{r} 7 \cdot 90 - 8 \cdot 80 \\ 7 \cdot 70 - 8 \cdot 65 \end{array}$	$8 \cdot 36 \pm 0 \cdot 02 \\ 8 \cdot 05  0 \cdot 07$	0·180 0·245
Length, M <sup>1-3</sup>	57 ♂♂ 12 ♀♀	$4 \cdot 50 - 5 \cdot 10$ $4 \cdot 50 - 5 \cdot 30$	$_{4\cdot 84\pm 0\cdot 02}^{4\cdot 84\pm 0\cdot 02}_{4\cdot 73\ 0\cdot 05}$	0·140 0·170
P.C./B.L., as per cent	52 ♂♂ 12 ♀♀	$\begin{array}{rrrr} 26 \cdot 8 & -30 \cdot 7 \\ 27 \cdot 5 & -30 \cdot 4 \end{array}$	$\begin{array}{c} 28 \cdot 4 \\ 28 \cdot 9 \\ 0 \cdot 25 \end{array} \pm \begin{array}{c} 0 \cdot 11 \\ 0 \cdot 25 \end{array}$	0·80 0·88
Z.B./B.L., as per cent	-48 ♂♂ 11 ♀♀	$61 \cdot 2 - 67 \cdot 7$ $62 \cdot 0 - 67 \cdot 5$	${64\cdot 6 \atop 64\cdot 8} {\pm 0\cdot 22 \atop 0\cdot 54}$	1 · 53 1 · 79
P.L./B.L., as per cent	52 ♂♂ 11 ♀♀	$51 \cdot 1 - 56 \cdot 1$ $53 \cdot 7 - 56 \cdot 1$	$54 \cdot 5 \pm 0 \cdot 13 \\ 54 \cdot 9  0 \cdot 24$	0·95 0·81
B.M./B.L., as per cent	49 ởở 12 ♀♀	$33 \cdot 9 - 37 \cdot 6$ $35 \cdot 3 - 39 \cdot 3$	$35 \cdot 5 \pm 0 \cdot 10 \\ 36 \cdot 6  0 \cdot 37$	0·73 1·28
P.C./Z.B., as per cent	50 ♂♂ 11 ♀♀	$\begin{array}{r} 41 \cdot 6 \ -46 \cdot 2 \\ 42 \cdot 8 \ -47 \cdot 6 \end{array}$	$\begin{array}{c} 43 \cdot 9 \\ 44 \cdot 5 \\ 0 \cdot 40 \end{array} \pm \begin{array}{c} 0 \cdot 18 \\ 0 \cdot 40 \end{array}$	$\begin{array}{c}1\cdot 25\\1\cdot 32\end{array}$
		)		

## Cranial measurements of Antechinus stuartii stuartii (Loch Valley series) Measurements in millimetres

latter is there a high degree of fusion on the pes. Here fusion is not accompanied by a reduction in size, on the contrary, the compound pad may be even slightly larger than the combined length of typical separate pads.

The structure of the claws also appears to be related to habit. In both *A. flavipes* and *A. stuartii* they are light, yet strongly curved and very sharp, and are thus well suited to climbing (see Figure 5).

#### (c) Pelage

The pelage of A. stuartii is noticeably softer than that of A. flavipes, evidently because the guard hairs are less strongly developed and hence do not affect the overall texture to the same degree. In other respects the physical components of the pelage of the two species are similar.

In dorsal colouration *A. stuartii* is a dull drab brown, lightly flecked with fawn, the latter colour occurring as a subterminal band in the main pile. Antero-posterior differentiation is absent or scarcely developed. In some specimens the ventrum is pale grey with a tinge of fawn; in others it is more strongly tinged across the midbelly with a pale brown. Ear patches and orbital crescents are absent or faint and ill-defined. The dorsal surface of both manus and pes is drab brown to grey brown. The dorsal surface of

the tail near the base is brown, flecked with fawn; below it is greyish fawn.

Some specimens are noticeably darker in colour, and in general this appears to be related to habitat. The description given above applies to specimens from coastal areas and dry sclerophyll forests. Dark specimens with warmer tonings were taken in wet sclerophyll forests in central Victoria, at Mount Macedon (3300 feet), Mount Arnold (4300 feet) and Loch Valley (1900 feet).

The few skins of A. s. adustus that we examined could not be distinguished from these latter specimens.

## X. SKULL AND DENTITION OF ANTECHINUS STUARTII

Apart from significant size differences, the skull of *A. stuartii* closely resembles that of *A. flavipes* (see Figures 6 and 8). However, when fully adult series are compared, the bones of *A. stuartii* skulls appear lighter and the supra-occipital crests are much less developed than in *A. flavipes.* The form and structure of the bulla are the same in both species.

The respective dentitions, including the deciduous premolars, are identical except that the incisor gradient  $I^2>I^3>I^4$  is less marked in *A. stuartii*; in some specimens the upper incisors are equal in size.

Measurements and proportions of a series of skulls of *A. stuartii* from Victoria, New South Wales and southeastern Queensland are set out in Table 13. Data on a series from Loch Valley are included separately to show variation in a relatively large sample from one locality (see Table 14).

The skulls of both sexes have similar proportions, but those of males average about 4 per cent larger in linear dimensions than those of females. The sexual differences in skull length and width are both of high statistical significance (P < 0.001 in each case).

There is some indication from the data that the length of the molar row diminishes in a metrical cline from Queensland to Victoria, as 11 specimens from Mount Glorious (27°20' S., 152°45' E.) in Queensland averaged  $5.50 \pm 0.05$  mm.; 10 specimens Irvine (33°30' from Mount S... 150°27' E.) in the Blue Mountains of New South Wales  $5.33 \pm 0.04$  mm.; 7 specimens from Argalong (38°18' S.,  $148^{\circ}26'$  E.)  $5 \cdot 29 \pm 0.04$  mm.; and the Loch Valley series  $4.83 \pm 0.02$ mm. However, as the northern and south-western extremities of the species range are each represented by only a few specimens, the full range of this clinal variation is unknown.

Comparison of Tables 13 and 6 show that the skull and teeth of A. stuartii are much smaller than those of A. flavipes, but, in our interpretation of speciation within Antechinus, difference in skull size is not considered to be of primary importance. However the two species differ considerably in one cranial proportion; in relation to length, the skull of A. stuartii is slightly narrower in zygomatic breadth, but is also much broader in post-orbital width (see Z.B./B.L. and P.C./B.L. in Tables 13 and 6). In these two tables the proportion Postorbital Constriction/Zygomatic Breadth is given, and in this the difference between the two species is a high order of significance of (P < 0.001).

## XI. BREEDING CONDITION OF ANTECHINUS STUARTII

Females. Marlow (1961) gave very full descriptions of the external changes associated with breeding in this species, but under the name A. f. flavipes. These changes are similar to those noted in other species of Antechinus (Wakefield and Warneke, 1963; and

Vict. Nat.—Vol. 84

TABLE 1
---------

		Range	Mean±S.E.	Standard Deviation		
Head-body	13 33 25 99	$ \begin{array}{rrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$143 \cdot 0 \pm 2 \cdot 86$ 121 · 9 1 · 79	10·33 8·94		
Tail	13 33 25 99	$\begin{array}{rrr} 105 & -146 \\ 86 & -115 \end{array}$	$\begin{array}{c} 123 \cdot 8 \pm 3 \cdot 02 \ * (87 \%) \\ 104 \cdot 1 \ 1 \cdot 64 \ (86 \%) \end{array}$	10·90 8·18		
Ear	3 33 4 99	$   \begin{array}{r}     16 & - & 19 \\     14 \cdot 5 - & 16 \cdot 5   \end{array} $	$\begin{array}{ccc} 17 \cdot 3 & (12 \%) \\ 15 \cdot 5 & (13 \%) \end{array}$			
Pes	13 ♂♂ 25 ♀♀	21 - 27 19 - 24	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	1 · 74 1 · 64		

Body measurements of Antechinus godn	nan
Measurements in millimetres	

\* Per cent of head-body length.

this paper). The timing of events in Marlow's captive colony is similar to that which may be deduced from Victorian field records, as follows:

Birth occurs at about mid-September. The actual date was recorded in one instance where a female gave birth in captivity, on September 14-15, seventeen days after capture. Gestation in this species takes 32 to 34 days (Marlow, 1961: 208; and observations by R.M.W.).

The young remain attached to the nipples for about five weeks, and subsequent to October 20 the pouch area of all lactating females trapped was found unoccupied. A female, FWD. No. D.449, trapped on December 9, 1962, was still lactating, indicating that the young are suckled until early December at least.

The number of nipples in *A. stuartii* varies from locality to locality, but appears to be constant in certain areas. For example, all females from the Cape Otway and Portland areas of south-western Victoria had 6; those from the Grampians had 8; from the Ballarat area 10; from the ranges east of Melbourne (including Loch Valley) 8; and from north-eastern Gippsland (Murrindal and Wulgulmerang) 10.

March, 1967

To date all reliable counts from New South Wales specimens have been 8. Two females of *A*. *s*. *adustus* from Mount Spec, near Townsville, had 6 and 8 nipples respectively.

Males. The comments on A. flavipes in Section VI apply also to A. stuartii.

## XII. PLASTIC CHARACTERS OF ANTECHINUS GODMANI

#### (a) Body measurements

Thirty-eight specimens were available to us, all of which were collected in or near the Ravenshoe district on the Atherton Tableland, close to the type locality. Thomas's published measurements of the type (Thomas, 1923), an adult male, have been included in Table 15, as these were the only other reliable data on an authentic specimen that was not seen by us. The statistical analysis of head-body and pes length in this species, made by Horner and Taylor (1959), was based on a series of specimens which we have since examined and found to include A. flavipes as well as A. godmani. These authors accepted Tate's evaluation of godmani as a subspecies of A. flavipes (Tate, 1952) but drew attention to the fact that Tate also

recognized *adustus* (actually a form of *A. stuartii*, see Section VII) which occurs sympatrically with *A.* godmani—as a subspecies of *A. flavi*pes.

Table 15 shows a marked disparity between the sexes in body size, which is in keeping with the sexual dismorphism noted in other *Antechinus* (Troughton, 1941: 26; Horner and Taylor, 1959; Marlow, 1961; Wakefield and Warneke, 1963).

As far as can be judged from the series available, *A. godmani* is the largest species of *Antechinus* in eastern Australia (see also Thomas, 1923). It is slightly larger than the Cape York form of *A. flavipes* and is considerably larger than the North-east Queensland form of that species.

## (b) Manus and pes

In structure the manus and pes are closely comparable with those of A. flavipes and A. stuartii. However, as in the latter, there is a relatively high frequency of fusion of the first interdigital and inner metatarsal pads of the pes (see Table 5). In eleven of twenty-six specimens these pads were fused on both feet and in four of the others on one foot only. The first interdigital and inner metacarpal pads of the manus were distinctly separate in all twenty-six specimens.

On each of the pes of ten specimens, three small striated super-numerary pads were noted. These were on the sole, immediately below and in line with the second, third and fourth interdigitals respectively—an occurrence without parallel in the other *Antechinus* examined by us.

The form of the claws is similar to that of A. *flavipes* and A. *stuartii*.

## (c) Pelage

The main pile is moderately fine in texture and the guard hairs, though profuse, are relatively short. The general dorsal colour is a subdued cinnamon brown, with conspicuous glints of fawn due to a sub-terminal band of that colour in the main pile; the basal zone is slate grey. A sheen is imparted to the whole by the guard hairs, which are glistening black, fading to light brown at the tips. Antero-posterior differentiation in dorsal colour appears to be stronger in males than in females. with the rump noticeably warmer in tone than the head and upper back. In his description of the species, Thomas (1923) made no distinction between the sexes on this point. In both sexes the fur of the ventrum is a uniform greyish fawn at the tips and slate grey beneath.

Orbital crescents and basal ear patches are lacking. The sides of the face, especially at the angle of the jaw, are rufescent. The dorsal surface of both manus and pes is ochraceous brown or cinnamon brown. The tail is unlike that of any other eastern Australian *Antechinus*. The dorsal surface is almost naked, having only a sparse covering of short adpressed brown hairs, while on the underside the hairs are longer and more dense, and form a ventral crest which extends beyond the tail tip.

## XIII. SKULL AND DENTITION OF ANTECHINUS GODMANI

Because this species has been confused with *A. flavipes* (see Section XII), the following comments emphasize those features of the skull and dentition which are of diagnostic importance. Other features may be seen in Figures 6 and 9.

On the average the skull is larger than that of A. flavipes, is narrower in relation to length and has a proportionately longer rostrum. The postorbital constriction is noticeably broader in relation to zygomatic breadth and the frontal region is depressed as in A. swainsonii. The anterior palatine

Vict. Nat.-Vol. 84

		Range	Mean±S.E.	Standard Deviation
Basalar length (B.L.)	7 33 7 99	$\begin{array}{r} 30 \cdot 00 - 32 \cdot 85 \\ 27 \cdot 70 - 29 \cdot 60 \end{array}$	$ \begin{array}{r} 31 \cdot 18 \pm 0 \cdot 38 \\ 28 \cdot 24  0 \cdot 29 \end{array} $	$\begin{array}{c} 1\cdot 000\\ 0\cdot 696\end{array}$
Zygomatic breadth (Z.B.)	6 33 8 99	$17 \cdot 55 - 19 \cdot 75$ $16 \cdot 20 - 17 \cdot 40$	${}^{18\cdot73}_{16\cdot82} {}^{\pm0\cdot27}_{0\cdot14}$	0.669 0.396
Postorbital constriction (P.C.)	8 33 8 99	$7 \cdot 55 - 7 \cdot 95$ $7 \cdot 40 - 8 \cdot 05$	$7.76 \pm 0.04$ 7.70  0.07	0·118 0·185
Palatalar length (P.L.)	7 ♂♂ 8 ♀♀	$16 \cdot 50 - 18 \cdot 30$ $15 \cdot 55 - 16 \cdot 75$	$17 \cdot 20 \pm 0 \cdot 22$ 16 \cdot 02 \cdot 16	$\begin{array}{c} 0\cdot 590\\ 0\cdot 463\end{array}$
Anterior palatine foramen	6 ♂♂ 9 ♀♀	$2 \cdot 50 - 3 \cdot 00$ $2 \cdot 25 - 3 \cdot 15$	$2 \cdot 81 \pm 0 \cdot 09 \\ 2 \cdot 71  0 \cdot 11$	0·216 0·352
Breadth at M <sup>3</sup> (B.M.)	7 33 8 99	9.65-10.65 9.60-10.30	${}^{10\cdot24}_{9\cdot92}{}^{\pm0\cdot13}_{0\cdot07}$	0·351 0·208
Length, M <sup>1-3</sup>	8 33 8 99	$5 \cdot 95 - 6 \cdot 25$ $5 \cdot 80 - 6 \cdot 30$	$6 \cdot 11 \pm 0 \cdot 04 \\ 6 \cdot 03  0 \cdot 06$	0·118 0·159
P.C./B.L., as per cent	7 ♂♂ 7 ♀♀	$\begin{array}{r} 23 \cdot 6 & -25 \cdot 7 \\ 25 \cdot 8 & -28 \cdot 4 \end{array}$	$\begin{array}{c} 25 \cdot 0 \\ 27 \cdot 2 \end{array} \begin{array}{c} \pm 0 \cdot 26 \\ 0 \cdot 34 \end{array}$	0·69 0·91
Z.B./B.L., as per cent	7 33 7 99	$58 \cdot 1 -61 \cdot 5$ $58 \cdot 4 -61 \cdot 2$	$\begin{array}{c} 59 \cdot 6 \\ 59 \cdot 7 \\ 0 \cdot 43 \end{array} \pm \begin{array}{c} 0 \cdot 51 \\ 0 \cdot 43 \end{array}$	1·36 1·17
P.L./B.L., as per cent	7 33 7 99	$53 \cdot 2 - 56 \cdot 1$ $56 \cdot 0 - 58 \cdot 4$	$\begin{array}{c} 55 \cdot 2 \\ 56 \cdot 7 \\ 0 \cdot 29 \end{array} \pm \begin{array}{c} 0 \cdot 39 \\ 0 \cdot 29 \end{array}$	1·04 0·77
B.M./B.L., as per cent	7 33 7 99	$31 \cdot 6 -34 \cdot 0$ $33 \cdot 6 -36 \cdot 2$	$\begin{array}{c} 32 \cdot 9 \\ 35 \cdot 1 \end{array} \begin{array}{c} \pm 0 \cdot 30 \\ 0 \cdot 37 \end{array}$	0·80 0·99
P.C./Z.B., as per cent	6 ♂♂ 8 ♀♀	$39 \cdot 2 -43 \cdot 5$ $43 \cdot 7 -48 \cdot 1$	$\begin{array}{c} 41 \cdot 5 \\ 45 \cdot 8 \\ 0 \cdot 54 \end{array} \pm \begin{array}{c} 0 \cdot 53 \\ 0 \cdot 54 \end{array}$	$ \begin{array}{r} 1 \cdot 31 \\ 1 \cdot 52 \end{array} $

Cranial measurements of Antechinus godmani Measurements in millimetres

foramina are short. The bulla is similar to that of *A*. *flavipes* and *A*. *stuartii* in structure, but the inflation of the alisphenoid does not extend as far anteriorly.

Of the incisors,  $I^1$  is of medium size, only slightly proödont and separated from  $I^2$  by a slight diastema;  $I^{2-4}$  are sub-equal, compressed laterally and not crowded together. The upper premolars are compressed laterally and are not separated by interspaces.  $M^4$ has a small distinct metacone.  $I_3$  bears a small accessory cusp on the buccal aspect near the heel, which slightly overlaps the canine.

Deciduous teeth were not found in any of the skulls examined by us.

In the outline of the skull as viewed

from above; in the form and position of I<sup>1</sup>; in the presence of the accessory cusp on I<sub>3</sub> and a metacone on M<sup>4</sup>, *A. godmani* shows a marked resemblance to *A. minimus* (see Wakefield and Warneke, 1963: 218).

## XIV. BREEDING CONDITION OF ANTECHINUS GODMANI

Females. On each of eight skins of females obtained in late November-December, 1921, the pouch area was greatly enlarged and most of the nipples were elongated, indicating that young were still being suckled. The appearance of the mammary region was essentially the same as that of other species at the same stage of development, and the period of year

when they were captured corresponds to the time of lactation in other species.

In four specimens 3 pairs of nipples were plainly visible, while in each of the other four specimens one nipple had regressed to a very small size but in each case was readily located by reference to the symmetry of the others. Thus the nipple number of A. godmani is 6.

Males. No information was available to us.

## XV. APPENDIX

Amended distribution of Antechinus minimus maritimus.

Since the publication of distributional data (Wakefield and Warneke, 1963), the following additional specimens of A. m. maritimus have been located:

- FWD. No. D.208. 9. Anglesea, Victoria. 28 Jan. 1957.
- FWD. No. D.582. J. Between Lighthouse Track and Waterloo Bay, Wilson's Promontory, S.E. Victoria. 5 Jan. 1965.
- FWD. No. D.647. Glennie Island, S.E. Victoria. 14 June 1966.

These records treble the known range of the mainland Australian race of A. minimus.

## XVI. ACKNOWLEDGEMENTS

We wish to acknowledge the cooperation of the Directors and staff members of the museums and other institutions, in Australia and overseas, who assisted with loan material, information and other facilities in connexion with the research and preparation of this paper.

The Commonwealth Bureau of Census and Statistics, Melbourne, gave advice in regard to the analysis of sets of measurements.

In connexion with the publication

of the paper, a major part of the cost of printing and of illustrations has been defrayed by a grant from the M. A. Ingram Trust.

### XVII. REFERENCES

- Brass, L. J., 1953. Results of the Archbold Expeditions. No. 68. Summary of the 1948 Cape York (Australia) Expedition. Bull. Amer. Mus. Nat. Hist. 102: Art. 2, pp. 135-206.
- Finlayson, H. H., 1958. A case of duplex convergent resemblance in Australian mammals, with a review of some aspects of the morphology of Phascogale (Antechinus) swainsonii Waterhouse and Phascogale (Antechinus) flavipes Waterhouse. Proc. Roy. Soc. S. Aust. **81**: 141-51.
- Fisher, R. A., and Yates, F., 1957. Sta-tistical tables for biological, agricultural and medical research. 5th Ed. Oliver & Boyd, London.
- Gould, John, 1863. Mammals of Australia. 3 vols., London.
- Gould, John, 1854. Mammals of Australia. Part vi (vol. 1, pl. 37), London.
- Gray, J. E., 1841. Contributions towards the geographic distribution of the mammals of Australia with notes on recently discovered species. some In Grey, G., Journals of two expeditions . . . in . . . Australia during the years 1837, 38, and 39. Appendix "C" vol. 2: 397-414.
- Horner, B. Elizabeth, and Taylor, J. Mary, 1959. Results of the Archbold Expeditions No. 80. Observations on the biology of the Yellow-footed Marsupial Mouse Antechinus flavipes flavipes. Amer. Mus. Nov., No. 1972.
- Iredale, Tom., and Troughton, E. Le G., 1934. A check-list of the mammals recorded from Australia. Mem. Austr. Mus. VI: 1-122.
- Jones, F. Wood, 1923. Mammals of South Australia. Part 1. Govt. Printer, Adelaide.
- Karmel, P. H., 1959. Applied statistics for economists. Pitman, Melbourne.
- *for economists.* Pitman, Merodume.
  Krefft, Gerard, 1866. On the vertebrated animals of the lower Murray and Darling, their habits, economy, and geographical distribution. *Trans. Phil. Soc. N.S.W.*, 1862-65: 1-33.
  Le Souef, A. S., and Burrell, H., 1926. *Wild animals of Australasia.* Harrap & Co. London
- & Co., London.
- Macleay, W. S., 1841. Notice of a new genus of Mammalia discovered by J.

Vict. Nat.-Vol. 84

Stuart, Esq., in New South Wales. Ann.

- Mag. Nat. Hist. VIII: 242, pl. vii. Macleay, W. S., 1842. Additional particulars respecting Antechinus Stuartii, a new marsupial quadruped. Ann. Mag. Nat. Hist. VIII: 337.
- Marlow, B. J., 1961. Reproductive behaviour of the marsupial mouse Antechinus flavipes flavipes Waterhouse (Marsupialia) and the development of the pouch young. Aust. J. Zool. 9
- (2): 203-18.
  Tate, G. H. H., 1947. Results of the Archbold Expeditions No. 56. On the anatomy and classification of the Dasyuridae (Marsupialia). Bull. Amer. Mus. Nat. Hist. 88 (3): 101-55.
- Tate, G. H. H., 1952. Mammals of Cape York Peninsula. Bull. Amer. Mus. Nat. Hist. 98: 567-616.
- Thomas, Oldfield, 1904. On a collection of mammals made by Mr. J. T. Tun-ney in Arnhem Land, Northern Terri-

tory of South Australia. Nov. Zool. x1: 229.

- Thomas, Oldfield, 1923. The Godman Exploration Fund: List of mammals from North Queensland collected by Mr. T. V. Sherrin. Ann. Mag. Nat. Hist. (series 9), XI: 174.
- Troughton, E. le G., 1941. Furred animals of Australia. Angus & Robertson, Sydney.
- Wakefield, N. A., 1954. Observations on marsupial mice. Vict. Nat. 71 (1): 6-8.
- Wakefield, N. A., 1960. Forlorn Hope and Reedy River. Vict. Nat. 77 (1): 4-11.
- Wakefield, N. A., and Warneke, R. M., 1963. Some revision in Antechinus (Marsupialia)—1. Vict. Nat. 80 (7): 194-219.
- Waterhouse, G. R., 1838. Characters of some new species of the genera Mus and Phascogale. Proc. Zool. Soc. Lond., 1837: 75.





# **Biodiversity Heritage Library**

Wakefield, N.

A.

٦

and Warneke, R. M. 1967. "Some Revision in Antechinus (Marsupialia)— 2." *The Victorian Naturalist* 84, 69–99.

View This Item Online: <a href="https://www.biodiversitylibrary.org/item/178977">https://www.biodiversitylibrary.org/partpdf/146536</a> Permalink: <a href="https://www.biodiversitylibrary.org/partpdf/146536">https://www.biodiversitylibrary.org/partpdf/146536</a>

Holding Institution Smithsonian Libraries and Archives

**Sponsored by** Biodiversity Heritage Library

## Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.