

Wall. ex Dc. var. *kurgensis* (Clarke) Noot., Rev. Sympl. 229. 1975.

Small tree, twigs sericeous. Leaves oblong to elliptic, $3-12.8 \times 0.8-5$ cm, acuminate, margins minutely spinous denticulate, base rounded or cuneate (on the same twig), young leaves sparsely pubescent both sides, older leaves glabrous except the midrib on the lower surface; nerves 7-11 pairs, arched. Petioles 0.9-2 cm long, sericeous. Inflorescence in axillary spikes, 5-9.1 cm long, rachis densely tawny tomentose. Bracts 3×1.5 mm, deltoid, acute, densely tawny tomentose both sides, caducous; bracteoles small, caducous. Flowers white, scented. Calyx 5 lobed; tube glabrous, 1 mm long; lobes 5, ovate, $2-3 \times 1-2$ mm, valvate, spathulate, tomentose without except the scarious margins. Corolla 5 lobed; lobes $5-6 \times 3$ mm, oblong, obtuse,

glabrous. Stamens 50-98, 3-5 mm long; anthers basifixed, 2 lobed. Pistil 6 mm long, style cylindric, base broad and hirsute; stigma capitate, disk 5 glandular. Fruits ovoid-cylindrical, $13-20 \times 6-8$ mm, stone with shallow lengthwise grooves, depressed at one side towards the base (description of fruit adopted from literature).

Specimens examined: INDIA: Kerala, Chandanathode, 710 m, 5-12-1967, Ellis 29476.

ACKNOWLEDGEMENTS

We wish to thank Dr. H. P. Nooteboom for confirming our identification and for the valuable information communicated through his letter. We are thankful to Dr. N. C. Nair, Dy. Director, for his help and encouragement.

SOUTHERN CIRCLE,
BOTANICAL SURVEY OF INDIA,
COIMBATORE 641 002,

January 4, 1980.

G. V. SUBBA RAO
R. GOPALAN

25. HOST PARASITE RELATIONSHIPS IN *DENDROPHTHOE FALCATA* (LINN.F.) BETTINGH (*LORANTHUS LONGIFLORUS* DESR.)

A wide spectrum of host plants known for *Dendrophthoe falcata* (Linn.f.) Ettingh (Loranthaceae) seems to be quite unique in the whole angiospermic parasites. This fact has been substantiated by a large number of host plants reported so far for this taxon (Fischer 1926, Sambandam 1966, Sampathkumar and Kunchithapatham 1969, Sampathkumar 1970). It has also been pointed out earlier that although there existed no specificity in the selection of host plants for *D. falcata*, the selection of host plants was not entirely promiscuous since the seeds of the parasite could not establish successfully in some monocots and also a few dicots (Sampathkumar 1970) where they initially germi-

nated but failed to establish later. One of the factors deciding the establishment of the parasite was thought to be osmotic pressure relationships between the host and the parasite (Sampathkumar 1970) for which experimental evidence presented in this paper lends support to this view.

The osmotic pressure relationships of the host as well as of the parasite in each case revealed in unambiguous terms that the parasite tends to have higher osmotic pressure than the host in question. Determination of osmotic pressure was made by plasmolytic method, using different molar concentrations of sucrose in which epidermal peelings of the leaves of

MISCELLANEOUS NOTES

host and the parasite were immersed separately for 30 minutes. By using the formula $OP = CRT$ (C = molar concentration ; R = gas constant = 0.082; and T = Absolute temperature = $273^{\circ} + \text{lab temperature}$) the osmotic pressures of the host and the parasite were determined. The osmotic pressure differences between the host and the parasite were calculated in terms of atmospheres. From the data presented in Table 1, it is evident that invariably in all the cases, there is higher

osmotic pressure in the parasite, as compared with the host, the difference in the pressure being very low (4.87 atm.) to very high (22.27 atm.), depending upon the host. Another interesting conclusion emerging from the present study is that there exists a narrow range in the osmotic pressure of the parasites on different host plants, with an average OP of 32.14 atmospheres (range : 29.47 to 36.91 atm.). It is also likely that this difference might well represent cases of physiologically distinct strains.

TABLE 1

Name of the host	Osmotic pressure in atmospheres		
	Host	Parasite	Difference
<i>Annona squamosa</i> Linn.	.. 15.38	30.80	15.42
<i>Ficus religiosa</i> Linn.	.. 18.56	32.08	13.52
<i>Cordia rothii</i> R. & S.	.. 8.38	30.65	22.27
<i>Citrus aurantium</i> Linn.	.. 21.76	34.61	12.85
<i>Mangifera indica</i> Linn.	.. 18.54	33.32	14.78
<i>Mimusops hexandra</i> Roxb.	.. 15.35	30.75	15.40
<i>Psidium guajava</i> Linn.	.. 12.34	33.32	20.98
<i>Oncoba spinosa</i> Linn.	.. 16.90	36.91	20.01
<i>Crescentia cujete</i> Linn.	.. 19.81	29.52	9.71
<i>Punica granatum</i> Linn.	.. 24.60	29.47	4.87

DEPARTMENT OF BOTANY,
GOVT. ARTS COLLEGE,
KARUR-639 005.

C. NARAYANASAMY

DEPARTMENT OF BOTANY,
ANNAMALAI UNIVERSITY,
ANNAMALAINAGAR-608 101,

R. SAMPATHKUMAR

November 29, 1979.

REFERENCES

FISCHER, C. E. C. (1926) : Lorantheae of Southern India and their host plants. *Rec. Bot. Surv. India* 11 (1) : 159-195.

SAMBANDAM, C. N. (1966) : Some new combinations of *Loranthus longiflorus* Desr. and host species. *Annamal Univ. Agric. Mag.* 6 : 63-64.

SAMPATHKUMAR, R. AND KUNCHITHAPATHAM, J. (1969) : Observations on the host range in *Loranthus longiflorus* Desr. *J. Bombay nat. Hist. Soc.* 65 (3) : 804-805.

————— (1970) : Further studies on the host range in *Loranthus longiflorus* Desr. *J. Bombay nat. Hist. Soc.* 67 (2) : 360-361.



Narayanasamy, C and Sampathkumar, R. 1981. "Host Parasite Relationships in *Dendrophthoe falcata* *Loranthus longiflorus*." *The journal of the Bombay Natural History Society* 78, 192–193.

View This Item Online: <https://www.biodiversitylibrary.org/item/187326>

Permalink: <https://www.biodiversitylibrary.org/partpdf/151426>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.