216 JOURNAL, BOMBAY NATURAL HIST. SOCIETY, Vol. 65 (1)

I observed the bird for perhaps two hours altogether, in conditions of both full sunlight and overcast skies.

The south-west monsoon (light showers) arrived on June 7 in this area (Hyderabad Dist.), but I cannot say whether this may have had some influence in the occurrence of this western sea-coast bird so far inland.

SHAMSABAD, Hyderabad District, Andhra Pradesh, July 8, 1967.

GEORGE F. NEAVOLL

6. THE FEMALE OF MOLESWORTH'S TRAGOPAN TRAGOPAN BLYTHI MOLESWORTHI BAKER

(With a plate)


Molesworth's Tragopan, Tragopan blythi molesworthi Baker, has so far been known from only two specimens, both males. The holotype was taken by Capt. A. L. M. Molesworth at Dengan La (alt. c. 2438 m.), c. 27° 11' N. 92° 1' E., Scherechopka country, south-eastern Bhutan, on 31 March 1914. Ludlow (1944)¹ procured the second specimen from Shingkar (alt. c. 2591 m.), Louri District, eastern Bhutan (c. 64 km. north-west of Dengan La, the type-locality), on 1 April 1936. Its female has so far been unknown.

During a recent ornithological survey of Bhutan undertaken jointly with Dr. Sálim Ali since 1966, I have been able to collect a female Tragopan which clearly belongs to *Tragopan blythi*, but differs from the nominate subspecies in some important details. There can, therefore, be little doubt that my specimen represents *T. b. molesworthi*, and the first known female at that (Plate).

The specimen was taken by me above the Bulfai Pass (alt. c. 2621 m.), c. 27° 14' N. 91° 31' E., Manas Valley, eastern Bhutan, on 30 March 1966. It was shot while skulking in a patch of rhododendron forest with thick undergrowth of various herbs and shrubs, and a few scattered clumps of ringal bamboo. It was a lone example busily feeding at about 8.00 hours when my presence there disturbed it.

This female specimen differs from the females of the nominate subspecies in having the whole upper plumage darker, the blackish marks being larger and deeper coloured, grey-brown markings much deeper grey and less brown, ear coverts subtipped rufous, rufous-centred cheek-

¹ LUDLOW, F. (1944): The birds of south-eastern Tibet. Ibis 86: 380-381.



Biswas: Molesworth's Tragopan



Molesworth's Tragopan (Tragopan blythi molesworthi Baker). Female.



Biswas, Biswamoy. 1968. "The Female of Molesworth's Tragopan Tragopan Blythi Molesworthi Baker." *The journal of the Bombay Natural History Society* 65, 216–217.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/186224</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/152805</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.