MISCELLANEOUS NOTES

4. SIGHTING OF GREY-HEADED LAPWING VANELLUS CINEREUS (BLYTH) IN HYDERABAD, ANDHRA PRADESH, INDIA

R. SREEKAR¹ AND RUDRA RAM²

¹306, Shanthi Soudha Apts., Erramanzil, Hyderabad 500 082, Andhra Pradesh, India. Email: sreekar1988@gmail.com
²Asian Wildlife Foundation, G 11, HUDA Swarna Jayanthi Complex, Ameerpet, Hyderabad 500 016, Andhra Pradesh, India. Email: ramrudra@yahoo.com

A solitary Grey-headed Lapwing Vanellus cinereus was sighted at Hussain Sagar (17° 25' 7" N; 78° 28' 3" E) located in the centre of Hyderabad city, Andhra Pradesh, India, twice on January 26, 2008, once on January 27, 2008, and twice on January 29, 2008. It was seen on the northern shores of the lake abutting Sanjeevaih Park. Its grey head, yellow beak with black tip and white secondaries differentiated it readily from the other Lapwings (Red-wattled Lapwing Vanellus indicus), which were also present in the area. This is the first record of the Grey-headed Lapwing from Hyderabad. However, it has been reported earlier from Andhra Pradesh from the following locations:

1. Machlipatnam, Krishna district on February 17, February 23, March 13 and April 03 in 2001 (Conroy 2003).

2. Thatipudi Reservoir, Visakhapatanam district on February 07, 2003 (Aasheesh Pittie, pers. comm. email dated September 02, 2008).

The sighting in Hyderabad seems to be an inland extension of range for the Grey-headed Lapwing, which was earlier reported from the coasts of Andhra Pradesh.

REFERENCE

CONROY, C. (2003): Grey-headed Lapwing Vanellus cinereus seen around Machlipatnam, Krishna district, Andhra Pradesh, India. J. Bombay Nat. Hist. Soc. 100(1): 122-123.

5. OCCURRENCE AND BREEDING RECORD OF THE FOREST OWLET HETEROGLAUX BLEWITTI FROM YAWAL WILDLIFE SANCTUARY, MAHARASHTRA, INDIA

RUSHIKESH A. CHAVAN¹ AND KISHOR D. RITHE²

¹A/11 'Safalya', Veera Desai Road, Andheri (W), Mumbai 400 058, Maharashtra, India. Email: rushikeshc@gmail.com ²'Pratishtha', Bharat Nagar, Akoli Road, Near Sainagar, Amravatti 444 607, Maharashtra, India. Email: kishor.rithe@gmail.com

The Forest Owlet *Heteroglaux blewitti* is one of the least known birds of India. It was considered extinct for 113 years, until its rediscovery on November 25, 1997, by Ben King, Pamela Rasmussen, and David Abbott in Toranmal Reserve Forest of Shahada in Nandurbar district, Maharashtra (King and Rasmussen 1998; Rasmussen and Collar 1998). The Forest Owlet is protected under Schedule I of the Indian Wildlife (Protection) Act of 1972, and is listed as critically endangered.

We were working to prepare a management plan for the buffer zone of Yawal Wildlife Sanctuary in Jalgaon district of Maharashtra situated in the Satpuda Mountain range. The 177.52 sq. km Sanctuary was established in 1969, and is located between Melghat Tiger Reserve and Toranmal Reserve Forest. Since the habitat of Yawal WS is similar to both these reserves, which hold a population of the Forest Owlet (Ishtiaq and Rahmani 2000; Jathar and Rahmani 2004), we speculated the presence of this critically endangered bird in the area.

On May 21, 2004, we were travelling in our vehicle towards the western side of the buffer zone through compartment number 166 (21° 36' 94" N; 75° 53' 147" E). The area is under the jurisdiction of the Forest Development Corporation of Maharashtra (FDCM). The FDCM has carried out Teak *Tectona grandis* plantations in these compartments about eight years ago. The average height of the trees is about 8-10 m. There was very little undergrowth, probably due to the continuous contour trenching (soil and water conservation work) carried out by the FDCM through out the area.

At about 1715 hrs an owlet flew past in front of our vehicle. The plumage pattern and colour of the owlet seemed to be different. We could not locate the bird as it disappeared in the nearby forest. We went in the direction the Owlet flew, but could not locate it. So we decided to play the call of the

MISCELLANEOUS NOTES

Forest Owlet, recorded from Melghat Tiger Reserve by the second author; the call was verified with records at the Bombay Natural History Society. As soon as we played the call, the Owlet immediately responded and came closer to investigate. It sat on a leafless teak *Tectona grandis* tree for about 10 minutes. We continued playing the call and to our surprise at around 1725 hrs one more bird responded to the call. Therefore, in all two birds were present in the vicinity. One of them approached us, up to a distance of *c*. 150 m. We got an opportunity to take some photographs of this individual and carefully observe its plumage. The crown, nape and back were unspotted brown, sharply contrasting with the broadly blackish and white banded wings. The breast looked entirely dull brown, contrasting sharply with the white belly.

We observed another Forest Owlet later in the day close to the site where we first observed it. The belly of the owlet was distinctly blotched. The body colour appeared darker with a stumpy tail. This suggested that the owlet was an immature and that the earlier birds could have been one of the parent birds. Therefore, we conclude successful breeding of the Forest Owlet in Yawal Wildlife Sanctuary. We have been able to identify the major stress factors in Yawal WS, which would possibly affect the Forest Owlet population in the Sanctuary. The stressors are listed below:

- Clearing of land for agriculture within the Sanctuary, some nesting trees may have been affected.
- Infiltration of villagers from Madhya Pradesh for collection of fire wood and timber.
- Encroachment for agriculture and new settlements within the Sanctuary.
- Forest fires in the Sanctuary.
- Many proposed and existing minor irrigation dams in the last remaining open scrub forest.

ACKNOWLEDGEMENTS

We are very grateful to all the support staff that was present during the survey. Thanks are due to Dr. Girish Jathar and Dr. Asad R. Rahmani for spending their valuable time to comment on the draft of this note.

REFERENCES

KING, B.F. & P.C. RASMUSSEN (1998): The rediscovery of the Forest Owlet Athene (Heteroglaux) blewitti. Forktail 14: 51-53.

RASMUSSEN, P.C. & N.J. COLLAR (1998): Identification, distribution and status of the Forest Owlet *Athene (Heteroglaux) blewitti. Forktail* 14: 41-49.

ISHTIAQ, F. & A.R. RAHMANI (2000): Further information on status and

distribution of Forest Owlet Athene (Heteroglaux) blewitti. Forktail 16: 172-173.

JATHAR, G.A. & A.R. RAHMANI (2004): Ecological studies of the Forest Spotted Owlet *Athene (Heteroglaux) blewitti*. Technical report. Bombay Natural History Society, Mumbai, India. Pp. 77.

6. ULTRAMARINE FLYCATCHER FICEDULA SUPERCILIARIS IN KACHCHH, GUJARAT

J.K. TIWARI¹

¹Centre for Desert and Ocean, Village Moti Virani, Taluka Nakhtrana, Kutch 370 665, Gujarat, India. Email: cedoindia@yahoo.com

Pingleshwar temple area, on the southern coast of Kachchh, Gujarat, India, has perhaps the largest tall tree groves in the area. During a bird watching trip on October 30, 2008, I came across an Ultramarine Flycatcher *Ficedula superciliaris* in a tree grove at Pingleshwar (area 200 sq. m).

So far, the Ultramarine Flycatcher has been reported from other parts of Gujarat, but not from Kachchh (Ali 1945; Grimmett and Inskipp 1998; Kazmierczak 2000; Rasmusseen 2005). There are five records of this bird from Gujarat (Bakul Trivedi pers. comm.).

The earlier records were by Bakul Trivedi at Jessore Sloth Bear Sanctuary, Polo Vijaynagar and by Bharat Jethva at Indroda Park, Gandhinagar.

This sighting of the Ultramarine Flycatcher in Pingleshwar temple area is thus a new record for Kachchh.

REFERENCES

ALI, S. (1945): Birds of Kutch. Pp.1-175. Oxford University Press, Mumbai.

GRIMMETT, R., C. INSKIPP & T. INSKIPP (1998): Birds of Indian Subcontinent. 1st edition. London; Christopher Helm, A & C Black.

KAZMIERCZAK, K. (2000): A Field Guide to the Birds of India, Sri Lanka,

Pakistan, Nepal, Bhutan, Bangladesh and the Maldives. Om Book Service. New Delhi.

RASMUSSEEN, P.C. & J.C. ANDERTON (2005): Birds of South Asia. The Ripley Guide. 2 vols. Washington D.C. and Barcelona: Smithsonian Institution & Lynx Edicions.



Chavan, Rushikesh A and Rithe, Kishor D. 2010. "Occurrence and Breeding Record of the Forest Owlet Heteroglaux Blewitti from Yawal Wildlife Sanctuary, Maharashtra, India." *The journal of the Bombay Natural History Society* 106, 207–208.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/188386</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/154899</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.