REFERENCES

CHOUDHURY, A.U. (1994): The decline of the wild water buffalo in northeastern India. *Oryx* 28(1): 70-73.

CHOUDHURY, A. U. (1997a): Checklist of the mammals of Assam 2nd edn. Gibbon Books & ASTEC, Guwahati.

> 5. OCCURRENCE OF INDIAN PANGOLIN MANIS CRASSICAUDATA IN ALIGARH, UTTAR PRADESH

Neither of the two pangolins reported from the Indian subcontinent are known to occur in Aligarh area. The Indian pangolin Manis crassicaudata is reported from peninsular India, while the Chinese pangolin M. pentadactyla occurs in northeastern India. Therefore, it was quite surprising that a pangolin was sighted in Aligarh (27° 29' to 28° 11' N and 77° 29' to 28° 39' E) in May 1997. The animal was captured and subsequently killed by the inhabitants of a local village called Barola Jafrabad 3 km from the Aligarh Muslim University (AMU) campus. The pangolin was identified as Manis crassicaudata from the measurements noted below by the AMU Museology Department where it is now on display.

6. STRANDED WHALES ON THE GUJARAT COAST

In this note, we report six stranded whales seen between 1983 to 1998. We had personally seen four specimens in Kutch district. There are reports of cetaceans, dolphins, porpoises and whales in the waters of Gujarat. Whales, when seen dead in deep waters, are brought to the coast by inquisitive fishermen, or by the natural force of tides. The reports of such dead whales in the newspapers attract hundreds of people to see these gigantic marine mammals.

S.no.	Place	Date & year	Remarks
1.	Asharmata	April 1983	15.24 m
2.	Sindhrodi	March 1984	15.24 m
3.	Kalumbhar Island	August 1988	8.38 m*
4.	Bhadreshwar coast	February 1977	13.72 m
5.	Ganga creek	July 1997	6.10 m
6.	Jakhau coast	July 1998	4.27 m

*Blue Whale

CHOUDHURY, A.U. (1997b): Survey of grasslands in some parts of central and southern Assam: to assess their biodiversity and socio-economic problem. Unpubl. Report. WWF-India NE Regional Office, Guwahati.

IN ALIGARH, UTTAR PRADESHolins reported from
known to occur inDate of collection15.v.1997Total length
Length of head105 cm10.5 cm

rotarionBin	1050111	
Length of head	10.5 cm	
Tail	50.5 cm	
Forelimb	8.5 cm + 5.5 cm (claw)	
Hindlimb	7 cm + 0.75 cm (claw)	
Width:		
Centre	48 cm	
Shoulder	47.5 cm	
Lower (hind) Portion	50 cm	

I thank Dr. Abdur Raheem, Lecturer, Department of Museology, AMU, for providing detailed measurements.

July 27, 2000 H.S.A. YAHYA Department of Wildlife Sciences, Aligarh Muslim University, Aligarh 202 002, Uttar Pradesh, India.

It seems that most of the stranded whales seen in Gujarat coast are the blue whales *Balaenoptera musculus*; sometimes sperm whales *Physeter catodon* may also be observed, e.g. specimen No. 6 seen on Jakhau coast.

The cause of death of the whales is not known. Interestingly, three specimens out of the six were seen from February to April and the remaining from July to August.

April 8, 2000	J.K. TIWARI
· · · · · · · · · · · · · · · · · · ·	Sea Water Farms Eritrea,
P.O. Box 40	06, Massawa, Eritrea, E. Africa
	S.N. VARU
Juna	vas, Temple Street, Madhapur,
ĸ	utch 370 020, Gujarat, India.



Tiwari, J. K. and Varu, Shantilal N. 2001. "Stranded Whales on the Gujarat Coast." *The journal of the Bombay Natural History Society* 98, 272–272.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/189534</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/155176</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.