

Hesperomys arviculoides, Pict.* *H. olivaceo-brunneus*, nigro-adspersus, subtus pedibusque canus, cauda dimidii corporis longitudine, dense pilosa; vellere nitido molli.—Corpus 5" 5^{'''}, cauda 2" 8^{'''}.
 Brasilia. Mus. Monac.

Hesperomys orobinus, Wagn. *H. supra brunneo-flavus*, subtilissime nigro-adspersus, subtus flavido-albescens, auriculis postice nudis, pedibus fuscis; cauda corpore multum brevior, squamata, nigricante.—Corpus 5", cauda 3" 5^{'''}.
 Brasilia. Mus. Monac.

Hesperomys subflavus, Wagn. *H. supra brunneo-flavidus*, subtus albidus, pedibus sordide albidis; cauda longitudine corporis, squamata, raripilosa.—Corpus 6", cauda 6".
 Brasilia. Mus. Monac.

Dasyprocta nigricans, Natt. *D. tota e nigro alboque variegata*, tergo concolore.—Corpus 22".
 From Borba on the Madaïro, and from Rio Negro above Cocuy.

VII.—*Descriptions of Chalcidites discovered by C. DARWIN, Esq.*
 By FRANCIS WALKER, Esq., F.L.S.

Thoracantha Latreillii, Fem. *Nigro-cyanea*, antennæ piceæ, pedes fulvi, femora fusca, alæ limpidæ.

Thoracantha Latreillii, Guérin, *Icon. Règne Anim., Insectes*, pl. 67.

Corpus gibbum, altum, compressum, incurvum, convexum, nigro-cyaneum, nitens, glabrum: caput parvum, transversum, declive, subtilissime striatum; vertex sat latus; frons impressa, ad os angustior: oculi rufi, mediocres, non extantes: ocelli 3, remoti, rufi, triangulum fingentes; medius perparum antepositus: antennæ piceæ, parvæ, graciles, extrorsum crassiores, apice fulvæ, capite vix longiores; articulus 1^{us} longus, sublinearis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, approximati; clava conica, articulo 10^o duplo longior: thorax maximus: prothorax brevissimus; sternum parvum; dorsum minimum, supra non conspicuum: mesothoracis scutum transverse striatum, longitudine latius, per medium obsolete sulcatum; parapsidum suturæ bene determinatæ, postice approximatae; scutellum transversum, breve, subtilissime rugulosum, per medium obsolete sulcatum: metathoracis dorsum maximum, subtilissime rugulosum, quasi bipartitum; pars antica longitudine vix latior, per medium obsolete sulcata; pars postica longior, fissa; fissura postice dehiscens, similitudine *Melvidis* elytrorum, abdominis apicem attingens: petiolus vix conspicuus: abdomen compressum, apice subtilissime rugulosum, basi læve, thorace paullo brevius et angustius; segmentum 1^{um} dorsum et sternum omnino obtegens: pedes fulvi, graciles, simplices, subæquales; coxæ nigro-cyaneæ; trochanteres picei; femora fusca; tarsi articuli 1^o ad 4^{um} curtantes, 5^{us} 4^o paullo longior; ungues et pulvilli minuti: alæ limpidæ, parvæ; squamulæ fulvæ; nervi albi, vix conspicui, basi fulvi; nervus humeralis ulnari multo longior, radialis vix ullus, cubitalis brevissimus; stigma minimum. (Corp. long. lin. 2; alar. lin. 3.)

Brazil.

* I have retained the name of *Mus. arviculoides*, Pictet, which was written on the label of the specimen when purchased.

Micromelus Silanus, Fem. *Æneo-niger, cyaneo-varius, antennæ fulvæ, pedes fulvi, alæ fuscæ.*

Corpus æneo-nigrum, subtus cyaneo-nigrum, breve, latum, convexum, parum nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace latius; vertex latus; frons impressa, abrupte declivis: oculi picei, mediocres, non extantes: ocelli vertice triangulum fingentes; medius perparum antepositus: antennæ fulvæ, ad os insertæ, basi graciles lineares: os piceum: thorax ovatus: prothorax brevissimus: mesothoracis scutum longitudine latius; parapsides scuto fere in unum confusæ; paraptera et epimera magna; scutellum subrhombiforme, postice latius: metathorax brevis, declivis, postice angustus: petiolus brevissimus: abdomen subrotundum, æneum, nitens, læve, supra planum, subtus carinatum, thorace paullo latius et multo brevius: pedes fulvi, simplices, subæquales: alæ fuscæ, mediocres; squamulæ piceæ; nervi fusci; nervus humeralis ulnari duplo longior, radialis ulnari brevior cubitali vix longior, cubitalis sat longus; stigma mediocre. (Corp. long. lin. $\frac{3}{4}$; alar. lin. $1\frac{1}{4}$.)

Mount Wellington.

VIII.—*Descriptions of Chalcidites discovered in St. Vincent's Isle by the Rev. Lansdown Guilding. By FRANCIS WALKER, Esq., F.L.S.*

Decatoma Oretilia, Mas et Fem. *Fulvo-flava ferrugineo-varia, antennæ fulvæ, pedes flavi, alæ limpidæ.*

Corpus convexum, sublineare, fulvo-flavum, nitens, subtilissime punctatum, parce hirtum: caput transversum, breve, thorace paullo latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ fulvæ, clavatæ, thorace non longiores; articulus 1^{us} longus, gracilis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 9^{um} latescentes; clava fusiformis, acuminata, articulo 9^o multo latior et plus duplo longior: thorax ovatus: prothorax maximus, transversus, subquadratus, postice ferrugineo-maculatus: mesothoracis scutum brevissimum; parapsidum suturæ vix conspicuæ; scutellum subrotundum; discus ferrugineus: metathorax transversus, brevis: petiolus brevis: abdomen ovatum, læve, *mas* thoracis longitudine, *fem.* thorace paullo longius subtus carinatum apice acuminatum fuscum; vitta per dorsum interrupta fusca: pedes pallide flavi, simplices, subæquales: alæ limpidæ; squamulæ fulvæ; nervi fulvi; nervus humeralis ulnari fere duplo longior, radialis ulnari non brevior cubitali plus duplo longior; stigma minutum. (Corp. long. lin. 1; alar. lin. $1\frac{1}{2}$.)

Pteromalus Helice, Fem. *Æneo-viridis, antennæ fulvæ, pedes flavi, alæ limpidæ.*

Corpus convexum, æneo-viride, nitens, scitissime squameum, parce hirtum: caput transversum, breve, thorace latius; vertex latus; frons impressa, abrupte declivis: oculi rufi, mediocres, non extantes: antennæ fulvæ, extrorsum crassiores, thorace vix longiores; articulus 1^{us} longus, gracilis; 2^{us} cyathiformis; 3^{us} et 4^{us} minimi; 5^{us} et sequentes breves, usque ad 10^{um} latescentes; clava longiconica, acuminata, articulo 10^o duplo longior: thorax ovatus: prothorax transversus, brevissimus: mesothoracis scutum longitudine latius; parapsidum suturæ vix conspicuæ; scutellum subconicum, antice truncatum: metathorax obconicus, mediocris, declivis: petiolus brevissimus: abdomen fusiforme, læve, supra depressum, subtus carinatum, apice acuminatum, thorace longius et angustius: pedes læte flavi, simplices, subæquales: alæ limpidæ; squamulæ fulvæ; nervi flavi; nervus humeralis



Walker, Francis. 1843. "Descriptions of Chalcidites discovered by C. Darwin, Esq." *The Annals and magazine of natural history; zoology, botany, and geology* 12, 45–46. <https://doi.org/10.1080/03745484309442484>.

View This Item Online: <https://www.biodiversitylibrary.org/item/19370>

DOI: <https://doi.org/10.1080/03745484309442484>

Permalink: <https://www.biodiversitylibrary.org/partpdf/15764>

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.