

CHROMOSOME NUMBERS OF SISYRINCHIUM (IRIDACEAE) IN EASTERN NORTH AMERICA¹

ROYCE L. OLIVER AND WALTER H. LEWIS

Stephen F. Austin State College, Nacogdoches, Texas

Chromosome numbers for 18 species of *Sisyrinchium* with basic complements of $x = 8, 9$, and 17 , have been reported (Vilmorin and Simonet, 1927; Maude, 1940; Bowden, 1945; Covas and Schnack, 1946; Sermonet, 1948; Skottsberg, 1953; Lewis and Oliver, 1961). The 8 and 9 series contain both diploid and high polyploid species, but only diploid species are found in the secondary $x = 17$ series.

MATERIALS AND METHODS.—Immature flower buds were collected in the field and were fixed in modified Carnoy's solution (4: 3: 1). As soon as possible after fixation the buds were stored at refrigerator temperatures for periods up to 8 months; only after 7 months was deterioration noted in some cells. Buds were squashed in 1% acetic-orcein and PMCs or more occasionally somatic cells were studied. Representative chromosomes were drawn with the aid of a camera lucida at X2300. The first set of voucher specimens are filed at the Southern Methodist University herbarium (SMU). Field work for this study was supported in part by the National Science Foundation, G-9800.

For one collection seeds were soaked in water until the radicle appeared and then sown on a culture medium (Lewis and Elvin-Lewis, 1961) to facilitate more rapid growth. After pretreatment for 1-2 hours in supersaturated paradichlorobenzene, the seedlings were fixed and stained following the procedure used for the buds and mitosis was studied in root tips and apical meristems.

OBSERVATIONS.—Fifteen species and one hybrid of *Sisyrinchium* from a total of 50 populations are listed in Table 1 with their chromosome numbers and voucher data. For 11 species the collections represent a wide range of morphological variation including atypical individuals, but in each case the chromosome numbers were found to be consistent. Chromosome numbers for 9 species are reported for the first time. These include *S. minus* Engelm. & Gray, $n = 5$; *S. fibrosum* Bickn., $n = 8$; *S. campestre* Bickn., $n = 16$; *S. intermedium* Bickn., $n = 16$; *S. laxum* Otto, $n = 16$; *S. mucronatum* Michx., $n = 16$; *S. sagittiferum* Bickn., $n = 16$; *S. arizonicum* Roth., $2n = 34-36$; *S. atlanticum* Bickn., $n = 48$. Those for the remaining species, *S. albidum* Raf., *S. bermudiana* L. (as *S. angustifolium* Mill.), *S. ensigerum* Bickn., *S. langloisii* Greene, *S. mi-*

¹ EDITOR'S NOTE. It was not until after this paper went to press that I was able to reach a conclusion about the correct names of the introduced annuals. I believe that *S. laxum* of this paper is correctly *S. rosulatum* Bicknell, and *S. micranthum* is rather *S. exile* Bicknell. See "Annual *Sisyrinchiums* (Iridaceae) in the United States," this issue, pp. 32-42. —L. H. Shinnars.

cranthum Cav., and *S. pruinatum* Bickn., agree with counts by Bowden (1945) and Lewis and Oliver (1961).

The number for *S. minus* Engelm. & Gray adds a new basic number of $x = 5$ to the genus (Fig. 1, 2). Although an undetermined diploid species in the $x = 8$ series has been reported from South America (Bowden, 1945), the number of $n = 8$ for *S. fibrosum* Bickn. (Fig. 3) is the first report of a diploid North American species in this series. All other species studied, with the exception of *S. arizonicum* Roth., occur in the $x = 8$ series at either the tetraploid or the dodecaploid level. Only a tentative count of $2n = 34$ or 36 is reported for *S. arizonicum* Roth.

There is little difference in chromosome size for most species of *Sisyrinchium* (Fig. 1-9), although *S. bermudiana* L. (Fig. 11) has larger chromosomes than *S. atlanticum* Bickn. (Fig. 10) in the same basic series.

Meiotic "irregularities" were rarely observed except for the extreme bunching of chromosomes. This phenomenon was observed for most collections and consequently only a small proportion of the meiotic metaphase and anaphase plates could be accurately interpreted. Despite this, pollen were usually normal in appearance except for one collection from 1.5 miles west of the Neches River and Highway 94, Trinity Co., Texas (Oliver, 312). In a sample of 100 pollen grains from each of several plants, pollen was non-staining, appeared shriveled, and micropollen were frequent. Meiosis was not observed, but the number of microspores per PMC at the tetrad stage, and frequency based on a random sample of 100 PMCs, was 4 microspores (6%), 5 microspores (26%), 6 microspores (36%), 7 microspores (12%), 8 microspores (12%), 9 microspores (4%), and 10 microspores (4%). With only 6% normal tetrad formation, meiosis was probably highly irregular. These plants are morphologically intermediate between *S. laxum* Otto and *S. micranthum* Cav., which were both growing in the immediate vicinity, and are assumed to be hybrids between these species.

LITERATURE CITED

- BOCHER, T. W., AND K. LARSEN 1950. Chromosome numbers of some arctic or boreal flowering plants. Medd. Gronland 147: 6.
- BOWDEN, W. M. 1945. A list of chromosome numbers in higher plants. I. Acanthaceae to Myrtaceae. Amer. Journ. Bot. 32: 81-92.
- COVAS, G., AND B. SCHNACK. 1946. Numero de cromosomas en Antofitas de region de cuyo (Republica Argentina). Rev. Argent. Agron. 13: 153-166.
- LEWIS, W. H., AND M. ELVIN-LEWIS. 1961. Medium for growing small rubiaceous seeds from herbarium material. Castanea 26: 146-155.
- _____, AND R. L. OLIVER. 1961. Meiotic chromosomes in six Texas and Mexican *Nemastylis* and *Sisyrinchium* (Iridaceae). Southwest. Nat. 6: 45-46.
- MAUDE, P. F. 1940. Chromosome numbers in some British plants. New Phytol. 39: 17-32.
- SERMONTI, G. 1948. Osservazioni sul kariogramma di *Sisyrinchium striatum*. Caryologia 1: 79-82.
- SKOTTSBERG, C. 1953. Chromosome numbers in Hawaiian flowering plants. Ark. f. Bot. 3: 63-70.
- VILMORIN, R., AND M. SIMONET. 1927. Nombre des chromosomes dans les genres *Lobelia*, *Linum*, et chez quelques autres especes vegetales. Compt. Rend. Soc. Biol. 96: 166-168.

Table 1. CHROMOSOME NUMBERS OF SISYRINCHIUM IN
EASTERN NORTH AMERICA.

Taxon	Gametic No.	Voucher
<i>S. minus</i> Engel. & Gray	5	TEXAS. Nacogdoches Co., Nacogdoches, <i>Oliver</i> 317 (tepals purple), <i>Oliver</i> 318 (tepals white); Brazos Co., 10.7 miles SE of College Station, <i>Oliver</i> 296.*
<i>S. fibrosum</i> Bickn.	8	GEORGIA. Pike Co., 0.5 miles N of Pike Co.-Upson Co. line on Hwy. 19, <i>Oliver</i> 327.
<i>S. albidum</i> Raf.	16	LOUISIANA. Bossier Par., 5.6 miles E of Bossier City, <i>Oliver</i> 272†. TEXAS. Nacogdoches Co., Fern Lake Fire Tower, <i>Oliver</i> 243†, 0.6 miles S of Nacogdoches, <i>Oliver</i> 241†.
<i>S. campestre</i> Bickn.	16	ARKANSAS. Ouachita Co., 7 miles SW of Y city, <i>Oliver</i> 302. KANSAS. Wayne Co., 1.7 miles E and 1 mile S of Wayne, <i>Morley</i> , 12 May 1961; Republic Co., 1 mile E and 0.7 miles N of Belleville, <i>Morley</i> , 13 May 1961.
<i>S. ensigerum</i> Bickn.	16	TEXAS. Hays Co., 1 mile S of San Marcus, <i>Osborne</i> 39; Lampasas Co., 8.9 miles S of Lampasas, <i>Osborne</i> 42; Wilson Co., 10 miles S of Stockdale, <i>Osborne</i> 38.
<i>S. intermedium</i> Bickn.	16	TENNESSEE. Polk Co., 1 mile N of Hwys. 64 and 30 junction, <i>Oliver</i> 325*.
<i>S. langloisii</i> Greene	16	TEXAS. Brazoria Co., 3.7 miles S of Freeport, <i>Lewis</i> 5571, 4 miles S of Freeport, <i>Lewis</i> 5569B*; Chambers Co., 2.3 miles SW of Hwys. 121 and 87 junction, <i>Oliver</i> 252; Panola Co., Lake Murvaul, <i>Oliver</i> 276; San Augustine Co., 3 miles E of Attoyac River on Hwy. 21, <i>Oliver</i> 248.
<i>S. laxum</i> Otto	16	TEXAS. Angelina Co., 5 miles S of Lufkin, <i>Oliver</i> 310; Jasper Co., Hwys. 96 and 1004 junction, <i>Lewis</i> 5626; Nacogdoches Co., Stripling Island, <i>Oliver</i> 304; Newton Co., 4.7 miles S of Newton, <i>Lewis</i> 5618.



Figs. 1-11. Meiotic chromosomes of *Sisyrrinchium* originally drawn at X2300 and reduced by ca. 28% in reproduction. Fig. 1. *S. minus*, $n = 5$, Oliver 296. Fig. 2. *S. minus*, $n = 5$, Oliver 318. Fig. 3. *S. fibrosum*, $n = 8$, Oliver 327. Fig. 4. *S. campestre*, $n = 16$ (one side of metaphase II), Oliver 302. Fig. 5. *S. campestre*, $n = 16$, Morley, 13 May 1961. Fig. 6. *S. pruinatum*, $n = 16$, Lewis 5615. Fig. 7. *S. sagittiferum*, $n = 16$, Oliver 241. Fig. 8. *S. sagittiferum*, $n = 16$, Oliver 269. Fig. 9. *S. intermedium*, $n = 16$, Oliver 325. Fig. 10. *S. atlanticum*, $n = 48$, Oliver 253. Fig. 11. *S. bermudiana*, $n = 48$, Oliver 292.

<i>S. micranthum</i> Cav.	16	TEXAS. Angelina Co., 5 miles S of Lufkin, <i>Oliver</i> 309; Hardin Co., 1.4 miles SW of Votaw, <i>Lewis</i> 5630; Trinity Co., 1 mile N of Neches River, <i>Oliver</i> 311.
<i>S. mucronatum</i> Michx.	16‡	CANADA. ONTARIO. Bruce Co., Oliphant, <i>Heimbürger</i> , 17 June 1961.
<i>S. pruinatum</i> Bickn.	16	ARKANSAS. Hempstead Co., 1 mile SW of Hope, <i>Lewis</i> 5615†. TEXAS. Aransas Co., 4.2 miles SW of Aransas Pass, <i>Lewis</i> 5591, 0.2 miles S of Aransas Wildlife Refuge Headquarters, <i>Lewis</i> 5597, 1 mile W of Aransas Wildlife Refuge Headquarters, <i>Lewis</i> 5592, 1 mile NE of Rockport, <i>Lewis</i> 5575†; Ellis Co., 1 mile N of Italy, <i>Oliver</i> 281; Henderson Co., 5.3 miles W of Hwys. 175 and 31 junction, <i>Osborne</i> 48; Lampasas Co., 4 miles W of Lampasas, <i>Osborne</i> 45; Matagorda Co., 4.7 miles NE of Markham, <i>Lewis</i> 5572; San Patricio Co., 0.2 miles W of Welder Wildlife Foundation Headquarters, <i>Lewis</i> 5580†; Van Zandt Co., 3 miles SE of Wills Point, <i>Oliver</i> 278†; Wharton Co., Louise, <i>Lewis</i> 5599.
<i>S. sagittiferum</i> Bickn.	16	LOUISIANA. La Salle Par., 2.5 miles SE of Gene, <i>Oliver</i> 269. TEXAS. Nacogdoches Co., 10 miles S of Nacogdoches, <i>Oliver</i> 237†; Orange Co., 3.8 miles SW of Orange, <i>Oliver</i> 255.
<i>S. laxum</i> X <i>micranthum</i>	16‡	TEXAS. Trinity Co., 1 mile N of Neches River and Hwy. 94, <i>Oliver</i> 312.
<i>S. arizonicum</i> Roth.	17-18‡	MEXICO. DURANGO. 19 miles SE of Durango, <i>Waterfall</i> 15541 (SMU).
<i>S. atlanticum</i> Bickn.	48	TEXAS. Chambers Co., 1 mile NE of Hwys. 121 and 87 junction, <i>Oliver</i> 253*, 5 miles NE of Ferry Landing, <i>Oliver</i> 251*.

* Chromosomes of 3 plants examined; otherwise the number is based on the study of 1 plant.

† Atypical collection.

‡ Chromosome number from somatic cells.

S. bermudiana L.

48 LOUISIANA. West Feliciana Par., 10 miles S of La.-Miss. state line on Hwy. 61, *Oliver* 266. TEXAS. Nacogdoches Co., Goodman's Bridge over Angelina River, *Oliver* 308, 1 mile NE of Nacogdoches, *Oliver* 316, 10 miles S of Nacogdoches, *Oliver* 292.



Oliver, Royce L and Lewis, Walter H. 1962. "CHROMOSOME NUMBERS OF SISYRINCHIUM (IRIDACEAE) IN EASTERN NORTH AMERICA." *SIDA, contributions to botany* 1, 43–48.

View This Item Online: <https://www.biodiversitylibrary.org/item/38228>

Permalink: <https://www.biodiversitylibrary.org/partpdf/162700>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Botanical Research Institute of Texas

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.