

**BRICKELLIA SONORANA (ASTERACEAE), A NEW SPECIES FROM  
MÉXICO BELONGING TO THE SUBGENUS PHANEROSTYLIS**

B. Turner

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

ABSTRACT

A new annual species, *Brickellia sonora* B. Turner, is described from Sonora, México. It belongs to the *Barroetea* complex of *Brickellia* and has been previously, but incorrectly, referred to as *Barroetea laxiflora* [= *Brickellia laxiflora* (T. Brandege) B. Turner, the type from Puebla, México]. The Barroeteoid complex of *Brickellia* is composed of four closely related species: *B. sonora*, *B. subuligera* (Schauer) B. Turner, *B. laxiflora* and *B. pavonii* (A. Gray) B. Turner. A key to these taxa is presented and a map showing their distributions is provided.

KEY WORDS: *Brickellia*, *Barroetea*, Eupatorieae, Asteraceae, México

Preparation of a treatment of *Barroetea* for the Asteraceae of México has convinced me that this genus, phyletically speaking, belongs under the taxonomic fabric of *Brickellia* (cf. Turner, *et al.* 1991). In connection with this study it has proved necessary to describe the following novelty.

***Brickellia sonora* B. Turner, sp. nov.** TYPE: MÉXICO. Sonora: San Bernardo, Río Mayo, hill slope in short-tree forest, 21 Feb 1935, *Howard Scott Gentry 1326* (HOLOTYPE: TEX!; Isotypes: ARIZ!, LL!).

*Brickelliae laxiflorae* (T. Brandege) B. Turner similis sed capitulescentia laxa, pedunculis ultimis plerumque 2-6 cm longis (vs. 0.1-0.5 cm) differt.

Stiffly erect annual herbs 30-70 cm high, arising from rather small, shallow, tap roots. Stems terete, striate, densely puberulent to glabrate. Leaves opposite throughout, mostly 1.5-5.0 cm long, 1-3 cm wide; petioles 5-20 mm long; blades triangular to somewhat subcordate, trinervate from the base, sparsely to moderately puberulent, sparsely to densely glandular punctate beneath, the



margins serrate, ciliate, most of the serrations with callous, usually ciliate, appendages. Heads few to numerous, purplish, arranged in very lax leafy cymes or cymelike panicles, the ultimate peduncles mostly 2-5(7) cm long. Involucres mostly 8-10 mm high, the bracts 4-8 seriate, graduate, linear-lanceolate, 2-5 striate, puberulent to glabrous, the apices narrowly acute to apiculate. Receptacle plane, epaleate. Florets 15-21, the corollas tubular, purplish above, ca. 5 mm long, the lobes 5, ca. 0.5 mm long, usually with 1 or more sessile glandular hairs. Style branches with linear-ob lanceolate purplish appendages which are essentially smooth at magnification of 20x. Achenes tangentially flattened, the abaxial surface with a single principal nerve, the adaxial surface with 3 principal nerves (lesser nerves sometimes intercalated between the principal nerves on the abaxial surface); pappus of 20-25 sparsely barbellate bristles 4-6 mm long.

ADDITIONAL SPECIMENS EXAMINED: MÉXICO. Sinaloa: Culiacán, 14 Nov (w/o year), *Brandegge s.n.* (UC); 23 mi N of Culiacán, 28 Jan 1964, *Flyr 110* (TEX); 4 mi S of turnoff to La Cruz on México highway 15, 29 Jan 1964, *Flyr 121* (TEX); Imala, 20 Nov 1939, *Gentry 5001* (ARIZ); Cerro Colorado, 4 Feb 1940, *Gentry 5485* (ARIZ); Cerro Tecomate, 29 Feb 1940, *Gentry 5765* (ARIZ); 59 mi N of Culiacán, 22 Dec 1971, *Norris, et al. 20113* (LL). Sonora: SW edge of Alamos, 22 Sep 1973, *Fish 134* (TEX); Canyon Sapopa, Río Mayo, 15 Oct 1934, *Gentry 1040* (F); Río Mayo, 21 Feb 1935, *Gentry 1326* (ARIZ); Alamos, 16 Oct 1936, *Gentry 2923* (ARIZ); Sierra Alamos, 3 Nov 1939, *Gentry 4864* (ARIZ); 109° 19' x 28° 16', 26 Mar 1983, *Sanders 3697* (ARIZ); Sierra Alamos, 30 Dec 1983, *Van Devender 83-119* (ARIZ, TEX).

This taxon was included within *Barroetea laxiflora* T. Brandegge by Robinson (1911) in his revision of *Barroetea*. That name, however, is typified by *Purpus 4128* (HOLOTYPE: UC!) from Coxcatlán, Puebla, a markedly different taxon with a lax capitulescence of congested heads and having cordate leaves with more numerous serrations. I have transferred this species and related taxa to *Brickellia* (Turner, et al. 1991).

A key to the Barroeteoid species of *Brickellia*, along with complete synonymy and a map showing distributions are presented below.

#### KEY TO SPECIES OF THE BARROETEA COMPLEX OF BRICKELLIA

1. Stems with glandular trichomes; denticulations of leaves not setose. ....  
..... *B. problematica*
1. Stems glabrous or puberulent, without glandular trichomes; denticulations of leaves setose. .... (2)
  2. Leaves sessile or nearly so. .... *B. pavonii*
  2. Leaves with petioles mostly 3-30 mm long. .... (3)



3. Peduncles of individual heads mostly 2-6 cm long; Son, Sin. *B. sonora*
3. Peduncles of individual heads mostly 0-2 cm long; widespread. .... (4)
4. Achenes with 1-2(3) well developed ribs on adaxial faces; florets 15-27 per head; midstem leaves with petioles 1-3 cm long; se Pue, Oax. .... *B. laxiflora*
4. Achenes without well developed ribs on adaxial faces; florets 30-50 per head; midstem leaves with petioles 0.5-1.0 cm long; widespread. .... *B. subuligera*

*Brickellia laxiflora* (T. Brandege) B. Turner, *Phytologia* 71(1):48. 1991. BASIONYM: *Barroetea laxiflora* T. Brandege, *Univ. Calif. Publ. Bot* 4:93. 1910. TYPE: MÉXICO. Puebla: Coxcatlán, Sep 1909, C.A. Purpus 4128 (HOLOTYPE: UC!).

*Barroetea brevipes* B.L. Robins., *Proc. Amer. Acad. Arts* 47:205. 1911. TYPE: MÉXICO. Oaxaca: Valley of Cd. Oaxaca, 5200-6800 ft, 2 Oct 1894 (HOLOTYPE: GH!; Isotype: US!).

As indicated in Fig. 1, this species is found in central Veracruz, southern Puebla and northern Oaxaca, mostly occurring from 500-1500 m, and flowering from October-November.

Robinson distinguished *Barroetea brevipes* from *B. laxiflora* by its reportedly sessile heads arranged in rather laxly branched panicles, but branching patterns of the capitulescence in *B. laxiflora* is very variable, as evident from the ten or more collections of this taxon (LL, TEX) examined by the present author, most of these relatively recent collections.

Chromosome number,  $2n = 18$ .

*Brickellia pavonii* (A. Gray) B. Turner, *Phytologia* 71(1):48. 1991. BASIONYM: *Barroetea pavonii* A. Gray, *Proc. Amer. Acad. Arts* 17:206. 1882. TYPE: MÉXICO. w/o locality, 1787-1804, M-Pavon Herb., Sesse, et al. s.n. (numbered 1499 and labeled "*Eupatorium setigerum*;" Photoholotype: F!, TEX!). Gray also cited a collection in the Pavon herbarium annotated as "*Eupatorium cuspidatum*," which is numbered 4053 (photograph F!, TEX!). Both sheets bear plants with sessile leaves and clearly belong to the same taxon.

*Brickellia pavonii* is a commonly encountered weedy species, largely confined to tropical deciduous forests of southcentral México (Fig. 1), occurring from near sea level to about 2200 meters, flowering from October-November. Numerous collections are on deposit at LL and TEX.

Chromosome number,  $2n = 18$  (Sundberg 2971, TEX).






Figure 1. Distribution of the Barroeteoid species of *Brickellia*.

*Brickellia problematica* B. Turner, *Phytologia* 71(1):48. 1991. Based upon *Barroetea glutinosa* T. Brandege, *Zoe* 5:262. 1908. Not *Brickellia glutinosa* A. Gray, *Proc. Amer. Acad. Arts* 21:385. 1886.

This species was originally described as *Barroetea glutinosa* and complete synonymy is given in Turner, *et al.* (1991).

*Brickellia problematica* is largely confined to southernmost Puebla (Fig. 1) and adjacent northern Oaxaca where it occurs in relatively xeric woodlands dominated by species of *Juniperus*, *Eysenhardtia*, *Montanoa*, *Croton*, *Agave*, etc., occurring mostly from 1800-2400 m; flowering from August to November. While originally thought to be rare, recent collections (seven or more) show the species to be locally common in the mountainous areas about Caltepec, Puebla and elsewhere in this region, depending upon rains.

*Brickellia subuligera* (S. Schauer) B. Turner, *Phytologia* 71(1):48. 1991. BASIONYM: *Bulbostylis subuligera* S. Schauer, *Linnaea* 19:718. 1847. *Barroetea subuligera* (S. Schauer) A. Gray, *Proc. Amer. Acad. Arts* 15:29. 1879. TYPE: MÉXICO. Hidalgo: Zimapan, w/o date, *Aschenborn* 260 (HOLOTYPE: B, destroyed?; Isotype: P, according to McVaugh [1984]).

*Barroetea setosa* A. Gray, *Proc. Amer. Acad. Arts* 15:29. 1879. TYPE: MÉXICO. San Luis Potosí: mountains southeast of the city of San Luis Potosí, 1830-2440 m, Jan-Feb 1878, *Parry & Palmer* 353 (HOLOTYPE: GH!).

*Barroetea subuligera* (S. Schauer) A. Gray var. *latisquamea* Greenm., *Proc. Amer. Acad. Arts* 40:35. 1904. TYPE: MÉXICO. Jalisco: hills above Etzatlán, 27 Oct 1903, *C.G. Pringle* 8773 (HOLOTYPE: MO; Isotypes: LL-[2]!).

This is by far the most widespread variable species within the *Barroetea* complex, occurring mostly on the Central Plateau of México in dry deciduous forests from 500-2000 m; flowering from September-January.

Chromosome number,  $2n = 18$  (*Barrie* 1106, TEX).

#### ACKNOWLEDGMENTS

I am grateful to Dr. Guy Nesom for the Latin diagnosis, and to him and Dr. A.M. Powell for reviewing the paper.



## LITERATURE CITED

- McVaugh, R. 1984. *Barroetia*, in *Flora Novo-Galiciana* 12:116-119.
- Robinson, B.L. 1911. Revision of the genus *Barroetia*. *Proc. Amer. Acad. Arts* 47:202-206.
- Turner, B.L., K.-J. Kim, & J. Norris. 1991. Taxonomic status of *Barroetia glutinosa* (Asteraceae, Eupatorieae) and its allies: Morphological evidence for the transfer of *Barroetia* to *Brickellia*. *Phytologia* 71(1):38-50.

*Phytologia* (July 1991) 71(1):56.

NEW NAME FOR *SENECIO GESNERIFOLIUS* TURNER (SENECIONEAE,  
ASTERACEAE)

B.L. Turner

Department of Botany, University of Texas, Austin, Texas 78713 U.S.A.

## ABSTRACT

*Senecio mezquitalanus* Turner is provided as a replacement for *S. gesnerifolius* B. Turner which is a later homonym of *S. gesnerifolius* Cuatrecasas.

KEY WORDS: *Senecio*, Senecioneae, Asteraceae, nomenclature

*Senecio mezquitalanus* B. Turner, *nom. nov.*, a new name for *S. gesnerifolius* B. Turner. BASIONYM: *Senecio gesnerifolius* B. Turner, *Phytologia* 62:75. 1987. Not *Senecio gesnerifolius* Cuatrecasas, *Fieldiana Bot.* 27:33. 1950.

## ACKNOWLEDGMENTS

I am grateful to Dr. C. Jeffery of KEW for calling the above oversight to my attention. He also informs me that he intends to position the species concerned under the genus *Roldana* H. Robins. & Brett.



Turner, B. L. 1991. "Brickellia sonorana (Asteraceae), a new species from Mexico belonging to the subgenus *Phanerostylis*." *Phytologia* 71, 51–56.  
<https://doi.org/10.5962/bhl.part.16699>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/47375>

**DOI:** <https://doi.org/10.5962/bhl.part.16699>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/16699>

#### **Holding Institution**

New York Botanical Garden, LuEsther T. Mertz Library

#### **Sponsored by**

The LuEsther T Mertz Library, the New York Botanical Garden

#### **Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.