

THE EARLIEST TISSOTIID AMMONITE

by W. J. KENNEDY and O. H. BAYLISS

ABSTRACT. *Pseudotissotia* (*Pseudotissotia*) *inopinata* sp. nov. is described from the Upper Cenomanian phosphatic *Calycoceras naviculare*/*Eucalycoceras pentagonum* Zone fauna of Division C of the Cenomanian Limestone of Shapwick Grange, Devon. It is the earliest occurrence of the otherwise exclusively Turonian genus *Pseudotissotia* Peron, 1897, and the Turonian to Coniacian Family Tissotiidae Hyatt, 1900. The family is otherwise unknown in the Cretaceous of the United Kingdom.

THE ammonite family Tissotiidae Hyatt, 1900, comprises approximately twenty genera with a widespread, largely Mesogean distribution. The group as a whole is characterized by a tendency towards loss of ornament, by the development of an entire, simple keel or keels, involute coiling, and a suture line which commonly shows a tendency towards multiplication of sutural elements and over-all simplification. *Tissotia* H. Douvillé, 1878, and *Buchiceras* Hyatt, 1875, are perhaps the best-known genera, and are classic Cretaceous pseudoceratites (e.g. Hyatt 1903), homoemorphous with Triassic ammonoids in their development of entire saddles and denticulate lobes. The group is chiefly known from the Tethyan Realm, especially North and East Africa, the Middle East, Central and South America, Texas and Spain (Bayle 1878; Douvillé 1912; Peron 1897; Hyatt 1903; Reyment 1954*a, b*, 1955; Barber 1957; Collignon 1958, 1965*a*; Freund and Raab 1969; Choffat 1898; Benavides-Cáceres 1956; see palaeogeographic maps in Matsumoto 1973*a, b*). They occur rarely in the U.S. Western Interior (e.g. Cobban and Scott 1972) and are also known from as far south as Madagascar (e.g. Collignon 1965*b*) and Borneo. In Western Europe the family becomes progressively rarer as one moves northward, the most northerly record hitherto being the occurrence of *Pseudotissotia* (*Pseudotissotia*) *galliennei* (d'Orbigny) in the Turonian of Sarthe and Touraine.

The previously known stratigraphic range of the tissotiids is Turonian to Coniacian. The actual dating of their first appearance in southern areas is difficult to relate to northern European faunal successions, but records by Freund and Raab (1969), Reyment (1955), Barber (1957), Wiedmann (1959), and others suggest it to be some way up in the Lower Turonian.

So far as evolutionary origins are concerned, Wright in 1952 (p. 221, n. 35) considered the family to be polyphyletic to a minor degree, with their source presumed to be in the Mammitinae. The work of Reyment (1954*a, b*, 1955) on the rich Nigerian faunas led him (1955, text-fig. 31) to suggest the subgenus *P.* (*Bauchioceras*) Reyment, 1954 as the rootstock of the group, whilst Barber (1957) suggested that this in turn derived from the vasococeratid *Gombeoceras* Reyment, 1954, during the early Turonian. Subsequently, Collignon (1965*a*, p. 179) suggested an alternative origin for the group in the vasococeratids, when he demonstrated that the early Turonian genus *Disco-vasoceras* Collignon, 1958 bore three keels when young, and could equally be regarded as intermediate between the two families.

It is therefore of great interest to record what appears to be a *P.* (*Pseudotissotia*) from the English Upper Cenomanian, apparently pre-dating the vasoceratids from which certain other tissotids have been claimed, or demonstrated, to be derived, especially as the family has never before been recorded from this country.

SYSTEMATIC DESCRIPTION

Superfamily ACANTHOCERATACEAE Hyatt, 1900

Family TISSOTIIDAE Hyatt, 1900

Subfamily PSEUDOTISSOTIINAE Hyatt, 1903

Genus and Subgenus PSEUDOTISSOTIA Peron, 1897

Type species. *Ammonites galliennei* d'Orbigny, 1850.

Pseudotissotia inopinata sp. nov.

Plate 120, figs. 1a-d, 2a-c, text-fig. 1

Derivation of name. Latin *inopinata*—unexpected.

Holotype. Formerly O. H. Bayliss Collection, no. 257; now B.M. (N.H.). No. C80436.

Diagnosis. A *Pseudotissotia* with a depressed whorl section (breadth to height ratio 1.16), strong umbilical bullae which project into the umbilicus and give rise to broad, subdued, triangulate ribs which efface at mid-flank, and a venter bearing three strong, equal keels.

Description. The holotype consists of a phosphatic internal mould of just over one-third of a whorl, is partially septate, and retains a considerable amount of phosphatized shell, which largely obscures the sutures.

The coiling appears to have been quite involute, with a moderately deep umbilicus; the umbilical wall is vertical, with an abruptly rounded shoulder. The whorl section is depressed, with a whorl breadth to height ratio of 1.16; the greatest breadth is at the umbilical bullae. The flanks are flattened and convergent, with rounded ventrolateral shoulders. The venter is broad (approximately 80% of maximum whorl breadth), and bears three distinct, entire, strong keels, the two lateral ones being rather more narrowly rounded than the siphonal. The keels are separated by quite deep, rounded grooves, of somewhat greater breadth.


Flank ornament consists of strong, rounded umbilical bullae which project slightly into the umbilicus. From these bullae arise broad, flat, triangulate ribs, which efface completely by mid-flank.

EXPLANATION OF PLATE 120


Figs. 1a-d, 2a-c. *Pseudotissotia* (*Pseudotissotia*) *inopinata* sp. nov. The holotype, Bayliss Collection no. 257, now B.M. (N.H.) no. C80436, a partly septate phosphatic internal mould retaining phosphatized shell.

Specimen from the phosphatic *Calycoceras naviculare*/*Eucalycoceras pentagonum* Zone fauna of Division C of the Cenomanian Limestone, Shapwick Grange, near Lyme Regis, Dorset. 1a-d, $\times 1$; 2a-c, $\times 2$.

Figs. 3a-c. *Pseudotissotia* (*Pseudotissotia*) *galliennei* (d'Orbigny). Specimen from the Turonian Tuffeau of Poncé, Sarthe, France. Muséum d'Histoire Naturelle, Paris, ex Bourgeois Collection, $\times 1$.



KENNEDY and BAYLISS, earliest tissotiid ammonite



TEXT-FIG. 1. Partial suture of the holotype of *Pseudotissotia* (*Pseudotissotia*) *inopinata* sp. nov. $\times 6$.

Little of the suture line can be deciphered, but there appears to have been a moderate degree of incision to both lobes and saddles (text-fig. 1).

Discussion. Identification of this fragment as a tissotiid is based upon whorl section, rib-style, and the prominent equal keels. Erection of a new species based on the fragment is justified by the distinctive ornament, and by the great age compared with most other members of the group.

Three subgenera have been distinguished within *Pseudotissotia*. Of these, *P. (Bauchioceras)* Reyment, 1954 has crenulate keels, and a siphonal keel which may be very feeble, together with ribs which extend across the whole of the flank at comparable diameters. Species such as *P. (B.) nigeriensis* (Woods), *P. (B.) tricarinata* (Reyment) are thus readily distinguishable from *P. (P.) inopinata*. *P. (Wrightoceras)* Reyment, 1954 has compressed outer whorls which lack a siphonal keel, whilst the umbilical region may be inflated. Species such as *P. (Wrightoceras) wallsi* (Reyment), *P. (W.) mirabilis* (Pervinqui re), *P. (W.) munieri* (Pervinqui re), *P. (W.) llareni* Karrenberg, and *P. (W.) gagnieri* Faraud may be distinguished from the present form by these criteria. When compared with typical *P. (Pseudotissotia)*, a specimen of the type species of which is figured for comparison (Pl. 120, fig. 3a-c), the attribution of our specimen to the restricted subgenus is quite clear, for it has three distinct, entire keels. When compared with *P. (P.) galliennei*, *P. (P.) inopinata* is depressed and broad, rather than high-whorled and compressed, whilst all specimens of *P. (P.) galliennei* we have seen completely lack umbilical bullae and ribs at comparable diameters; the long, low, fold-like ribs of the adult instead appear at much larger diameters, as in Peron's specimen (Peron 1897, p. 28, pl. 2, fig. 3; pl. 3, fig. 1). The suture of *P. (P.) inopinata* is, in addition, more divided than that of *P. (P.) galliennei*. Collignon (1957, p. 15) has described, but not figured, a var. *inflata* of *P. (P.) galliennei*, based on a rather worn specimen from the Turonian of Tinhret (Fezzan). It has a breadth:height ratio of 0.83, substantially less than the present species, but further comparison is difficult in the absence of illustrations of Collignon's variety.

Occurrence. The holotype and only known specimen of *Pseudotissotia inopinata* is from Division C of the Cenomanian Limestone exposed in the large chalk pit at Shapwick Grange Farm, Devon, near Lyme Regis, Dorset (National Grid Reference SY 313918).

Age. The sequence at Shapwick Grange is closely similar to that on the Devon Coast described by Kennedy (1970, p. 657; text-fig. 15) who also summarizes previous work. Large collections from the succession have enabled us to place the present specimen and associated material precisely in sequence. Division B of the Cenomanian Limestone at Shapwick has yielded scarce *Acanthoceras* of Middle Cenomanian age. This unit is terminated by a phosphatic veneer, and overlain by the glauconitic chalk of Division C; from the base of this comes a rich phosphatic fauna including, in addition to *P. (P.) inopinata*, the following ammonites in similar preservation (O. H. Bayliss Collection): abundant *Schloenbachia lymense* Spath, *Calycoceras* spp., including *C. (C.) naviculare* (Mantell), *C. (C.) guerangeri* (Spath), *C. (Lotzeites) aberrans* (Kossmat), *Eucalycoceras pentagonum* (Jukes-Browne), *E. rowei* (Spath), *Protacanthoceras* including *P. compressum* (Jukes-Browne) and *P. bunburianum* (Sharpe), *Thomelites* spp., including *T. aff. sornayi* (Thomel), *Forbesiceras* spp., *Hamites* cf. *simplex* (d'Orbigny) and abundant *Scaphites equalis* J. Sowerby. This is an Upper Cenomanian, *C. naviculare*/*E. pentagonum* Zone fauna.

Occurring somewhat higher in Division C is an essentially indigenous fauna preserved as glauconite-coated moulds. This includes *Sciponoceras gracile* (Shumard), *C. naviculare*, *Allocrioceras annulatum* (Shumard), *Pseudocalycoceras dentonense* (Moreman), and *Metoicoceras geslinianum* (d'Orbigny) (= *M. gourdoni* de Grossouvre). This fauna indicates the latest Cenomanian *S. gracile* Zone (for a discussion of zonal nomenclature see Kennedy and Juignet 1976, and Rawson *et al.* in press).

DISCUSSION

The recognition of a *Pseudotissotia* in the English Cenomanian would seem to confirm Wright's (1952) suggestion that the Tissotiidae are polyphyletic, for *Bauchioceras* appears to be a quite definite derivative of the early Turonian vascoceratid *Gombeoceras*, as Reyment (1955) and Barber (1957) demonstrated, whilst Collignon's *Discovascoceras* is a further contender as a vascoceratid ancestor for other tissotids. The origin of *P. inopinata* remains a problem. The morphologically closest contemporary ammonites appear to be amongst the Acanthoceratinae. Certain compressed *Acompsoceras* from the Lower to Middle Cenomanian develop a low siphonal keel and sharp ventral shoulders which are sometimes tuberculate, e.g. *Acompsoceras essendiense* (Schlüter) and *A. renevieri* (Sharpe) (Kennedy 1971, p. 67 *et seq.*, pl. 30, pl. 31, fig. 2). These forms may also bear umbilical bullae and ribs, which die out on the flank. Mr. C. W. Wright has also drawn our attention to certain compressed, feebly ornamented variants of *Acanthoceras rhotomagense* (Brongniart) which show similar flank ornament and develop an incipient siphonal keel; we would suggest, therefore that, within the Cenomanian acanthoceratid radiation, these forms were a possible source for *P. (P.) inopinata*.

Acknowledgements. We thank Dr. J. M. Hancock and Mr. M. R. Cooper for useful discussions, and Mr. C. W. Wright for advice, and for critically reviewing the manuscript of this article.

REFERENCES

- BARBER, W. 1957. Lower Turonian ammonites from north-eastern Nigeria. *Bull. geol. Surv. Nigeria*, **26**, 86 pp., 34 pls.
- BAYLE, E. 1878. Fossiles principaux des Terrains. *Explic. Carte géol. France*, **4** (1) (Atlas), 158 pls.
- BENAVIDES-CÁCERES, V. E. 1956. Cretaceous system in northern Peru. *Bull. Am. Mus. nat. Hist.* **108**, 357–498, pls. 31–66.
- CHOFFAT, P. 1898. Recueil d'études paléontologiques sur la faune crétacique du Portugal. 1, Espèces nouvelles ou peu connues. 2, Les ammonées du Bellasian, des Couches à *Neolobites vibrayeanus*, du Turonien et du Sénonien. *Trav. géol. Portugal*, **1898**, 41–86, pls. 3–22.
- COBBAN, W. A. and SCOTT, G. R. 1972. Stratigraphy and ammonite fauna of the Graneros Shale and Greenhorn Limestone near Pueblo, Colorado. *Prof. Pap. U.S. geol. Surv.* **645**, 108 pp., 41 pls.
- COLLIGNON, M. 1958. Céphalopodes néocrétacés du Tihret (Fezzan). *Annls Paléont.* **43**, 115–136, pls. 16–18.
- 1965a. Nouvelle ammonites néocrétacés Saharienne. *Ibid.* **51**, 165–202, pls. A–H.
- 1965b. *Atlas des Fossiles Caractéristiques de Madagascar* (Ammonites) fasc. **12** (Turonien), pp. 1–82, pls. 376–413. Service Géologique, Tananarive.
- DOUVILLÉ, H. 1912. Évolution et classification des Pulchelliidés. *Bull. Soc. géol. Fr.* (4) **11**, 285–320.
- FREUND, R. and RAAB, M. 1969. Lower Turonian ammonites from Israel. *Spec. Pap. Palaeont.* **4**, 83 pp., 10 pls.
- HYATT, A. 1903. Pseudoceratites of the Cretaceous. *Monogr. U.S. Geol. Surv.* **44**, 351 pp., 47 pls.
- JUIGNET, P. and KENNEDY, W. J. 1977. Stratigraphie comparée du Cénomaniien du sud d'Angleterre et de la Haute Normandie. *Bull. Soc. Géol. Normandie et Amis du Muséum du Havre* (in press).
- KENNEDY, W. J. 1970. The correlation of the uppermost Albian and the Cenomanian of south-west England. *Proc. Geol. Ass.* **81**, 613–677.
- 1971. Cenomanian ammonites from Southern England. *Spec. Pap. Palaeont.* **8**, 133 pp., 64 pls.
- PERON, A. 1896–1897. Les ammonites du Crétacé Supérieur de l'Algérie. *Mem. Soc. géol. Fr.* **6** (1896), 1–24, pls. 1–6; **7** (1897), 25–88, pls. 7–18.
- RAWSON, P. *et al.* *A correlation of the Cretaceous Rocks of the British Isles*. Geological Society of London (in press).
- REYMENT, R. A. 1954a. New Turonian (Cretaceous) ammonite genera from Nigeria. *Colon. Geol. Min. Resour.* **4**, 149–164, 4 pls.
- 1954b. Some new Upper Cretaceous Ammonites from Nigeria. *Ibid.* 248–270, 5 pls.
- 1955. The Cretaceous Ammonoidea of southern Nigeria and the southern Cameroons. *Bull. Geol. Surv. Nigeria*, **25**, 112 pp., 25 pls.
- WIEDMANN, J. 1959. Le Crétacé Supérieur de l'Espagne et du Portugal et ses céphalopodes. *C.R. Congrès des Sociétés Savantes-Dijon 1959: Colloque sur le Crétacé supérieur français*, 709–764, 8 pls.
- WRIGHT, C. W. 1952. A classification of Cretaceous ammonites. *J. Paleont.* **26**, 213–222.
- 1957. In ARKELL, W. J., *et al.* *Treatise on Invertebrate Palaeontology*, Part L, **4**, 490 pp., 558 figs. University of Kansas Press and Geological Society of America. Lawrence, Kansas and New York.

W. J. KENNEDY

Department of Geology and Mineralogy
Parks Road
Oxford

O. H. BAYLISS

Spring House
Uplyme
Dorset

Typescript received 12 May 1976

Revised typescript received 19 July 1976



Kennedy, W. J. and Bayliss, O. H. 1978. "The earliest tissotiid ammonite."
Palaeontology 20, 901–906.

View This Item Online: <https://www.biodiversitylibrary.org/item/197363>

Permalink: <https://www.biodiversitylibrary.org/partpdf/173404>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.