

THE PALAEOECOLOGY OF THE GONIATITE BED AT COWLOW NICK, CASTLETON, DERBYSHIRE

by TREVOR DAVID FORD

ABSTRACT. The Cowlow Nick Goniatite Bed is shown to be an accumulation of randomly oriented, hollow, or spar-filled goniatite shells of several species in a matrix of calcilutite. Of very limited dimensions, the bed is surrounded by algal limestones of the fore-reef facies of Upper B₂ age (Lower Carboniferous). Suggestions as to the mode of accumulation are discussed and it is concluded that the bed represents a specialized drifted assemblage of floating shells which were washed gently into an inactive surge channel or submarine cave. Some comparable occurrences elsewhere in Derbyshire are noted.

THE small bed of limestone crowded with goniatites occurring in Cowlow Nick, west of Castleton in Derbyshire (Nat. Grid Ref. SK/142824) is well-known to workers on the Carboniferous Limestone. It was first recorded by Bisat (1934) who described species found there by G. B. Alexander. Subsequently it was included in the Castleton Limestones (of reef facies) by Shirley and Horsfield (1940) who included these in the B₂ sub-zone, resting unconformably against 'cliffs' of the massif-facies D zone limestones. Parkinson (1947) agreed in placing the Cowlow Nick beds in the B₂ sub-zone, but thought that this reef facies was equivalent in age to the massif-facies limestones to the south which are of D₁ age. Both Hudson and Cotton (1945) and Parkinson (1947) discussed further subdivision of the B₂ sub-zone, but as there is some discrepancy between the results of these authors and with the indications of the fauna listed herein, there seems to be some revision needed.

None of the above workers discussed the palaeoecology of the Goniatite Bed except as a very broad interpretation of all the pockets of fossils in the reef limestones as drifted assemblages. Wolfenden (1958) described the palaeoecology of the whole reef complex of B₂/D₁ age by analysing the distribution of the various elements of the fauna in terms of fore-reef, reef, back-reef, and shelf biotopes. The existence of a basinal biotope was noted, but not analysed. Goniatites were noted as being almost confined to the fore-reef facies (and basin by inference). Wolfenden challenged the drifted-assemblage concept of Hudson and Cotton (1945) and said that as the goniatites occurred throughout the fore-reef in all sizes and were well preserved, and were not often transported out of that biotope, they were probably preserved where they lived—a life assemblage. Wolfenden, however, did not discuss the Cowlow Nick Goniatite Bed nor did he mention localized accumulation of goniatites to the almost total exclusion of other members of the biota.

It is the purpose of this short communication to analyse the composition of the fauna in the Cowlow Nick Goniatite Bed, to examine its relationship with surrounding deposits and to attempt to draw conclusions about the mode of accumulation.

THE FAUNA

The material collected by Alexander and described by Bisat (1934) included: *Beyrichoceras submicronotum* Bisat, *B. vesciculiferum* (de Koninck), *Goniatites maximus* Bisat, [Palaeontology, Vol. 8, Part 1, 1965, pp. 186–91, pl. 20.]

Dimorphoceras gilbertsoni (Phillips), *Sagittoceras acutum* Hind (= *discus* Roemer), *Pronorites cyclolobus* (Phillips), *Prolecanites discoides* Foord and Crick.

Shirley and Horsfield (1940), Hudson and Cotton (1945), and Parkinson (1947) failed to add to this, though other species were found in the Castleton Limestones at the nearby localities of Cavedale and Treak Cliff. Wolfenden's list (1958) noted facies distribution, but not individual localities.

Further collecting by the writer and by Dr. J. R. L. Allen has yielded: *B. rectangularum* Bisat, *G. cf. hudsoni* Bisat, *P. serpentinus* Phillips, *P. cf. P. discoides* Foord and Crick, ? *Discitoceras* sp., 'Orthoceras' spp.

A sample of the bed was broken up and the following abundances of the various species were noted:

<i>B. submicronotum</i>	29
<i>B. vesciculiferum</i>	2
<i>B. rectangularum</i>	8
<i>G. maximus</i>	29
<i>G. cf. hudsoni</i>	1
<i>D. gilbertsoni</i>	1
<i>P. serpentinus</i>	1
<i>P. cf. P. discoides</i>	3
'Orthoceras' spp.	6
? <i>Discitoceras</i> sp.	1
	—
Total	81

Other samples produced much the same proportions, showing a slight preponderance of *Beyrichoceras* spp. over *Goniatites* spp. Owing to the difficulties of extraction of the goniatites in an undamaged condition (particularly of young forms) no reliable size frequency distributions could be recorded. However, both polished and thin sections show all sizes to be present, although adults are predominant with only a few young. The sizes of the adults in the different species groups differ considerably. In the Beyrichoceratids the diameters range from 15 mm. to 20 mm. whilst in the *Goniatites* spp. the range is from 18 mm. to 25 mm. Orthoceratids are scattered throughout and vary up to 100 mm. in length and 20 mm. diameter.

The Goniatic Bed has only a few other macrofossils, except for a small lens of young brachiopods near the roof. This is seen only for about 30 cm. in length and up to 10 cm. thick, and is composed of young *Dielasma hastata*, all much the same size, about 7–8 mm. in length. A few productids are scattered through the bed, and also a few gastropods, some of which show traces of original colouring. The microfossils present are bryozoan fragments, and very occasional young brachiopods, foraminifera, and ostracods.

STRATIGRAPHICAL RELATIONSHIPS

The Cowlow Nick Goniatic Bed reaches a maximum of nearly 1 metre thickness in the mouth of an old lead-vein working, now partly collapsed. As far as can be judged from the irregular bedding it has a dip of some 15° NE. which is rather less than the hill slope and considerably less than the dips in the neighbouring outcrops of reef-complex

limestones, which range between 20° and 40° within a few hundred metres. As a result of these relationships, the Goniatile Bed disappears into the hillside on the up-dip side, whilst down dip its base is lost by the steep hill-slope. Along the strike exposures are discontinuous and the bed is lost beneath the turf only 3 m. to either side of the lead vein. The vein is, however, oblique to the strike and in the north-west wall the bed appears to wedge out in both directions, with about 3.5 m. between the extremities. The nearest exposures in all directions are calcite-mudstones of the reef-complex frequently showing algal banding, and with pockets of reef-brachiopods. Lenses of crinoidal debris appear lower down the hill slope.

Thus the detailed stratigraphical relationships of the bed are obscure, but in the general sense it may be said that it occurs in steeply dipping reef-complex limestones, which by projection south-eastwards of Wolfenden's (1958) mapping in the Winnats Pass are within the fore-reef facies, close to the reef wall.

Fortunately within the lead-vein opening parts of the roof and floor of the bed are visible, though only for a short distance laterally. The roof is composed of calcite-mudstone with algal banding. Immediately beneath this is the lens of young brachiopods in a matrix of calcite-mudstone. A metre below is the floor of the Goniatile Bed, which is composed of either a fragmental limestone with small crinoid ossicles, shell-fragments and calcite-mudstone fragments or of calcite-mudstone with algal banding. These are seen only for a length of about 0.5 m. and to a similar depth below the goniatile bed, and within this grain-size is variable up to about 8 mm.

PETROGRAPHY


The goniatites in the Cowlow Nick bed are of varied sizes as discussed above. They are randomly oriented in a fine-grained calcite-mudstone matrix. The shells are almost undamaged and the ornament is not worn. Body chambers can be seen in sections, but can rarely be extracted undamaged. No evidence of size sorting has been obtained.

Diagenetic alterations obscure much of the detail and the following features have been observed: (a) frequent veinlets of calcite and occasional fluorite cutting both goniatites and matrix; (b) patches of calcite spar, some with carbonaceous impurities, sometimes surrounded by algal reef 'tufa'; (c) fine calcite crystalline mosaic in the matrix obscuring details of original depositional textures; (d) recrystallization of parts of the shells to form either coarse calcite spar or fine crystal mosaic, and (e) scarce authigenic quartz crystals and patches of limonite staining. In places brecciation appears to be associated with the mineral vein.

The goniatile shells are mostly filled with coarse calcite spar. A few are still hollow or with only some chambers hollow and lined with calcite crystals. In a few cases there are crystals of fluorite or blebs of bitumen in either the spar filling or in the hollow shells. Some shells, which appear to be broken on a polished section, are seen in thin section

EXPLANATION OF PLATE 20

Polished vertical section of the Cowlow Nick Goniatile Bed. Approximately twice natural size. The section shows the random orientation of goniatile shells of various sizes in a calcilutite matrix. Both primarily damaged and diagenetically recrystallized shells are present. Cavity fillings of calcite, fluorite, and carbonaceous matter are also present.



FORD, Cowlow Nick goniatite bed

to pass into the fine crystal mosaic of the matrix. No evidence has been found of original sedimentary filling of the shells other than in body chambers.

The fine grained matrix appears to be largely due to diagenetic recrystallization, but in places traces of original textures remain, chiefly irregular algal banding.

PALAEOECOLOGY

From the above it may be surmised that the Goniatite Bed is an assemblage of several species of goniatites, of varying sizes, preserved virtually undamaged and randomly oriented in a lenticular bed of restricted dimensions in the reef-complex close to the probable position of the algal reef-wall. In many of its characters the Bed compares with the well-known reef-brachiopod beds of the reef-complex on Treak Cliff, &c. Three possible explanations may now be examined:

- A. The Goniatites were gregarious sessile benthonic animals, not, as usually supposed, free-swimmers, and they were preserved as a 'life-assemblage', possibly by some catastrophe.
- B. The Goniatites were, as usually supposed, free-swimming reef-grazers, which at times of storm, &c., retired to a sheltered hollow in the reef, where they were overwhelmed by a catastrophe.
- C. The Goniatites are an unusual drifted assemblage, for which some mechanism must be found to account for a concentration of floating shells in the midst of largely algal limestones.

The first of these is difficult to uphold as it is completely in contrast to most opinion of the habits of ammonoids. Also one would expect such sessile gregarious colonies to have been preserved much more frequently. The same species are found in shale facies elsewhere associated with planktonic and epi-planktonic faunas, and they are also found as scattered individuals in association with reef brachiopods elsewhere in the reef complex. Finally, gregarious sessile benthonic associations are usually formed of a single species.

The possibility B seems more likely at first as it does not require any unusual means for concentrating shells if they retired to this shelter of their own accord. However, one would again expect to find such occurrences more often. If the goniatites had retired to a sheltered hollow during some time of stress only to be trapped there by some catastrophe such as the collapse of flanking deposits, it would be expected that the shells would be crushed together, not dispersed at random through the Bed. Also such a sheltered hollow might be expected to be populated with a sessile fauna before being occupied by the goniatites.

The third possibility C, that of a special type of drifted assemblage, must now be examined, and a perusal of literature on modern reefs has given a suggestion. In the Bikini Atoll Reports, Emery and others (1954, part A pl. 51, and part C figs. 92 and 93) have shown diagrams of surge channels in the active reef edge, and they have described there the gradual closure and abandonment of such surge channels by algal growth over the top and outwards around the mouths of the channels, leaving submarine caves. It is thus possible that the Cowlow Nick Goniatite Bed represents a concentration of goniatite shells in a surge channel, in a Lower Carboniferous reef complex. The fact that



Ford, Trevor David. 1965. "The palaeoecology of the goniatite bed at Cowlow Nick, Castleton, Derbyshire." *Palaeontology* 8, 186–191.

View This Item Online: <https://www.biodiversitylibrary.org/item/196343>

Permalink: <https://www.biodiversitylibrary.org/partpdf/174535>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.