Phytologia (July 1995) 79(1):22-24.

THE STATUS OF QUERCUS ARKANSANA SARG. (FAGACEAE) IN TEXAS

David M. Hunt

New York Natural Heritage Program, 700 Troy-Schenectady Road, Latham, New York 12110-2400 U.S.A.

&

Michael H. MacRoberts and Barbara R. MacRoberts

Bog Research, 740 Columbia, Shreveport, Louisiana 71104 U.S.A.

ABSTRACT

The status of *Quercus arkansana* Sarg. in Texas is discussed. It is known to occur in Cass County, and there is an historical record for Jasper County.

KEY WORDS: Quercus arkansana, Texas, Fagaceae, phytogeography

Quercus arkansana Sarg., an uncommon Coastal Plain oak with scattered populations ranging from southwestern Georgia and northwestern Florida to southwestern Arkansas and northwestern Louisiana, has not been known for Texas (Bill Carr, Texas Natural Heritage Program, pers. comm.; Correll & Johnston 1970; Johnston 1990; Stanley D. Jones, Botanical Research Center (BRCH), pers. comm.; Hunt 1986; Little 1977; Nixon 1985; Sargent 1965; Simpson 1988; Vines 1977). Hunt (1990) lists the species as occurring in Texas on the basis of an historical specimen collected in 1903 in Jasper County (W.W. Ashe 1 [NCU]). More recently, two small populations of Q. arkansana have been found in Cass County.

In 1990, Hunt (*Hunt TX218* [to be distributed to TEX and ND]) first confirmed *Quercus arkansana* from Texas, identifying it as "*Q. arkansana* tending toward *Q. nigra.*" This collection, representing one sapling, was from the south side of TX 77, 1 km west of the Louisiana state line in sandy loam pine-oak woods dominated by *Pinus echinata* P. Mill., *Quercus falcata* Michx., *Q. nigra* L., and *Liquidambar styraciflua* L. This find was the result of searches stimulated by the presence of a large population in Louisiana about 3 km away, documented initially by G.H. Ware (*Ware 1492* [DAV]) in 1965 (Hunt 1990).

In 1995, the MacRobertses found several small trees (up to 7 meters high) in a young, densely stocked, even-aged pine plantation on deep sandy soils along a 1 km

stretch of Cass County Road 4561 about 5 km northeast of McLeod, Texas, and 5 km west of Rodessa, Louisiana, which they identified as *Quercus arkansana* (*MacRoberts & MacRoberts 2875* [NCSC], 2874, 2877 [BRCH], 2879 [VDB], 2881 [LSUS], 2873 [ND], 2878 [DAV]). Some of these specimens were sent elsewhere for confirmation. Hunt determined 2876 as "probably *Quercus arkansana* × *Q. nigra;*" 2875 and 2873 were identified as *Q. arkansana* by J.W. Hardin and Richard Jensen, respectively. Julia Larke determined an unnumbered specimen to be *Q. arkansana*. This population is within 2 km of a large population of *Q. arkansana* in Louisiana centering on a point where Black Bayou and State Line Creek cross LA 168 about 3 km west of Rodessa (Teague & Wendt 1994). Associated species for this site include *Quercus marilandica* Muenchh., *Q. stellata* Wang., *Q. incana* Bartr., *Pinus taeda* L., *P. echinata*, Vitis spp., Sassafras albidum (Nutt.) Nees, and Liquidambar styraciflua. The area was heavily shaded, had a dense pine litter and almost no herbaceous understory, and was badly damaged by commercial forestry.

The environment of the Cass County sites is typical for the species: midslope of eroding sandhills adjacent to headwater tributaries. That *Quercus arkansana* occurs in Cass County is not surprising since it occurs in adjacent parishes and counties in Louisiana and Arkansas (Hunt 1990; Louisiana Natural Heritage 1995; Smith 1988). Hunt (1986) has documented increased introgression between Q. arkansana and Q. nigra at the range edge of the former. Thus the mixed traits found in the Cass County collections are expected: short petioles (Q. nigra), scurfy leaf and twig pubescence (Q. arkansana), and leaf shape varying between the two provide strong evidence for putative hybridization between these species.

In addition to these recent finds, Hunt located a specimen from Jasper County (W.W. Ashe 1 [NCU]) collected in 1903 that is probably Quercus arkansana but may be a hybrid between Q. nigra and Q. velutina Lam. (Hunt 1990). This specimen was originally identified as Q. velutina \times Q. laurifolia Michx. Unfortunately, a specific locality is not given and, although much potential mesic sandhill habitat exists in the northern third of the county, relocating this population, if it still exists, may be difficult. The east Texas landscape has been drastically altered over the past century, and Q. arkansana is notorious for its patchy distribution and its inconspicuousness, which has always made it difficult to locate (Hunt 1986).

These populations represent westward range extensions for *Quercus arkansana*. Additional populations of this oak are expected and should be sought in Bowie, Marion, and Cass counties. The rarity of this species in Texas and throughout its range makes it a good candidate for any protected plant list for the state, and an effort to protect its habitat should be made to ensure its survival in Texas.

ACKNOWLEDGMENTS

Stanley Jones (BRCH) and Bill Carr (Texas Natural Heritage Program) supplied information on the distribution of *Quercus arkansana*. J.W. Hardin (NCSC), Richard Jensen (ND), and Julia Larke (Louisiana Natural Heritage Program), confirmed the MacRobertses' initial identification. Hunt's research was supported by NSF Grant BSR-8414419 to David E. Giannasi and David M. Hunt. Thanks are due to Lori

Hunt and Michael Moore for assistance in the field, and Nancy Coile for handling herbarium loans. Richard Jensen and Stanley Jones reviewed the manuscript.

LITERATURE CITED

Correll, D.S. & M.C. Johnston. 1970. Manual of the Vascular Plants of Texas. Texas Research Foundation, Renner, Texas.

Hunt, D.M. 1986. Distribution of *Quercus arkansana* in Georgia. Castanea 51:183-187.

Hunt, D.M. 1990. A Systematic Review of *Quercus* Series *Laurifoliae*, *Marilandicae* and *Nigrae*. Ph.D. dissertation, University of Georgia, Athens, Georgia.

Johnston, M.C. 1990. The Vascular Plants of Texas. A list, up-dating the Manual of the Vascular Plants of Texas. Published by the author, Austin, Texas.

Little, E.L. 1977. Atlas of United States Trees. Vol. 4. Minor Eastern Hardwoods. Misc. Publ. 1342. USDA Forest Service, Washington D.C.

Louisiana Natural Heritage Program. 1995. Rare plant species of Louisiana. Unpublished report. Louisiana Department of Wildlife and Fisheries, Baton Rouge, Louisiana.

Nixon, E.S. 1985. Trees, Shrubs, & Woody Vines of East Texas. Bruce Lyndon Cunningham Productions, Nacogdoches, Texas.

Sargent, C.S. 1965. Manual of the Trees of North America. Dover Press, New York, New York.

Simpson, B.J. 1988. A Field Guide to Texas Trees. Texas Monthly Press, Austin, Texas.

Smith, E.B. 1988. An Atlas and Annotated List of the Vascular Plants of Arkansas. Published by the author.

Teague, J. & T. Wendt. 1994. Caddo and Bossier Parishes, LA: Natural areas survey. Unpublished Report. The Nature Conservancy. Baton Rouge, Louisiana Vines, R.A. 1977. *Trees of East Texas*. University of Texas Press, Austin, Texas.



Hunt, David M, Macroberts, Michael H , and Macroberts, Barbara R . 1995. "The status of Quercus arkansana Sarg. (Fagaceae) in Texas." *Phytologia* 79, 22–24.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/183041</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/175289</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Biodiversity Heritage Library

Copyright & Reuse Copyright Status: In Copyright. Digitized with the permission of the rights holder Rights Holder: Phytologia License: <u>http://creativecommons.org/licenses/by-nc/3.0/</u> Rights: <u>https://www.biodiversitylibrary.org/permissions/</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.