

BLAINVILLEA BRASILIENSIS BLAKE TRANSFERRED TO
CALYPTOCARPUS (ASTERACEAE).

B.L. Turner

Dept. of Botany, University of Texas, Austin, TX 78713
U.S.A.

H. Robinson (1978, *Phytologia* 41:34. 1978), in his transfer of Blainvillea biaristata DC. into Calyptocarpus, alluded to "the transfer of all of the subgenus Oligogyne from Blainvillea to Calyptocarpus.", commenting thereafter that "A second South American species, C. bahiensis (DC.) Sch.-Bip. has previously been placed in the genus."

Robinson was apparently unaware that the latter is synonymous with the earlier Blainvillea brasiliensis (Nees & Mart.) Blake, as noted by Blake (1925, *Proc. Biol. Soc. Wash.* 38:85.). Thus the following new combination would appear to be in order:

CALYPTOCARPUS BRASILIENSIS (Nees & Mart.) B. Turner, comb. nov. based upon Galophthalmum brasiliense Nees & Mart. in Wied, *Nov. Act. Acad. Leop.-Carol. Nat. Cur.* 12:8. pl. 2, 1824.



Turner, B. L. 1988. "BLAINVILLEA-BRASILIENSIS BLAKE TRANSFERRED TO CALYPTOCARPUS ASTERACEAE." *Phytologia* 64, 214–214.

View This Item Online: <https://www.biodiversitylibrary.org/item/46821>

Permalink: <https://www.biodiversitylibrary.org/partpdf/176071>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.