

Additions to the List of Newton Noctuidæ.

The following names are to be added to those given in the list on p. 34-39 of the present volume of *PSYCHE*; the abbreviations are the same as in that list.

- | | |
|---|--|
| <p><i>Diphtera fallax</i>. R. l. Jl.
 <i>Apatela Radcliffei</i>. R.; raised. Jn.
 Lrv. Prunus, Sp.
 <i>Agrotis badicollis</i>. R. l. Au.
 <i>fennica</i>. Several s. Sp.
 <i>albipennis</i>. Two s. l. Sp.
 <i>incivis</i>. R.; three s. Sp.
 <i>trabalis</i>. Cocoon found under bark of <i>Pinus strobus</i> in April.
 Imago appeared in June.
 <i>Hadena impulsæ</i>. R. l. Jl.
 <i>apamiformis</i>. R. l. Jl. Au.
 <i>vultuosa</i>. C. l. Au.
 <i>leucoscelis</i>. R. s. Au.
 <i>Dryobota stigmata</i>. R. s. Sp.
 <i>Laphygma frugiperda</i>. C. s. Sp.
 <i>Prodenia flavimedia</i>. R. s. Sp.
 <i>lineatella</i>. R. l. Jl. [<i>P. ornithogalli</i> is to be stricken from the previous list.]
 <i>Gortyna limpida</i>. R. l. Sp.
 <i>Arzama densa</i>. R. l. Au.
 <i>Nonagria</i> sp. One l. Au.
 <i>Doryodes acutaria</i>. R. l. Au.
 <i>Platysenta atriciliata</i>. R. l. Jn. Jl.
 <i>Parastichtis minuscula</i>. R. l. s. Au.
 Sp.</p> | <p><i>Glaea viatica</i>. S. Oct. 19.
 <i>inulta</i>. Several s. Sp.
 <i>deleta</i>. (See below.)
 <i>Scopelosoma Graefiana</i>. Several s. Sp. Oct. Nov.
 <i>Lithophane semiusta</i>. R. s. Sp. 28, Oct. 18.
 <i>Georgii</i>. One s. Sp.
 <i>laticinerea</i>. C. s. Sp. Oct.
 <i>tepida</i>. R. s. Sp.
 <i>querquera</i>. R. s. Sp.
 <i>Baileyi</i>. (Grote, Can. Entom., ix, 86). Several s. Sp. Oct. Nov.
 <i>viridipallens</i> (Grote, Can. Entom., ix, 215). R. s. Sp. Oct.
 <i>Plusia formosa</i>. One l. Jl.
 <i>verruca</i>. One s. Au.
 <i>Galgula hepara</i>. One s. Nov.
 <i>Catocala neogama</i>. R. s. q. Au.
 <i>piatrix</i>. R. q. s. Au. Sp.
 <i>Homoptera Woodii</i> (Grote, Can. Entom., ix, 88-89). R. l. Jl.
 <i>Spargoloma 6-punctata</i>. R. l. Jl.</p> |
|---|--|

The new species mentioned in the above list is described as follows, by Mr. Grote: —

Glaea deleta n. sp. ♂. Belongs, with *G. tremula*, to the section which I have called *Epiglaea*, in which the thorax has a sharp scale-ridge. Resembles *G. viatica* at first sight, but belongs with *G. pastillicans* to a different section of the genus. Expanse, 46 mm. Of the same color as *G. viatica*, but smaller, and with the transverse posterior line punctate, and the subterminal also broken into spots. Stigmata pale ringed, well sized, concolorous; reniform stained at base. The markings are faint. Hind wings fuscous above; with discal spot and line beneath. Antennæ white at base. Abdomen with ochery anal hairs. (*A. R. Grote.*)

Rare, at sugar, in September. Several (poor) specimens were taken at sugar in October.

Noctuae have been abundant here this fall. Besides the additional species of *Lithophane* mentioned above, I have taken all those given in my former list. The larvæ of *Lithophane disposita* and *L. Bethunei* were found last June at the roots of apple and willow trees. *R. Thaxter.*



Thaxter, Roland. 1877. "Additions to the List of Newton Noctuidae." *Psyche* 2, 80–80. <https://doi.org/10.1155/1877/90712>.

View This Item Online: <https://www.biodiversitylibrary.org/item/44519>

DOI: <https://doi.org/10.1155/1877/90712>

Permalink: <https://www.biodiversitylibrary.org/partpdf/181228>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.