

**Drosos Koutsoubas (\*) Eleni Voultsiadou-Koukoura (\*)**

**THE OCCURRENCE OF *RAPANA VENOSA* (VALENCIENNES, 1846)  
(GASTROPODA, THAIDIDAE) IN THE AEGEAN SEA\*\***

KEY WORDS: Thaididae, *Rapana*, distribution, Aegean Sea, Mediterranean Sea

**Riassunto**

*Rapana venosa* (VALENCIENNES, 1846) è segnalata per la prima volta nel mare Egeo. Si precisa la distribuzione nota di questa specie nel Mediterraneo.

**Summary**

*Rapana venosa* (VALENCIENNES, 1846) is reported for the first time from the Aegean Sea. The hitherto distribution of this species in the Mediterranean Sea is given.

**Introduction**

*Rapana venosa* (VALENCIENNES, 1846) is a species of Pacific origin which entered the Black Sea, through human intervention (GHISOTTI, 1971), in the early fifties of this century and since then it has extended its distribution along the greatest part of its coasts (BILECIK, 1975; CIUHCIN, 1984; GROSSU, 1986; etc.). The colonization of the Black Sea coasts by this species resulted in a decrease of the natural populations of *Ostrea edulis* LINNAEUS in this area (POUTIERS, 1987) because, as it is well known, *R. venosa* is an active predator of bivalves and especially oysters and mussels (GHISOTTI, 1971).

About twenty years later, this species appeared in the Adriatic Sea (GHISOTTI, 1974) and since then, its presence has been recorded from various sites of its northern part (MEL, 1976; CUCAZ, 1983; RINALDI, 1985) where it seems to have been acclimatized (CESARI & PELLIZZATO, 1985). Excluding one more record from Elba Island in the Tyrrhenian Sea (TERRENI, 1980), this species has not been found in any other part of the Mediterranean.

(\*) Department of Zoology, University of Thessaloniki, 54 006 Thessaloniki, Greece

(\*\*) Lavoro accettato il 24 novembre 1990



## Material

In April 1986, one live specimen of *R. venosa* (Fig. 1 a, b) was found in the south-eastern coast of the Bay of Thessaloniki (Micra), in Thermaikos Gulf (North Aegean Sea). It was collected at a depth of 14 m, from a substrate of very coarse sand (median diameter = 1064  $\mu\text{m}$ ); organic carbon of the sediment was 2.45 o/o and water salinity near the bottom 33.2 o/oo. In January 1988, in the locality Chalastra, in the north-western coast of the same Bay, close to the natural oyster banks existing in the area, another live specimen was collected at a depth of 12 m, from a substrate of fine sand (median diameter = 235  $\mu\text{m}$ ); organic carbon of the sediment in this area was 2.33 o/o and water salinity near the bottom 33.5 o/oo.

## Results and Discussion

Both specimens of *R. venosa* found in Thermaikos Gulf had a light grey-brown shell with brown spiral lines which were darker in the body whorl. The dimensions of the two specimens were  $h=110$  mm,  $D=92$  mm and  $h=90$  mm,  $D=58$  mm. To the specimen coming from Micra, numerous individuals of the serpulid polychaete *Protula* sp., two individuals of the bivalve *Anomia ephippium* LINNAEUS and one individual of the barnacle *Balanus amphitrite* DARWIN were attached. A few individuals of *Protula* sp. were also found on the specimen from Chalastra.

The finding of *R. venosa* in Thermaikos Gulf represents its first record in the Aegean Sea and extends its distribution, as suggested by GHISOTTI (1971): «...fra breve tempo apparirà anche in Egeo, nel Mar Ionio...». The present distribution of this species in the Mediterranean Sea is given in the map of Fig. 2. *R. venosa* should be considered as the only species of the genus existing in the Mediterranean Sea, although the species *R. rapiformis* (BORN, 1778) has been reported from the Mediterranean coast of Israel by BARASH & DANIN (1977); the presence of this species in the Mediterranean is doubtful (BRUSCHI et al., 1985) and should be confirmed as it is based on the finding of a juvenile shell and no further reports exist.

The hitherto known distribution of *R. venosa* in the Mediterranean and the Black Sea suggests that this species is carried by boats (GHISOTTI, 1971) in different areas but survives and gets acclimatized only where natural populations or cultures of bivalves (oysters and mussels) exist, in order to be able to satisfy its feeding needs. Such areas are usually the estuaries, where the rich organic matter carried by the rivers allows the development of oyster and mussel populations. Thermaikos gulf seems to be a suitable environment for this species, due to the natural banks of the above bivalves existing there and also to the culture installations situated mainly in the western part of the Gulf.







Fig. 1. *Rapana venosa*: a, ventral view of the specimen collected in the area of Micra (h=110 mm, D=92 mm); b, dorsal view of the same specimen (photo by A. Koukouras).






Fig. 2. Records of *Rapana venosa* in the Mediterranean Sea. Derived from: Drapchin (1953) (according to GHISOTTI 1971), GHISOTTI (1971, 1974), BILECIK (1975), MEL (1976), Goncharov (1977) (according to CESARI & PELLIZZATO 1985), TERRENI (1980), CUCAZ (1983), CESARI & PELLIZZATO (1985), and GROSSU (1986).

## REFERENCES

- BARASH A. & DANIN Z., 1977 - Additions to the knowledge of Indo-Pacific Mollusca in the Mediterranean. *Conchiglie*, Milano, **13** (5-6): 85-116.
- BILECIK N., 1975 - La répartition de *Rapana thomasiana thomasiana* Grosse sur le littoral turc de la Mer Noire s'étendant d'Igneada jusqu'à Calti Burnu. *Rapp. Comm. int. Mer Médit.*, **23** (2): 169-171.
- BRUSCHI A., CEPPODOMO I., GALLI C. & PIANI P., 1985 - Catalogo dei Molluschi conchiferi viventi nel Mediterraneo. ENEA - *Collana Studi Ambientali*, Roma, 111 pp.
- CESARI P. & PELLIZZATO M., 1985 - Insediamento nella Laguna di Venezia e distribuzione Adriatica di *Rapana venosa* (VALENCIENNES) (Gastropoda, Thaididae). *Lavori - Soc. Ven. Sc. Nat.*, **10**: 3-16.
- CIUHCIN V.D., 1984 - Ecology of the gastropod molluscs of the Black Sea. *Acad. Sc. USSR, Kiev Naukova Dumka*, 175 pp. (in Russian).
- CUCAZ M., 1983 - *Rapana venosa* (VALENCIENNES, 1846) vivente nel Golfo di Trieste. *Boll. Malacologico*, Milano, **19**(9-12): 261-262.
- GHISOTTI F., 1971 - *Rapana thomasiana* GROSSE, 1861 (Gastropoda, Muricidae) nel Mar Nero. *Conchiglie*, Milano, **7** (3-4): 55-58.
- GHISOTTI F., 1974 - *Rapana venosa* (VALENCIENNES), nuova ospite Adriatica?. *Conchiglie*, Milano, **10** (5-6): 125-126.
- GROSSU A., 1986 - Gastropoda Romaniae. Caractere generale, istoricul si biologia gastropodelor. Subclasa PROSOBRANCHIA SI OPISTHOBRANCHIA. Editura Litera, Bucuresti, 524 pp.
- MEL P., 1976 - Sulla presenza di *Rapana venosa* (VALENCIENNES) e di *Charonia variegata sequenzae* (AR & BEN.) nell'Alto Adriatico. *Conchiglie*, Milano, **12** (5-6): 129-132.
- POUTIERS J.M., 1987 - Bivalves (Acephales, Lamellibranches, Pelécypodes). In: Fishes FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et Mer Noire. I. Végétaux et invertébrés. Ed. Fischer W., M. Scheider & M.-L. Bauchot. FAO, Rome, 760 pp.
- RINALDI E., 1985 - *Rapana venosa* (VALENCIENNES) spiaggiata in notevole quantità sulla spiaggia di Rimini (Fo). *Boll. Malacologico*, Milano, **21** (10-12): 318.
- TERRENI G., 1980 - Molluschi poco conosciuti dell'Arcipelago Toscano: 1° - Gasteropodi. *Boll. Malacologico*, Milano, **16** (1-2): 9-17.



Koutsoubas, Drosos and Voultsiadou-Koukoura, Eleni. 1991. "The occurrence of *Rapana venosa* (Valenciennes, 1846) (Gastropoda, Thaididae) in the Aegean Sea." *Bollettino malacologico* 26, 201–204.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/199621>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/184269>

#### **Holding Institution**

Smithsonian Libraries and Archives

#### **Sponsored by**

Biodiversity Heritage Library

#### **Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: Società Italiana di Malacologia

License: <http://creativecommons.org/licenses/by-nc/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.