

***Agrias rodriguezi*, new species.**

Body fuscous. Fore wings: apex broadly, outer and inner margins narrowly black; a large crimson space at base, reaching nearly the middle of wing on costa, its outer edge oblique to vein 2 postmedially, then angled and inbent, followed by a broad blue shade from subcostal vein to submedian. Hind wings black with a large deep blue space, not extending above vein 6, leaving costal, inner margin and base broadly black, the outer margin narrowly so. Underneath the coloration is similar to the female of *Agrias aedon* given in the *Biologia* on Pl. 31, fig. 6. Female: expanse 83 mm.

Hab.: Vera Paz, Guatemala.

Type: A female in the Schaus Collection in the United States National Museum.

Vaporous discharge by *Anisomorpha buprestoides*. (Orthoptera: Phasmidae).

A review of information on the secretions of this phasmid and their ejection was published by Samuel H. Scudder in 1876 (*Psyche*, Vol. 1, pp. 137-9). C. J. Maynard gives (*Contr. to Science*, Vol. I, 1889, pp. 31-35) a full account of the thoracic glands and their discharge. An interesting note on the subject will be found also under the name of H. M. Russell (*Proc. Ent. Soc. Wash.* Vol. XIV, No. 3, July-Sept., 1912, p. 117). Observations on the nature and source of the discharge by the insects vary; it is a milky fluid or a very fine mist; it is discharged from definite pores on the sides of the thorax or from some part of the abdomen. An observation differing somewhat from the previous accounts should be put on record, and the diversity of opinions should induce someone to make a thorough study of the matter and publish a full account. The observation to which I refer was made by Mr. C. H. M. Barrett, taxidermist of the United States Biological Survey, while on a collecting trip in Florida. At Miakka Lake, Manatee County, during a thunderstorm, June 19, 1918, Mr. Barrett found a number of pairs of *Anisomorpha buprestoides*, in copula, in the interstices of a pile of boards in a deserted cabin. When suddenly uncovered or otherwise disturbed, the insects discharged from the end of the abdomen vapor in the form of small puffs appearing two inches from the abdominal apex. The discharge in each case was immediately preceded by a crackling sound similar to that made by a small electric spark. Mr. Barrett's observation that *Anisomorpha buprestoides* discharges a vapor from end of abdomen and that the discharge is preceded by a peculiar crepitation, differs, so far as I am aware, from any previously recorded.—W. L. McATEE, Biological Survey, U. S. Dept. Agriculture, Washington, D. C.



McAtee, W. L. 1918. "Vaporous discharge by *Anisomorpha buprestoides* (Orthoptera: Phasmidae)." *Entomological news, and proceedings of the Entomological Section of the Academy of Natural Sciences of Philadelphia* 29, 388.

View This Item Online: <https://www.biodiversitylibrary.org/item/20206>

Permalink: <https://www.biodiversitylibrary.org/partpdf/18716>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.