

less sharply toothed leaves on broadly winged petioles and the achenes, although smaller than in var. *petiolata*, are not so small as in var. *gracilipes*, the outer being 4–5.3 mm. long, the inner 5–6.5 mm. long and with marginal awns 2.2–3.6 mm. long.

Very young specimens from peaty shores of the Little Ossipee River, Limington, Maine (*Fernald & Long*, no. 14,843) may belong here.—M. L. FERNALD, Gray Herbarium.

AN OMISSION IN THE PRELIMINARY LIST OF NEW ENGLAND RANUNCULACEAE.—By a regrettable oversight, chiefly my own, the one New England record for *Cimicifuga racemosa* (L.) Nutt., var. *dissecta* Gray was omitted from the list of New England *Ranunculaceae* published in RHODORA, xx. 182. The plant in question was collected by Dr. E. H. Eames at Stratford, Conn., in 1893 and was duly included in the Connecticut Flora. There is also a specimen in the Gray Herbarium. *C. racemosa*, var. *dissecta* should have been entered in the list and marked with a cross.—C. A. WEATHERBY, East Hartford, Connecticut.

Vol. 21, no. 244, including pages 73 to 88, was issued 9 May, 1919.



Weatherby, Charles Alfred. 1919. "An Omission in the preliminary List of New England Ranunculaceae." *Rhodora* 21, 104–104.

View This Item Online: <https://www.biodiversitylibrary.org/item/14492>

Permalink: <https://www.biodiversitylibrary.org/partpdf/188365>

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.