Centaurea Maculosa in Illinois.—Within the past year this species has been reported from Michigan by E. E. Sherff and from Indiana by E. D. Hull², hence its recent detection in two of the northern counties of Illinois is not at all surprising. The discovery of the spotted knapweed in 1945, by Dr. E. W. Fell of Rockford, brings the total number of centaureas now known in this state to seven, the others being C. calcitrapa L., C. cyanus L., C. jacea L., C. nigra L., C. solstitialis L. and C. vochinensis Bernh. The following specimens of C. maculosa L. are in the herbarium of the Illinois State Museum at Springfield, and through the courtesy of the curator, Professor Geo. D. Fuller, duplicates have been deposited in the herbarium of the University of Illinois. The collection-data are as follows:

Illinois: roadside, generally scattered over the county, three miles south of Rockford, Winnebago County, July 30, 1945, E. W. Fell No. 45485; roadside, three miles southeast of Cherry Valley, September 27, 1945, E. W. Fell, No. 45743.—George Neville Jones, University of Illinois.

¹ Rhodora 48: 98 (1946).

² Op. cit. 391.

Volume 49, no. 578, consisting of pages 25-60 and plates 1053 and 1054, was issued 14 February, 1947.



Jones, George Neville. 1947. "Centaurea maculosa in Illinois." *Rhodora* 49, 84–84.

View This Item Online: https://www.biodiversitylibrary.org/item/14519

Permalink: https://www.biodiversitylibrary.org/partpdf/189063

Holding Institution

Missouri Botanical Garden, Peter H. Raven Library

Sponsored by

Missouri Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: http://creativecommons.org/licenses/by-nc-sa/3.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.