

Banding pattern comparisons between *Akodon iniscatus*, and *Akodon puer* (Rodentia, Cricetidae)

By MARIA A. BARROS, ROSA C. LIASCOVICH, L. GONZALEZ, MARTHA S. LIZARRALDE,
and O. A. REIG

GIBE, Dpto. de Cs. Biológicas, FCEyN, Universidad de Buenos Aires, Argentina, Instituto de Evolución y Ecología, Universidad Austral de Chile, Valdivia, Chile, and Centro Austral de Investigaciones Científicas, CONICET, Ushuaia, Argentina

Receipt of Ms. 18. 08. 1988

Abstract

Described the chromosomes of *Akodon iniscatus* ($2n = 33-34$, $FN = 42$) and compared with the karyotypes of *A. puer* ($2n = 34$, $FN = 40$). 37 specimens of *A. iniscatus* and 11 of *A. puer* were studied on their C, G and fluorescent (Q and CMA) banding patterns. Whole chromosomes and chromosome arms were shared between both species, 38 arms showed the same G-banding pattern in both species. *A. iniscatus* contains mainly wide-spread centromeric heterochromatin that is absent in *A. puer* and rare in other akodonts. Chromomycin banding revealed that centromeric heterochromatin was homogeneous and GC poor in all chromosomes, whereas the telomeric heterochromatin was GC rich. The present results support the hypothesis of close relationship between *A. iniscatus* and *A. puer* and that the former probably derived from an *Akodon* stock originally differentiated in the low-lands of northeastern Argentina. Therefore, *A. iniscatus* is related with the varying $2n = 40$ akodont chromosome group.

Introduction

Akodon is a highly speciose genus of South American cricetids that is the most widely and better known of the genera of sigmodontine mice in its cytogenetics. Of the 38 recognized species of *Akodon* (REIG 1987), 29 are known in their karyotype and 17 have also been analysed in their banding patterns (BIANCHI and MERANI 1980; BIANCHI and MERANI 1984; BIANCHI et al. 1976; GALLARDO 1982; LOBATO et al. 1982; RODRIGUEZ et al. 1983; VITULLO et al. 1986; YONENAGA et al. 1975, 1976; YONENAGA-YASSUDA 1979).

Akodon iniscatus Thomas is one of the few species of the genus for which the chromosomal complement has not been described. Several specimens from Collón Curá, Neuquén Province, Argentina were karyotyped some time ago by two of the present authors (O. A. R. and L. G.), but these determinations remained unpublished. More recently, *A. iniscatus* was found to be the most common cricetid in the coastal region of Chubut Province, Argentina where its presence had not been reported previously. *A. iniscatus* karyotype was very similar to that of *Akodon puer* Thomas, (BARQUEZ et al. 1980; KAJON et al. 1984; VITULLO et al. 1986). Cytogenetic results on *A. iniscatus* and their comparison with new results from specimens of *A. puer* are presented here.

Material and methods

Cytogenetic analysis was performed on 37 specimens of *A. iniscatus* from Chubut and Neuquén Provinces and on 11 specimens of *A. puer* from Jujuy Province, Argentina. Localities for *A. iniscatus* were: 5 km S Pto. Madryn, Chubut (13 females and 18 males); Punta Lomas, 20 km S Pto. Madryn, Chubut (2 males), and Collón Curá, Neuquén (2 females and 2 males). *A. puer* was collected from León, Jujuy (7 females and 4 males). These specimens are deposited as skin and skull museum specimens in the Museo Municipal de Ciencias Naturales "Lorenzo Scaglia" of Mar del Plata,

Argentina (MMP), and in the Collection of Mammals of the Centro Nacional Patagónico (CNP), in Puerto Madryn, Argentina.

Additional specimens have been studied for taxonomic comparisons and identification. These include the type specimen of *A. nucus* at the British Museum of Natural History (BMNH 26.10.11-42, male from Chos Malal, Neuquén), the type of *A. iniscatus* (BMNH 3.7.9.64, female from Valle del Lago Blanco, Chubut), a topotypical series of *A. iniscatus*, and 4 specimens from 20 km S of Leleque, Chushamen, Chubut.

For taxonomic studies, cranial and molar tooth morphology were examined with a Wild M-5 stereomicroscope. Skull and teeth were measured with a digital caliper and/or with an eye-piece reticule of the stereomicroscope to the nearest 0.1 mm following standard conventions.

Metaphase chromosome preparations were obtained from bone marrow (BARROS and PATTON 1985) of animals injected with yeast 24 hs prior to sacrifice (LEE and ELDER 1980). C-bands were obtained following the procedure described in BARROS and PATTON (1985). SEABRIGHT's (1971) procedure was followed for G-bands. Chromomycin A₃/methyl green (CMA) bands were done according to HACK and LAWCE (1980). Quinacrine/actinomycin D (Q) banding was done according to HACK and LAWCE (1980). All fluorescent bands were photographed with Kodak Tri-X film exposed at ASA 800.

Diploid number and chromosome morphology were determined for all specimens, chromosomes were classified according to LEVAN et al. (1964). REIG and KIBLISKY's (1969) size proposal convention was followed. Fundamental number is defined as the number of autosomal arms.

Results

Taxonomy

The type specimens of *A. iniscatus* Thomas, 1919 and of *A. nucus* Thomas, 1926 were found in Valle del Lago Blanco, SW of Chubut and Chos Malal, N of Neuquén, respectively. The two localities are separated by nearly 935 km in the mountainous region of the southern Andean slopes of Argentina (Fig. 1). The type specimens show that the two taxa are very similar in external, skull and tooth features, though they differ markedly in size, *A. nucus* being noticeably larger than *A. iniscatus*. *A. nucus* also differs from *A. iniscatus* in having nasals longer than frontals, parietals comparatively longer, the anterior border of the zygomatic plate in an upward position, and the M3 relatively smaller (Fig. 2).

Judging from the holotypes and additional topotypical or referred specimens in the British Museum of Natural History, the differences found seem to justify full species




Fig. 1. Map showing localities of *A. iniscatus* and *A. nucus*. 1. Chos Malal (type locality for *A. iniscatus*); 2. Collón Curá; 3. El Maitén; 4. Leleque; 5. Lago Blanco (Type locality for *A. nucus*); 6. Puerto Madryn

distinction, as proposed by one of us (REIG 1987). However, CABRERA (1961) considered that *A. nucus* is merely a subspecies of *A. iniscatus*, a course implicitly followed by HONACKI et al. (1982), who did not list *A. nucus* as a separate species.

Our specimens from Collón Curá, a locality for which THOMAS (1927) cited *A. nucus*, are fairly similar to the holotype of this species, and share all its distinctive characters. Our small available series from Leleque, which comes from 42 km S of Maitén, the type locality of the local form *A. iniscatus collinus* Thomas, 1919, is almost undistinguishable from *A. iniscatus*, thus confirming the distinction of the two (Table 1).

However, our sample from localities near Puerto Madryn, a region where no *Akodon* of this group has been found previously, is quite intermediate between *A. nucus* and *A. iniscatus* in size and morphology (Table 1). Additionally, the karyotype of specimens from these localities are similar to those from Collón Curá reported in this paper. Thus, the intermediacy of the Puerto Madryn *Akodon* as regards typical *A. iniscatus* and *A. nucus*, sheds doubts on the species status of *A. nucus*. We are probably dealing with a geographically variable group of populations which are better considered, at the present state of knowledge, as a single polytypic species. It would support CABRERA's proposal of treating *A. nucus* as a subspecies of *A. iniscatus*. However, this issue is a tentative one, and must be tested by further morphological and cytogenetic studies. But at present, we prefer to allocate provisionally all our chromosomally studied specimens to *Akodon iniscatus* without further refinement.




Fig. 2. Skull in lateral, dorsal, and ventral views of the holotypes of *A. iniscatus* and *A. nucus*

Table 1. Measurements (mm) of *A. iniscatus* (AI) and *A. nucus* (AN) holotypes, and samples from Leleque, Puerto Madryn, and Collón Curá. N: sample size; X: mean; SD: standard deviation

Variable	AI	AN	Leleque sample			PTO. Madryn sample			Collón Curá sample		
	Type	Type	N	X	SD	N	X	SD	N	X	SD
Total length	145.0	181.0	4	140.0	6.4	11	153.0	8.3	4	173.0	8.6
Tail length	53.0	77.0	4	55.8	2.2	11	60.2	6.0	4	65.9	3.9
Hind foot	19.0	23.0	4	18.9	0.6	12	20.6	0.9	4	22.6	1.0
Ear	11.0	14.0	4	11.8	0.2	12	12.5	0.7	4	12.8	1.5
Skull total length	23.7	26.8	4	23.0	0.6	13	24.5	0.9	4	25.8	0.2
Condylar basal	21.7	25.2	4	20.8	0.5	13	22.6	0.9	4	24.0	0.1
I-M3	11.1	12.4	4	10.5	0.3	13	11.1	0.4	4	12.0	0.3
Diastema	5.6	6.7	4	5.5	0.3	13	6.0	0.3	4	6.4	0.2
Zygomatic breadth	12.4	14.7	4	11.7	0.3	13	12.8	0.4	4	13.3	0.1
Mastoid breadth	11.3	12.1	4	11.1	0.1	13	11.5	0.1	4	11.3	0.2
Inteorb. constrict.	4.0	4.4	4	4.4	0.2	13	4.3	0.1	4	4.2	0.1
Nasal length	8.4	10.3	4	7.8	0.7	13	8.7	0.4	4	9.6	0.2
Frontal length	9.1	9.7	4	9.1	0.2	13	9.0	0.3	4	9.4	0.5
Parietal length	5.1	6.4	4	4.4	0.2	13	5.0	0.4	4	5.2	0.2
Incisor foramina	5.5	6.4	4	5.3	0.3	13	5.6	0.3	4	6.1	0.3
M1M3 alveolar	3.5	4.3	4	4.0	0.1	12	4.1	0.2	4	4.4	0.1
m1m3 alveolar	3.8	4.5	4	4.1	0.1	13	4.2	0.2	4	4.5	0.3

Chromosomal analysis

35 specimens of *A. iniscatus* have a chromosome complement of $2n = 34$, $FN = 42$ (Fig. 3). These include 29 specimens from the Puerto Madryn area, 2 from Punta Lomas, and 4 from Collón Curá. Specimens from all three localities show the same karyotype. The first four pairs are large biarmed chromosomes distinguished from the remaining autosomes by a sharp size gap. Pair 1 is submetacentric, whereas pairs 2 to 4 are metacentric. Pairs 5 to 15 are telocentric. As in most akodontines, the last autosomal pair, (16), is a small metacentric. Pair 5 is medium-sized and pairs 6 to 16 are small autosomes gradually decreasing in size. The X and Y are small telocentrics, representing 5.12 % and 2.46 % of the complement respectively.

A male and a female from Puerto Madryn, which presented a heteromorphic Robertsonian variant of $2n = 33$, $FN = 42$ were found. The decrease of $2n = 34$ to 33 results from the presence of a metacentric chromosome whose G-band pattern matches the combined banding patterns of telocentric autosomes 9 and 11 of the $2n = 34$ karyotype (Fig. 3D).

C and G banding results refer only to the Puerto Madryn sample. Well-developed pericentromeric C-bands (Fig. 3B) are present on nearly all autosomes of *A. iniscatus*, in striking contrast to other species of the genus, which are usually C poor. Additionally pair 16 shows a band in the telomeric region of both arms, and pair 5 shows a small interstitial band. G-bands are illustrated in Fig. 3C.

A. puer has a karyotype of $2n = 34$, $FN = 40$ (Fig. 3). The first three pairs are large biarmed chromosomes, differing sharply from the remaining autosomes in size. Pair 1 is submetacentric and pairs 2 and 3 are metacentric. Pair 4 is medium-sized telocentric, and pairs 5 to 15 are small telocentrics. Pair 16 is a small metacentric one. The X chromosome is a medium-sized (6.87 %) subtelocentric and the Y is a small (2.20 %) telocentric chromosome.

C-bands (Fig. 3B) show that heterochromatin is found only in the short arm of the X and in the centromeric region of the Y. G-bands of *A. puer* are presented in Fig. 3C.

Fig. 4 shows the G-band correspondence between *A. puer* and *A. iniscatus*. This comparison reveals that 38 arms are shared by both species. Of the 38 shared arms, 28 belong to 24 whole chromosomes, two of which are biarmed. The remaining 10 shared




Fig. 3. *Akodon puer* and *A. iniscatus* giemsa stained karyotypes (A), C-bands (B), and G-bands (C). *A. iniscatus* heteromorphic pair of $2n = 33$ morph (D)




Fig. 4. G-banding pattern comparison between *A. puer* (left) and *A. iniscatus* (right). Pair numbers correspond to their position in each species' karyotype (A). G-banded ideogram comparison between both species (B). Diagram showing arms involved in monobrachial homology (AI = *A. iniscatus* and AP = *A. puer*) (C).

arms do not correspond to whole chromosomes in both species. Pair 9 of *A. iniscatus* and the short arm of the X of *A. puer* are unique elements to these species. Low G-band resolution of pair 16 did not clarify if this pair is shared between both species. Since the microchromosomal pair is present in most *Akodon* species, we conventionally take it as a shared pair. Although, the size difference between pair 16 of *A. iniscatus* and *A. puer* suggests an addition of heterochromatin in the former.

In order to analyze the shared chromosomes between both species we nominate *A. puer* as "AI" and *A. iniscatus* as "AI"; short arm as "p" and long arm as "q". Arm 1q and pairs 2 and 3 are shared by both species. The 1p of AP is shared with the 4p of AI. Chromosome 4 of AP is shared with 4q of AI. Chromosome 9 of AP is shared with 1p of AI. Pairs 5, 6, 7, 8, 10, 11, 12, 13, 14, and 15 are shared by both species. Thus, two biarmed autosomes of both species can be postulated as shared chromosomes. The remaining biarmed chromosomes (pairs 1 of both species and pair 4 of AI) result from Robertsonian rearrangements of different arms showing monobrachial homology (Fig. 4C) (BAKER and BICKHAM 1986; CAPANNA et al. 1985; CORTI et al. 1986).

Fluorescent bands are illustrated in Fig. 5. In the *A. iniscatus* karyotype (Fig. 5C), most CMA-positive bands occur within the euchromatin and in the telomeric heterochromatic region of pair 16. The remaining heterochromatic regions are CMA-negative. All CMA-positive bands seem to correspond with G-negative bands, with the exception of pair 16. Fluorescent CMA-bands suggest heterochromatin similarity in base composition among centromeric regions, as well as heterochromatin heterogeneity between these regions and the telomeric region of pair 16. *A. iniscatus* Q-bands (Fig. 5D) show that fluorescent and quenched Q-bands seem to correspond with G-positive and G-negative bands respectively, although better banding resolution is found in G-bands. The Q banding pattern is opposite to that obtained with Chromomycin A₃, except in the centromeric regions that are CMA-negative and Q-negative. *A. puer* CMA-bands (Fig. 5A) reveal that CMA-positive bands are located within the euchromatin, in regions which were identified as G-negative.

Heterochromatin observed in the sex chromosomes is CMA-negative. *A. puer* Q-bands (Fig. 5B) show that the Q-band pattern, as in *A. iniscatus*, is opposite to that with Chromomycin A₃. The same correspondence between Q and G bands as in *A. iniscatus* was found.

Discussion

Several results indicate that akodontine rodents underwent an intensive chromosomal evolution. The high degree of cladogenesis exhibited by 11 genera and 66 species belonging to the tribe (REIG 1987), is correlated with a high number of chromosomal rearrangements. However, this correlation does not imply an overall causal link between evolutionary divergence and chromosomal changes in the Akodontini. Within *Akodon*, a high range of chromosomal numbers is found, from $2n = 14$ in *A. arviculoides* (YONENAGA 1972) to $2n = 52$ in *A. illuteus* (LIASCOVICH et al., 1989), *A. longipilis* (SPOTORNO and FERNANDEZ 1976), *A. olivaceus* (GALLARDO 1982), etc. Species of the genus *Akodon* can be divided into two assemblages according to their overall karyotype characteristics. Species of the first group have a varying karyotype of around $2n = 40$ chromosomes which may be reduced by fusions, the second group comprises species with a uniform $2n = 52$, FN = 56 karyotype. A good knowledge of the degree of whole chromosome and arm sharing among the species is required for this study.

The term "shared chromosomes" refers to whole chromosomes or chromosome arms which share banding patterns. G-banding was first employed to analyze chromosomal homology of akodontines in a study of *A. azarae*, *A. molinae*, and "*A.*" *obscurus* (BIANCHI et al. 1976). Full chromosome arm homology of species sharing the $2n = 52$ karyotype was

demonstrated in *A. longipilis*, *A. olivaceus* and *A. sanborni* (GALLARDO 1982), *A. xanthorhinus* (RODRIGUEZ et al. 1983), and *A. illuteus* (LIASCOVICH et al. 1989). BIANCHI and MERANI (1984) studied the karyological relationships among several akodontine species of the varying karyotype group: *A. azarae*, *A. dolores*, *A. molinae*, *A. mollis*, "*A.*" *obscurus*,




Fig. 5. Fluorescent banded karyotypes. *A. puer* CMA-bands (A), and Q-bands (B); *A. iniscatus* CMA-bands (C), and Q-bands (D)

A. urichi, and *A. varius*. They demonstrated a high degree of arm sharing among species, which had mostly diverged by presumed Robertsonian processes. These studies are germane to the hypothesis suggested by BIANCHI et al. (1971) that chromosome evolution within the akodontine rodents proceeds from low to high numbers by centromeric dissociations, or conversely from high to low numbers by centromeric fusions. One of the present authors (REIG 1987; VITULLO et al. 1986) claimed that the second alternative is the most plausible. Our present results show a high degree of chromosomal homology between *A. iniscatus* and *A. puer*, which suggests close phylogenetic relations between these species. The reductional process of chromosomal number in these two species involved metacentrics, resulting from the fusion of the same autosomes, and then give fully shared chromosomes, and of metacentrics derived from the fusion of different telocentrics which show merely monobrachial homology. BAKER and BICKMAN (1986), CAPANNA et al. (1985), CORTI et al. (1986), and MORITZ (1986), have recently pointed out that monobrachial homology probably results in full hybrid sterility. The degree of meiotic breakdown depends on the number of fusions that are monobrachially homologous, and on the specific chromosomes involved. Generally, those studies are based on hybrids with monobrachial homology. The allopatric distribution presented by the species under study led us unable to find hybrids in nature. In order to evaluate the extent of monobrachial homology as a cause of hybrid sterility, studies on laboratory hybrids should be done.

The finding of a polymorphism for one Robertsonian rearrangement in two of the thirty seven studied specimens of *A. iniscatus* is in good keeping with the general occurrence of Robertsonian translocations in the group. In the Puerto Madryn population, this fusion polymorphism is not as extensive as it is in *A. dolores*, *A. molinae* (BIANCHI and MERANI 1980), and *A. simulator* (LIASCOVICH et al. 1989). In view of the extensive arm homology found between *A. iniscatus* and *A. puer*, it was of interest to investigate if other chromosomal changes revealed by more refined techniques were involved in the differentiation of these related species.

As species of *Akodon* usually show very little heterochromatin, it was unusual to find a wide-spread occurrence of centromeric heterochromatin in the *A. iniscatus* complement. The few previously known cases of centromeric heterochromatin in *Akodon* were in *A. mollis* (LOBATO et al. 1982), *A. arviculoides*, *A. aff. arviculoides* (YONENAGA et al. 1975; YONENAGA-YASSUDA 1979; but see HONACKI et al. 1982, and LIASCOVICH and REIG 1989, for correct naming of these forms), *A. (Deltamys) kempi* (SBALQUEIRO et al. 1984), and *A. (Hypsims) budini* (VITULLO et al. 1986), in which modest amounts of heterochromatin have been described. Relevant to our study is the fact that C-positive bands are almost nonexistent in *A. puer* and *A. azarae* (VITULLO et al. 1986), both of which are considered to be closely related to *A. iniscatus* (REIG 1987).

In other groups of mammals, G-banding patterns appear to be quite stable, but extensive interspecific variation in constitutive heterochromatin has been reported (ARNASON 1974; ELDER and LEE 1985; PATHAK et al. 1973; STOCK 1981). We have no indication of the processes which were involved in the presence of the centromeric and telomeric heterochromatin, or of its function in *A. iniscatus*, but it is relevant to these problems to have found sequence diversity in the heterochromatin of *A. iniscatus*.

Some fluorescent antibiotic dyes exhibit a nucleotide specificity with helical DNA. A guanine and cytosine (GC) specificity was found for chromomycin A₃ (SAHAR and LATT 1978, 1980), whereas a less clear adenine and thymine (AT) specificity has been proposed for quinacrine (COMINGS 1978). Despite the fact that the molecular mechanisms of interaction between these specific fluorescent dyes and DNA are not well known, they permit a cytological characterization of presumed heterogeneity of DNA sequences within the genome (BARROS and PATTON 1985).

Chromomycin A₃ has revealed heterogeneity between the heterochromatin of all chromosome pairs and the telomeric heterochromatin found in pair 16 of *A. iniscatus*

karyotype. The latter type of heterochromatin appears as predominantly GC rich, whereas the former is not. This telomeric heterochromatic region is not observed in *A. puer*'s pair 16. An increase/decrease of this GC rich region could be responsible for the size difference observed in pair 16 between the species analyzed.

JOHN et al. (1985) present a review of the concept of equilocality, the similar location of C-bands found between non-homologous chromosomes within the complement. By comparative fluorescence studies on heterochromatin of acridid grasshoppers, these authors found that certain groups of chromosomes that presented a similar C-band distribution also showed a similar fluorescence behaviour with specific dyes, although heterochromatin heterogeneity was also found between similar sites which differ in size and structure. A form of equilocality is present in *A. iniscatus* where all chromosomes, except pair 16, carry a C-band at a centric location, which is CMA and Q negative, while a different heterochromatic site (telomeres of pair 16) respond differentially to CMA.

Although centromeric regions in *A. iniscatus* are heterochromatic, while those in *A. puer* are euchromatic, no fluorescent staining difference was found between them. Findings of CMA and Q negative fluorescence in centromeric heterochromatin could reflect an insufficient number of AT or GC clusters of DNA (SCHWEIZER 1981; SAHAR and LATT 1980). While DNA composition is an important determinant in fluorescence staining, the possible role of chromosomal proteins remains unknown (COMINGS 1978). Comparison of the centromeric regions of *A. iniscatus* and *A. puer* therefore needs to be defined by further sequence analysis. Coincidentally, no composition differences were found between the remaining complement of both species, except for pair 16, which also confirms the chromosome correspondence analysis based on G-bands.

In biarmed complements, equilocality has been explained in terms of a specific ordering of the chromosomes during metaphase (SCHWEIZER and EHRENDORFER 1983), where a sequence transposition or convergence is assumed to occur between repeated sequences of chromosomes in proximity, but according to JOHN et al. (1985) the initial production of C-bands could occur by sequence amplification. Although we don't know if a non-random chromosome distribution is present in *A. iniscatus*, chromosome interactions between centromeric heterochromatic areas have been found to occur between several chromosomes at metaphase (Fig. 6). Following JOHN et al. (1985), an alternative basis for equilocal distribution of centric heterochromatin in *A. iniscatus* could be explained by concerted amplification of the centromeric C-bands between non-homologous chromosomes.

A peculiar result of this study is the finding of a reduced $2n = 34$ karyotype of the varying $2n = 40$ chromosome group in *A. iniscatus*, as this species inhabits an area where all the remaining akodontine mice show the constant $2n = 52$ karyotype. The latter was found in *A. andinus*, *A. brachiotis*, *A. hershkovitzi*, *A. illuteus*, *A. longipilis*, *A. olivaceus*, *A. sanborni*, *A. xanthorhinus*, and in species of *Geoxus* and *Chelemys* (but see REIG 1987, and LIASCOVICH et al. 1989), all of which inhabit the southern Andes and the Patagonian tableland. REIG (1984, 1986, 1987) recently postulated that the akodontine radiation took its origin in the Puna region during Miocene times from a $2n = 52$ *A. andinus*-like *Akodon* ancestor. From this area of original differentiation a chromosomally conservative phylad dispersed to the southern Andes and to Patagonia, maintaining the original $2n = 52$ karyotype unaltered in a cladogenetic process which involved species of the subgenera *Akodon* and *Abrothrix* and genera such as *Chelemys*, *Geoxus*, and *Notiomys*. Another main dispersing branch descended from the Puna to the southeastern lowlands, presumably from a $2n = 40$ *A. boliviensis*-like species, giving rise to the different species in central and northern Argentina, Paraguay, Brazil, and Uruguay. Regarding the origin of *A. iniscatus* and the related (and now considered co-specific) Neuquenian form *A. nucus*. Two alternative hypothesis were postulated by REIG (1987): both forms may have derived from the southern Andean branch or from the southeastern lowland one. REIG inclined himself




Fig. 6. Chromosome interactions between centromeric heterochromatin in *A. iniscatus*

toward the second alternative in view of the greater morphological similarities of these forms with *A. puer* and *A. azarae*. The results presented here indicating a high chromosomal similarity between *A. iniscatus* and *A. puer* are consistent with the latter hypothesis.

Acknowledgements

We thank the CENPAT mammal-team (D. ALOIA, S. DAHINTEN, L. KUBA, S. PEDRAZA, and A. TOYO) for help in the laboratory and field work. M. B. ESPINOSA, A. GOSZTONY, J. KIRSCH, S. MERANI, M. ORTELLS, O. SCAGLIA, and R. TAYLOR are greatly acknowledged for assistance in field work. We thank Dr. J. L. PATTON for valuable comments on the manuscript. Dr. H. CICILEO and Dr. L. ARISTARAIN provided institutional facilities at the CENPAT. C. HERKE has the credit of the drawings. This paper is mainly the result of the CONICET grant PID N° 3085300/85, granted to O. A. REIG.

Resumen

Comparación de patrones de bandeo entre Akodon iniscatus y Akodon puer (Rodentia, Cricetidae)

Se describen los cromosomas de *Akodon iniscatus* ($2n = 33-34$, $NF = 42$) y se comparan con el cariotipo de *A. puer* ($2n = 34$, $NF = 40$). Se estudian los patrones de bandeo C, G y fluorescentes (Q y CMA) de 37 ejemplares de *A. iniscatus* y 11 de *A. puer*. *A. iniscatus* presenta heterocromatina localizada principalmente en la región centromérica, la cual está ausente en *A. puer* y es poco frecuente en otros akodontinos. Las bandas de cromomicina revelan que la heterocromatina de la región centromérica de todos los cromosomas es homogénea y pobre en GC, mientras que la heterocromatina telomérica es rica en GC. Ambas especies comparten brazos cromosómicos y cromosomas completos, siendo 38 los brazos en los que se encuentra el mismo patrón de bandas G. Estos resultados apoyan la hipótesis de una cercana relación entre *A. iniscatus* y *A. puer* y sugieren que la primera probablemente deriva de un stock akodontino que originalmente se diferenció en las tierras bajas del noreste de Argentina. Por lo tanto, *A. iniscatus* estaría relacionada con el grupo de especies de *Akodon* de alrededor de 40 cromosomas.

Zusammenfassung

Vergleiche von Bandenmustern zwischen Akodon iniscatus und Akodon puer (Rodentia, Cricetidae)

Der Karyotyp von *Akodon iniscatus* ($2n = 33-34$, $FN = 42$) wird erstmals beschrieben und mit dem von *A. puer* ($2n = 34$, $FN = 40$) verglichen. Dazu wurden die C-, G- und Fluoreszenzbandenmuster

(Q, CMA) von 37 *A. iniscatus* und 11 *A. puer* untersucht. Eine Reihe ganzer Chromosomen und 38 Chromosomenarme besitzen bei beiden Arten gleiche G-Bandenmuster. Im Gegensatz zu *A. puer* und den meisten *Akodon*-Arten besitzt *A. iniscatus* auffälliges Heterochromatin in der Zentromeren-Region. Die CMA-Färbung zeigt, daß das Heterochromatin der Zentromeren-Region arm an den Basen CG ist, wogegen das telomerische Heterochromatin deren viele enthält. Der Vergleich der Chromosomen spricht insgesamt für eine enge Verwandtschaft von *A. iniscatus* und *A. puer*. *A. iniscatus* hat sich wahrscheinlich in tiefgelegenen Gebieten im Nordwesten Argentiniens differenziert und ist innerhalb von *Akodon* der Artengruppe mit etwa 40 Chromosomen zuzuordnen.

References

- ARNASON, V. (1974): Comparative chromosome studies in Cetacea. *Hereditas* **77**, 1–36.
- BAKER, R. J.; BICKHAM, J. W. (1986): Speciation by monobrachial centric fusions. *Proc. Natl. Acad. Sci.* **83**, 8245–8248.
- BARQUEZ, R. M.; WILLIAMS, D. F.; MARES, M. A.; GENOWAYS, H. H. (1980): Karyology and morphometrics of three species of *Akodon* (Mammalia: Muridae) from northwestern Argentina. *Ann. Carn. Mus. Nat. His.* **49**, 379–403.
- BARROS, M. A.; PATTON, J. L. (1985): Genome evolution in pocket gophers (genus *Thomomys*). III. Fluorochrome-revealed heterochromatin heterogeneity. *Chromosoma (Berl.)* **92**, 337–343.
- BIANCHI, N. O.; MERANI, M. S. (1980): Chromosome 1 polymorphism and viability in *Akodon* (Rodentia, Cricetidae). *Simp. Ecol. Gen. de la Esp. Anim. Univ. S. Bolivar, Caracas, Venezuela* 211–238.
- (1984): Cytogenetics of South American akodont rodents (Cricetidae). X. Karyological distances at generic and intergeneric levels. *J. Mammalogy* **65**, 206–219.
- BIANCHI, N. O.; REIG, O. A.; MOLINA, O. J.; DULOUT, F. N. (1971): Cytogenetics of the South American akodont rodents (Cricetidae). I. A progress report on Argentinian and Venezuelan forms. *Evolution* **21**, 724–736.
- BIANCHI, N. O.; VIDAL-RIOJA, L.; BIANCHI, M. S. (1976): Cytogenetics of South American akodon rodents (Cricetidae). II. Interspecific homology and G-banding patterns. *Cytologia* **41**, 139–144.
- CABRERA, A. (1961): Catalogo de los mamíferos de America del Sur. II (Sirenia, Perissodactyla, Artiodactyla, Lagomorpha, Rodentia). *Rev. del Mus. Arg. de Cs. Nat. "B. Rivadavia"*, (Zoologia) **4**, 309–732.
- CAPANNA, E.; CORTI, M.; NASCETTI, G.; BULLINI, L. (1985): Pre-and post-mating isolating mechanisms in the speciation of the European long-tailed house mouse *Mus musculus domesticus*. *Acta Zool. Fenica* **170**, 115–120.
- COMINGS, D. E. (1978): Mechanisms of chromosome banding and implications for chromosome structure. *Ann. Rev. Genet.* **12**, 25–46.
- CORTI, M.; CAPANNA, E.; ESTABROOK, G. F. (1986): Microevolutionary sequences in house mouse chromosomal speciation. *Syst. Zool.* **35**, 163–175.
- ELDER, F. F. B.; LEE, M. R. (1985): The chromosomes of *Sigmodon ochrognathus* and *S. fulviventer* suggest a realignment of *Sigmodon* species groups. *J. Mammalogy*, **66**, 511–518.
- GALLARDO, M. H. (1982): Chromosomal homology in southern *Akodon*. *Experientia* **38**, 1485–1487.
- HACK, M. S.; LAWCE, H. J. (1980): The association of cytogenetic technologist cytogenetics laboratory manual. Univ. of Calif. USA.
- HONACKI, J. H.; KINMAN, K. E.; KOEPL, J. W. (1982): Mammal species of the World. A taxonomic and geographic reference. Lawrence, Kansas: Allen Press and Association of Systematic Collections.
- JOHN, B.; KING, M.; SCHWEIZER, D.; MENDELAK, M. (1985): Equilocality of heterochromatin distribution and heterochromatin heterogeneity in acridid grasshoppers. *Chromosoma* **91**, 185–200.
- KAJON, A. E.; SCAGLIA, O. A.; HORGAN, C.; VELAZQUEZ, C.; MERANI, M. S.; REIG, O. A. (1984): Tres nuevos cariotipos de la tribu Akodontini (Rodentia, Cricetidae). *Rev. del Mus. Arg. de Cs. Nat. "B. Rivadavia"*, XIII **48**, 461–469.
- LEE, M. R.; ELDER, F. F. B. (1980): Yeast stimulation of bone marrow mitosis for catogenetic in investigations. *Cytogenet. Cell. Genet.* **26**, 36–40.
- LEVAN, A.; FREDGA, K.; SANDBERG, A. (1964): Nomenclature for the centromeric position on chromosomes. *Hereditas* **52**, 201–220.
- LIASCOVICH, R. C.; BARQUEZ, R. M.; REIG, O. A. (1989): A karyological and morphological reassessment of *Akodon (Abrothrix) illuteus*. *Thomas, J. Mammalogy*, **70** (2), 386–391.
- LIASCOVICH, R. C.; REIG, O. A. (1989): Low chromosome number in *Akodon cursor montensis* Thomas, and karyologic confirmation of *Akodon serrensis* Thomas in Misiones, Argentina. *J. Mammalogy* **70** (2), 391–395.
- LOBATO, L.; CANTOS, G.; ARAUJO, B.; BIANCHI, N. O.; MERANI, M. S. (1982): Cytogenetics of

- South American akodont rodents (Cricetidae). VIII. *Akodon mollis*: a species with XY females and B-chromosomes. *Genetica* **57**, 199–205.
- MORITZ, C. (1986): The population biology of *Gehyra* (Gekkonidae): chromosome change and speciation. *Syst. Zool.* **35**, 46–67.
- PATHAK, S.; HSU, T. C.; ARRIGHI, F. E. (1973): Chromosomes of *Peromyscus* (Rodentia, Cricetidae) IV. The role of heterochromatin in karyotype evolution. *Cytogenet. Cell. Genet.* **12**, 315–326.
- REIG, O. A. (1984): Distribuição geográfica e história evolutiva dos roedores muroideos sulamericanos (Cricetidae: Sigmodontinae). *Rev. Brasileira de Genética* **7**, 333–365.
- (1986): Diversity patterns and differentiation of High Andean rodents. Oxford Univ. Press 404–438.
- (1987): An assessment of the systematics and evolution of the Akodontini, with the description of new fossil species of *Akodon* (Cricetidae, Sigmodontinae). *Fieldiana, Zool.* **496**, 347–399.
- REIG, O. A.; KIBLISKY, P. (1969): Chromosome multiformity in the genus *Ctenomys* (Rodentia, Octodontidae). *Chromosoma* **28**, 211–244.
- RODRIGUEZ, M.; MONTOYA, R.; VENEGAS, W. (1983): Cytogenetic analysis of some Chilean species of the genus *Akodon* Meyen (Rodentia, Cricetidae). *Caryologia* **36**, 129–138.
- SAHAR, E.; LATT, S. A. (1978): Enhancement of banding patterns in human metaphase chromosomes by energy transfer. *Proc. Natl. Acad. Sci.* **75**, 5650–5654.
- (1980): Energy transfer and binding competition between dyes used to enhance staining differentiation in metaphase chromosomes. *Chromosoma* **79**, 1–28.
- SBALQUEIRO, I. J.; MATTEVI, M. S.; OLIVEIRA, L. F. B. (1984): An $X_1X_1X_2X_2/X_1X_2Y$ mechanism of sex determination in a South American rodent, *Deltamys kempi* (Rodentia, Cricetidae). *Cytogenet. Cell Genet.* **38**, 50–55.
- SCHWEIZER, D. (1981): Counterstained enhanced chromosome banding. *Hum. Genet.* **57**, 1–14.
- SCHWEIZER, D.; EHRENDORFER, F. (1983): Evolution of C-band patterns in Asteraceae-Anthemideae. *Biol. Zentralbl.* **102**, 637–655.
- SEABRIGHT, M. (1971): A rapid banding technique for human chromosomes. *Lancet* **2**, 971–972.
- SPOTORNO, O. A.; FERNANDEZ, R. (1976): Chromosome stability in southern *Akodon* (Rodentia, Cricetidae). *Mam. Chrom. Newsletter* **17**, 13–14.
- STOCK, A. D. (1981): Chromosomal variation and constitutive heterochromatin in three porpoise species (genus *Stenella*). *Cytogenet. Cell. Genet.* **31**, 91–100.
- THOMAS, O. (1919): On small mammals collected by Sr. E. Budin in North-Western Patagonia. *Annals & Magazine of Natural History* **9**, 199–212.
- (1926): Two new mammals from Northern Argentina. *Annals & Magazine of Natural History* **9**, 311–312.
- (1927): On a further collection of mammals made by Sr. E. Budin in Patagonia. *Annals & Magazine of Natural History* **9**, 545–554.
- VITULLO, A. D.; MERANI, M. S.; REIG, O. A.; KAJON, A. E.; SCAGLIA, O.; ESPINOSA, M. B.; PEREZ-ZAPATA, A. (1986): Cytogenetics of South American akodont rodents (Cricetidae): new karyotypes and chromosomal banding patterns of Argentinian and Uruguayan forms. *J. Mammalogy* **67**, 69–80.
- YONENAGA, Y. (1972): Chromosomal polymorphism in the rodent *Akodon arviculoides* ssp. ($2n = 14$) resulting from two pericentric inversions. *Cytogenetics* **11**, 488–499.
- YONENAGA-YASSUDA, Y. (1979): New karyotypes and somatic and germ-cell banding in *Akodon arviculoides* (Rodentia, Cricetidae). *Cytogenet. Cell. Genet.* **23**, 241–249.
- YONENAGA, Y.; FROTA-PESSOA, O.; KASAHARA, S.; DE ALMEIDA, E. J. C. (1976): Cytogenetic studies on Brazilian rodents. *Ciencia e Cultura* **28**.
- YONENAGA, Y.; KASAHARA, S.; ALMEIDA, E. J. C.; PERACCHI, A. L. (1975): Chromosomal banding patterns in *Akodon arviculoides* ($2n = 14$), *Akodon* sp. ($2n = 24$ and 25), and two male hybrids with 19 chromosomes. *Cytogenet. Cell. Genet.* **15**, 388–399.

Authors addresses: MARIA A. BARROS, ROSA C. LIASCOVICH, OSVALDO A. REIG, GIBE, Dpto. de Cs. Biológicas, FCEyN, Universidad de Buenos Aires, Pabellón 2, 4to. piso, Ciudad Universitaria, Núñez, 1428 Buenos Aires, Argentina; LUZ GONZALEZ, Instituto de Evolución y Ecología, Universidad Austral de Chile, Valdivia, Chile; MARTHA S. LIZARRALDE, Centro Austral de Investigaciones Científicas, CONICET, Ushuaia, Argentina



Reig, Osvaldo A et al. 1990. "Banding pattern comparisons between Akodon iniscatus, and Akodon puer (Rodentia, Cricetidae)." *Zeitschrift für Säugetierkunde : im Auftrage der Deutschen Gesellschaft für Säugetierkunde e.V* 55, 115–127.

View This Item Online: <https://www.biodiversitylibrary.org/item/163746>

Permalink: <https://www.biodiversitylibrary.org/partpdf/191943>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder.

Rights Holder: Deutsche Gesellschaft für Säugetierkunde

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.