FURTHER RECORDS OF PHORIDAE (DIPTERA) FROM COCOONS OF CIMBICIDAE (HYMENOPTERA)

By A. D. LISTON*

I have collected a further two Cimbex femoratus (L.) cocoons containing 6 and 17 puparia of Megaselia giraudii (Egger) from soil under birches, Whitadder Valley, East Lothian, 13.iv.1979. 2 44 emerged on 20.iv.1979 and were determined as M. giraudii by Dr. R. H. L. Disney. Also one cocoon of Trichiosoma lucorum (L.), contianing 3 old puparia apparently identical with those of giraudii, was removed from a birch twig at Hopes Reservoir, East Lothian, 5.xii.1979.

As previously stated (Liston, 1979), the only reason why Phoridae do not infest Trichiosoma lucorum to such an extent as they do Climbex femoratus appears to be because of the site chosen for overwintering. This is in the soil for all Cimbex spp. and some Trichiosoma, but the cocoon is spun on a twig of the hostplant in T. lucorum and T. tibiale. The remains of any dead Trichiosoma euonymphs in their cocoons above ground desiccate easily while those of Cimbex in the soil are kept damp and are presumably more attractive to female phorids because of the odour given off. Only 1 Trichiosoma lucorum cocoon containing phorid puparia has been found in over 330 examined. Conversely, I have examined only about 25 Cimbex cocoons, 3 of which have contained puparia.

Dr. Disney's determination of giraudii confirms his earlier tentative determination made from puparia. Disney (1979) lists numerous rearing of giraudii from insects (chiefly Lepidoptera),

most of which appear to have been moribund.

References

Disney, R. H. L., 1979. Natural History notes on some British Phoridae (Diptera) with comments on a changing picture. Entomologist's Gaz., 30: 141-150.

Liston, A. D., 1979. On Phoridae (Diptera) from Sawfly cocoons

(Hym.: Symphyta). Ent. Rec., 91: 303-305.

* 99 Clermiston Road, Edinburgh, EH12 6UU.

EGGS LAID BY A DECAPITATED MOTH. - A Muslin Moth, Diaphora mendica Clerck, which I caught on the Downs near Beachy Head, Sussex, on 31st May 1982 was unfortunately decapitated as I enclosed it in a box. Arriving home an hour or two later I was surprised to find a few eggs had been laid on the box-side where it was suspended. More were laid later, on the floor of the box, as I watched. Its head was also on the floor. Egg-laying continued for two more days; there were two separate groups as well as the few on the side. The larvae emerged a few days later and were reared on Soft Sowthistle until 1st July when, having to leave home, I released them on Seaford Head. — Miss B. A. KNELLER, 9 Ashurst Road, Seaford, E. Sussex BN25 1AH.



Liston, Andrew. 1982. "Further records of Phoridae (Diptera) from cocoons of Cimbicidae (Hymenoptera)." *The entomologist's record and journal of variation* 94, 221–221.

View This Item Online: https://www.biodiversitylibrary.org/item/94409

Permalink: https://www.biodiversitylibrary.org/partpdf/196095

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: http://creativecommons.org/licenses/by-nc-sa/3.0/

Rights: https://biodiversitylibrary.org/permissions

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.