- Bird, F. T., 1954. The use of a virus disease in the biological control of the European Spruce Sawfly, Diprion hercyniae (Htg.). Bi-monthly Prog. Rpt., Div. Forest Biol., Canada Dept. Agr., 10 (1): 2-3.
- Bretherton, R. F., and Chalmers-Hunt, J. M., 1981. The Immigration of Lepidoptera to the British Isles in 1980, with an Account of the Invasion of the Painted Lady : *Cynthia cardui* (L.). *Ent. Rec.*, 93:47-54.
- Buckler, W., 1895. The Larvae of the British Butterflies and Moths., V1:Plate 99.
- Kettlewell, H. B. D., 1944. Temperature Experiments on the Pupae of *Heliothis peltigera* Schiff. Proc. S. Lond. ent. nat. *Hist. Soc.*, **1943-44:** 69-79.
- Neilson, M. M., 1965. Effects of a Cytoplasmic Polyhedrosis on Adult Lepidoptera. J. Inver. Pathol., 7:306-314.
- Pratt, C. R., 1981. A History of the Butterflies and Moths of Sussex. In press.
- Rivers, C. F., 1976. Diseases. In British Moths and Butterflies (J. Heath, ed.), Vol.1, 57-70.
 Sacharov, N. L., 1930. Studies in Cold Resistence of Insects.
- Sacharov, N. L., 1930. Studies in Cold Resistence of Insects. Ecology, 11:3:505-517.
- Tanada, Y., and Tanabe, A.M., 1965. Resistance of Galleria mellonella (Linnaeus) to the Tipula Tridescent Virus at High Temperatures. J. Inver. Pathol., 7:184-188.

CEUTHORHYNCHIDIUS RUFULUS DUF., ETC., (COL.: CURCU-LIONIDAE) IN S. E. LONDON. - As this little weevil is considered very local, and I have always found it to be so (occurring as a rule by odd specimens only), it may be worth noting that it belongs to the fauna of this suburban part of N. W. Kent; especially as the four localities given in the VCH list for that county (Fowler, 1908) are far east of here. I have taken solitary examples at roots of low herbage at Upper Charlton (my garden), Lower Charlton (Thamesside), and Kidbrooke (on a field bank). Its foodplant - species of Plantago - was not noticed at any of the places, but was probably present in all. Elsewhere I have met with C. rufulus at Mickleham Downs (Surrey) and Clayton Downs (Sussex), by sweeping. I may mention further that C. barnevillei Bris., the prettiest of the group, also occurs in my district but extremely sparingly, despite the abundance of its foodplant (Achillea millefolium L.); the localities are Blackheath (twice) and Charlton, at roots of yarrow. East of here I have taken it singly at Erith Marshes and Faversham Creek; and once swept several from its host-plant in a very restricted spot in the Lea Valley at Cheshunt, Herts. - A. A. ALLEN.

ORANGE-TIPS IN PEEBLESHIRE. – Two specimens of Anthocharis cardamines L. were seen by Mr. David G. Long of the Royal Botanic Garden, Edinburgh, on 12th May 1981 at Dawyck Garden, by a stream in the Tweed valley south of Stobo, VC78, and NT13. Both sexes were seen. This observation confirms the recent spread and establishment of this butterfly in Scotland. – A. G. LONG, 33, Windsor Crescent, Berwick upon Tweed, TD15.



Allen, Anthony Adrian. 1981. "Ceuthorhynchidius rufulus Duf., etc., (Col.: Curculionidae) in S.E. London." *The entomologist's record and journal of variation* 93, 141–141.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/94414</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/196177</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Smithsonian

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: Amateur Entomologists' Society License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.