XXI. *Addendum to Strepsiptera* (*p.* 86.) *By the Rev. William Kirby, F. L. S.*

Read February 16, 1813.

Having received a clearly distinct species of *Stylops*, though in a very mutilated state, from Mr. George B. Sowerby, F. L. S. I am desirous of giving some account of it by way of appendix to my Paper on *Strepsiptera*, and shall begin by describing it as far as its condition enabled me.

Stylops tenuicornis

Aterrima, oculis subsessilibus, antennis tenuioribus piceis: ramis linearibus, alis nigricantibus.

Long. Corp. $\frac{1}{2}$ lin. circiter.

A Dom. G. B. Sowerby Julio ineunti 1811, in Araneæ telâ lectus.

This species is not only much smaller than *Stylops Melittæ*, but it is also strikingly distinguished from it by its subsessile eyes and slender antennæ. From its mutilated state it was impossible to give a more detailed description.

In a specimen of the pupa of *Stylops Melittæ*, sent by the same gentleman for my inspection, the membrane for three-fourths of the length down the back appeared to be loose in the middle, and was divided into six segments by transverse impressed lines. This I imagine is the rugose part in Tab. IX. fig. 17.

When

Kirby, William. 1815. "XXI. Addendum to Strepsiptera (p. 86)." *Transactions of the Linnean Society of London* 11, 233–234.

https://doi.org/10.1111/j.1096-3642.1813.tb00055.x.