

DIAGNOSES

Specierum Novarum Carnioliae,

a cl. Jos. Stussiner Labacensi collectarum.

Auctore Dr. Osk. Boettger.

1. *Vitrina (Semilimax) truncata* n. sp.

F. J. Schmidt, Land- und Süßw.-Conch. in Krain, Laibach 1847 p. 7 (elongata, non Drap.)

Fr. Erjavec, Malak. Verh. v. Görz, Görz 1877 p. 18 (brevis, non Fér.)

Char. T. quasi intermedia inter *V. diaphanam* Stud. et *brevem* Fér., discrepans a *V. brevi* Fér., specie re vera persimili, t. semper minore, reniformi, angustiore, superne distincte plana nec convexa, spira paulo majore, ca. $\frac{2}{5}$ diam. majoris testae aequans, sutura profundiore, anfr. 2, ultimo minus dilatato, apert. minus obliqua, margine supero valde arcuatim protracto, superne ad suturam distincte recedente, dextro subtruncato, superne inferneque fere angulato, basali latiore, peripheriae longe parallelo, limbo membranaceo angustissimo.

Alt. $2\frac{1}{8}$, diam. min. $3\frac{1}{8}$, maj. $4\frac{5}{8}$ mm.

Hab. in voragine »Boltačev brezen« nec non in specu »Ihanška« regionis »Moraentscher Gegend« dictae Carnioliae superioris.

2. *Vitrina (Helicolimax) Carniolica* n. sp.

F. J. Schmidt, l. c. p. 7 (pellucida, non Drap.)

Fr. Erjavec, l. c. p. 18 (elliptica, non Brown).

Char. T. valde affinis *V. majori* Fér., sed spira paulo magis conica, $\frac{4}{7}$ diam. majoris testae aequans, anfractibus subangulatis, angulo peripherico inframediano,

supra et infra angulum planioribus, minus convexis, ultimo distincte magis dilatato, apert. majore, axi aperturae majore magis deflexa. Caeterum *V. majori* simillima.

Alt. 4, diam. min. $5\frac{3}{4}$, maj. $7\frac{1}{4}$ mm.

Hab. ad clivum, in quo castellum situm est Labacense nec non in voragine »Boltačev brezen«, in specu »Ihanška« aliisque locis regionis »Moraetscher Gegend« dictae Carnioliae superioris.

3. *Acme Stussineri* n. sp.

T. minutissima, minor graciliorque quam *A. perpusilla* Reinh., cylindrata, pellucida, vitrea, nitidissima; apex obtusissimus, fere subtruncatus. Anfr. 5 pro longitudine testae alti, fere aequales, superi convexiusculi, inferi planiores, sutura submarginata, sat impressa disjuncti, laeves, ultimus $\frac{1}{4}$ altitudinis testae aequans, ante aperturam callo annulari non cinctus. Apert. obliqua, basi recedens, late ovata, marginibus simplicibus, hebetibus, callo tenuissimo junctis, columellari subincrassato, reflexiusculo, dextro regulariter curvato-porrecto.

Alt. fere $1\frac{1}{2}$, diam. $\frac{1}{2}$ mm.

Hab. in voragine »Volaufov Kevder« regionis dictae »Moraetscher Gegend« Carnioliae superioris, 7 specimina.

Observ. Minima et gracillima omnium specierum adhuc cognitarum generis *Acme* (*Acicula*) Hartm.



Boettger, O. 1884. "Diagnoses Specierum Novarum Carnioliae,"
Nachrichtenblatt der Deutschen Malakozoologischen Gesellschaft 16, 184–185.

View This Item Online: <https://www.biodiversitylibrary.org/item/97105>

Permalink: <https://www.biodiversitylibrary.org/partpdf/199318>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.