

Die vertikale Verbreitung der Lepidopteren in der Columbischen Ost-Cordillere.

(Schluß. Vgl. Nr. 11.)

Von *A. H. Fassl*, Teplitz.

	Westen	500	1000	1500	2000	2500	3000	3500	3000	2500	2000	1500	1000	500	Osten	
		0	500	1000	1500	2000	2500	3000	3500	3000	2500	2000	1500	1000	500	0
Anteros gyas Cr., renaldus Stoll, carus Stgr., cupris Hw., Mesenopsis melanochlora Gdm., Symmachia domitianus F., Mesene phareus Cr., monostigma Erichs., telephus Cr., Charis saphirina Stgr., macularia Bsd., holosticta Gdm., Panara barsacus Westw., Nymphidium lycorias Hw., grande Gdm., Theope publius Fld., Aricoris jansonii Btlr.																
Anteros chrysus Cr., formosus Cr., Charis monogramma Bat., Lemonias crispus Cr., Thisbe irenaea Cr., Nymphidium philone Godt., abaris Cr., Pandema pasiphae Cr., Stalachtis phlegia Cr., Tharops menander, isthmica Godm., Anatole zygia Hübn.																
Mesosemia mevania Hw., Diorhina pseacas Saund., Siseme alectryo Bat.																
Mesosemia maenades Hw., Imelda glauconia, Hyphilaria anophthalma Flg., Necyria manco Saund.																
Syrmatia aethiops Stgr., dorilas Cr., Esthemopsis strigosa Stgr., Lymnas passiena Hw., thyatira Hw., Hades noctula Westw., Necyria saundersii Hw., Ancyluris sogamuxi Fassl., Ithomeis corinna Stgr., Crocozona capicaris Butl., Aricoris phythioides Btl., Compso-tera cascella Hw.																
Lycaenidae überhaupt:																
Thecla nobilis H. Sch., imperialis Cr., gibberosa Hw., linus F., mavors Hübn., marsyas L.																
Thecla tuneta Hw., cypria Hbn. paphia Fld., balias Gdm., thalesa Hw., barajo Reak., arageus Cr., themon Cr., calus Hw., gabatha Hw., Theoroma eumenia Hw.																
Thecla pholus Cr., phaleros Cr.																
Thecla coronata Hw., telemus Cr., atys Cr.																
Thecla cadmus Fld., aegides Fld., Micandra platyptera Fld.																
Thecla polibetes Cr.																
Thecla loxurina Fld., commodus F.																
Hesperidae überhaupt:																
Erycides gnetus F.																
Telegonus actylos Cr., Erycides gaudialis Hw., maculosa Hw., Achlyodes chlorocephala Latr.																
Spathilepia thamyroides Feld., Entheus talaus, coeleste D. H., gentius Cr., Pyrrhopyga hadora Hw., Erycides cloanthes Latr., ochrope Ploetz, Carystus sergestes Cr., cymica Swains.																
Carystus catargyrea Fld., vulpina Fld., Erycides spatiosa Hw.																
Erycides sela Hw., Hesperia claudianus Latr.																
Pamphila kirschi Weym., Ancyloxypha melanoneura Fld., Mnes-theus ludens Feld.																
Castniidae überhaupt:																
Castnia evalthe F.																
Castn. atymnus Dalm., wagneri Busch., cycna Westw., diva Butl., cacica H. Sch.																
Castn. licus Drury, papilionaris Westw.																

	Westen	0	500	1000	1500	2000	2500	3000	3500	3000	2500	2000	1500	1000	500	Osten
<i>Rhescyntis norax</i> Druce, <i>Arsenura</i> arcaei Druce, <i>Dysdaemonia boreas</i> Cr., <i>tamerlana nobilis</i> Schs., <i>Copiopteryx semiramis</i> Cr., <i>Copaxa multifenestrata</i> H. Sch., <i>Lenomia concordia</i> Druce, <i>electra</i> Druce, <i>obliqua</i> Wlk., <i>aeneides</i> Druce, <i>Oxytenis honesta</i> Stoll, <i>Drepanopteryx sricula</i> Weym., <i>Attacus bolivar</i> M. u. W., <i>hesperus</i> L., <i>Automeris titania</i> Fld., <i>zozine</i> Druce, <i>egeus</i> Cr., <i>theresa</i> Weym., <i>euryopa</i> Möschl., <i>angulatus</i> Conte, <i>hübneri</i> Bsd., <i>Dirphia menander</i> Druce, <i>ursina</i> Wlk., <i>pomona</i> Fld., <i>Grammapella fuscovenosa</i> Weym.																
<i>Dysdaemonia timur</i> Weym., <i>Attacus belis</i> Wlk., <i>Automeris liberia</i> Cr., <i>illustris</i> Wlk., <i>venosus</i> Btl., <i>aulestes</i> H. Sch., <i>fassli</i> Weym., <i>Dirphia tarquinia</i> Cr.																
<i>Dirphia eumedide</i> Stoll, <i>Dysdaemonia pluto</i> Westw.																
<i>Dirphia semirosea</i> Wlk., <i>mota</i> Druce, <i>Oxytenis fulvofasciata</i> Weym.																
<i>Sagana zapatoza</i> Westw., <i>Attacus aricia</i> Wlk., <i>Automeris leucane</i> Hübn.																
<i>Attacus zacateca</i> Westw., <i>Automeris flexilineata</i> Dgn.																
<i>Automeris saturniata</i> Fld., <i>Dirphia litura</i> Wlk., <i>Hemileuca venosa</i> Wlk.																
Ceratocampidae überhaupt:																
<i>Eacles magnifica</i> Grote, <i>Sissphinx molina</i> Stoll																
<i>Eacles ormondei</i> Schs., <i>Citheronia mexicana</i> Grote, <i>Adelocephala fallax</i> Bsd., <i>columbia</i> Schs., <i>anthoni</i> H. Sch., <i>argyrocantha</i> Bsd.																
<i>Adeloceph. walkeri</i> Grote, <i>Eacles fenestrata</i> Rotsch.																
Notodontidae überhaupt:																
<i>Crinodes ritsemae</i> Bull., <i>beskei</i> Hübn., <i>Hapigia raatzi</i> notricornis Guen., <i>smerinthoides</i> Wlk., <i>Colas apulus</i> Cr., <i>Ophytes cyllota</i> Druce, <i>Rhuda phocula</i> Zelica thalassina																
<i>Rosema apicalis</i> Wlk.																
<i>Xyleutes putridus</i> Perch., <i>xylotribus</i> H. Sch., <i>Therinia transversaria</i> Druce, <i>lactucina</i> Cr.																
<i>Epia muscosa</i> , <i>Stenele translata</i> Wlk.																
Noctuidae überhaupt:																
<i>Orthia augias</i> H. Sch.																
<i>Palindia dominicata</i> Guen., <i>alabastaria</i> , <i>Gonodonta pyrgo</i> Cr., <i>Ophideres procius</i> L., <i>serpentera</i> Druce																
<i>Vespola caeruleifera</i> Wlk.																
<i>Plusia rogationis</i> Gn, <i>Darceta hesperina</i> H. S., <i>Palindia ilyrias</i> Cr.																
<i>Erebus odora</i> L.																
Geometridae überhaupt:																
<i>Urania fulgens</i> Wlk., <i>Sangala beata</i> Wlk., <i>Ametrix netrix</i> Cr., <i>Amblychia scolopacea</i> Drury, <i>Trigodes niobe</i> Druce, <i>musivaria</i> H. Sch.																
<i>Urania leilus</i> L., <i>Micropus longalis</i> Hbn., <i>Racheaspila albicoma</i> Fld.																
<i>Sangala gloriosa</i> Wlk., <i>Diopthis hesperioides</i> Wlk.																
<i>Diplochroa bicentraria</i> H. Sch., <i>Heterusia placilla</i> Druce																
<i>Siga liris</i> Cr.																



Fassl, Anton Heinrich. 1918. "Die vertikale Verbreitung der Lepidopteren in der Columbischen Ost-Cordillere." *Entomologische Rundschau* 35, 48–50.

View This Item Online: <https://www.biodiversitylibrary.org/item/38724>

Permalink: <https://www.biodiversitylibrary.org/partpdf/201815>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.