

- ANDREWS, R. M. 1979. The lizard *Corytophanes cristatus*: an extreme "sit-and-wait" predator. *Biotropica*, in press.
- ANDREWS, R. M. AND A. S. RAND. 1974. Reproductive effort in anoline lizards. *Ecology*, 55: 1317-1327.
- BEARD, J. S. 1949. *The Natural Vegetation of the Windward and Leeward Islands*. Oxford, Clarendon Press. 142 pp.
- BENNETT, A. F. AND W. R. DAWSON. 1976. Metabolism. In Gans, C. (ed.), *Biology of the Reptilia*, Vol. 5 (Physiology A), New York, Academic Press, pp. 127-223.
- BOND, JAMES. 1961. *Birds of the West Indies*. Boston, Houghton Mifflin Co. 256 pp.
- BRACH, V. 1976. Habits and food of *Anolis equestris* in Florida. *Copeia*, 1976: 187-189.
- BRATTSTROM, B. H. 1974. The evolution of reptilian social behavior. *Amer. Zool.*, 14: 35-49.
- CAMPBELL, H. W. 1973. Ecological observations on *Anolis lionotus* and *Anolis poecilopus* (Reptilia, Sauria) in Panama. *Amer. Mus. Novitates*, 2516: 1-29.
- CASE, T. J. 1975. Species numbers, density compensation, and colonizing ability of lizards on islands in the Gulf of California. *Ecology*, 56: 3-18.
- CASE, T. J. 1978. A general explanation for insular body size trends in terrestrial vertebrates. *Ecology*, 59: 1-18.
- COLE, L. C. 1954. The population consequences of life history phenomena. *Quart. Rev. Biol.*, 29: 103-137.
- CREWS, DAVID. 1973. Behavioral correlates to gonadal state in the lizard, *Anolis carolinensis*. *Hormones and Behavior*, 4: 307-313.
- ECHELLE, A. A., A. F. ECHELLE, AND H. S. FITCH. 1971. A new anole from Costa Rica. *Herpetologica*, 27: 354-362.
- FENCHEL, T. 1974. Intrinsic rate of natural increase: the relationship with body size. *Oecologia*, 14: 317-326.
- FITCH, H. S. 1972. Ecology of *Anolis tropidolepis* in Costa Rican cloud forest. *Herpetologica*, 28: 10-21.
- FITCH, H. S. 1973a. Observations on the population ecology of the Central American iguanid lizard, *Anolis cupreus*. *Carib. Jour. Sci.*, 13: 215-230.
- FITCH, H. S. 1973b. Population structure and survivorship in some Costa Rican lizards. *Occas. Papers Univ. Kans. Mus. Natur. Hist.*, 18: 1-41.
- FITCH, H. S. 1973c. A field study of Costa Rican lizards. *Univ. Kans. Sci. Bull.*, 50: 39-126.
- FITCH, H. S. 1976. Sexual size differences in the mainland anoles. *Occas. Papers Univ. Kans. Mus. Natur. Hist.*, 50: 1-21.
- FITCH, H. S., A. A. ECHELLE, AND A. F. ECHELLE. 1972. Variation in the Central American iguanid lizard, *Anolis cupreus*, with the description of a new subspecies. *Occas. Papers Univ. Kans. Mus. Natur. Hist.*, 8: 1-20.
- FITCH, H. S., A. F. ECHELLE, AND A. A. ECHELLE. 1976. Field observations on rare or little known mainland anoles. *Univ. Kans. Sci. Bull.*, 51: 91-128.
- FITCH, H. S. AND R. W. HENDERSON. 1976. A field study of the rock anoles (Reptilia, Lacertilia, Iguanidae) of southern Mexico. *J. Herpetol.*, 10: 303-311.
- FLEMING, T. H. AND R. S. HOOKER. 1975. *Anolis cupreus*: the response of a lizard to tropical seasonality. *Ecology*, 56: 1243-1261.

- GORMAN, G. C. AND R. HARWOOD. 1977. Notes on population density, vagility, and activity patterns of the Puerto Rican grass lizard, *Anolis pulchellus* (Reptilia, Lacertilia, Iguanidae). J. Herpetol., 11: 363-368.
- GORMAN, G. C. AND P. LICHT. 1974. Seasonality in ovarian cycles among tropical *Anolis* lizards. Ecology, 55: 360-369.
- GORMAN, G. C. AND P. LICHT. 1975. Differences between the reproductive cycles of sympatric *Anolis* lizards on Trinidad. Copeia, 1975: 332-336.
- HENDERSON, R. W. AND H. S. FITCH. 1975. A comparative study of the structural and climatic habitats of *Anolis sericeus* (Reptilia: Iguanidae) and its syntopic congeners at four localities in southern Mexico. Herpetologica, 31: 459-471.
- HICKMAN, J. C. 1975. Environmental unpredictability and plastic energy allocation strategies in the annual *Polygonum cascadenae* (Polygonaceae). J. Ecology, 63: 689-702.
- HIRSHFIELD, M. F. AND D. W. TINKLE. 1975. Natural selection and the evolution of reproductive effort. Proc. Natl. Acad. Sci., 72: 2227-2231.
- HODGE, W. H. 1954. Flora of Dominica, B.W.E. Lloydia, 17: 1-238.
- HOLDRIDGE, L. R. 1967. Life Zone Ecology. Tropical Science Center, San Jose, Costa Rica, rev. ed. 124 pp.
- JACKSON, J. F. 1973. Notes on the population biology of *Anolis tropidonotus* in a Honduran highland pine forest. J. Herpetol., 7: 309-311.
- JANZEN, D. H. 1973a. Sweep samples of tropical foliage insects: description of study sites, with data on species abundances and size distributions. Ecology, 54: 659-686.
- JANZEN, D. H. 1973b. Sweep samples of tropical foliage insects: effects of seasons, vegetation types, elevation, time of day, and insularity. Ecology, 54: 687-702.
- JANZEN, D. H. AND T. W. SCHOENER. 1968. Differences in insect abundance and diversity between wetter and drier sites during a tropical dry season. Ecology, 49: 96-110.
- JENSSEN, T. A. 1970. The ethoecology of *Anolis nebulosus* (Sauria, Iguanidae). J. of Herpetol., 4: 1-38.
- JENSSEN, T. A. 1973. Shift in the structural habitat of *Anolis opalinus* due to congeneric competition. Ecology, 54: 863-869.
- LAZELL, J. D., JR. 1972. The Anoles (Sauria, Iguanidae) of the Lesser Antilles. Bull. Mus. Comp. Zool., 143: 1-115.
- LESTON, D. 1970. Entomology of a cocoa farm. Ann. Rev. Entomol. 15: 237-294.
- LICHT, P. AND G. C. GORMAN. 1970. Reproductive and fat cycles in Caribbean *Anolis* lizards. Univ. Calif. Publ. in Zool., 95: 1-52.
- LISTER, B. C. 1976. The nature of niche expansion in West Indian *Anolis* lizards I: Ecological consequences of reduced competition. Evol., 30: 659-676.
- MACARTHUR, R. H. AND E. O. WILSON. 1967. The Theory of Island Biogeography. Princeton Univ. Press. 203 pp.
- MACARTHUR, R. H., J. M. DIAMOND, AND J. R. KARR. 1972. Density compensation in island faunas. Ecology, 53: 330-342.
- MARTEN, G. G. 1970. A regression method for mark-recapture estimation of population size with unequal catchability. Ecology, 51: 291-295.

- MEYER, J. R. 1968. Distribution and variation of the Mexican lizard, *Anolis barkeri* Schmidt (Iguanidae), with a redescription of the species. *Copeia*, 1968: 89-95.
- MIYATA, K. 1977. Report, In E. E. Williams (ed.), The Third *Anolis* Newsletter, Museum of Comparative Zoology, Harvard University, Cambridge.
- MURPHY, G. I. 1968. Pattern in life history and the environment. *Amer. Natur.*, 102: 391-403.
- MYERS, C. W. 1971. Central American lizards related to *Anolis pentaprion*: Two new species from the Cordillera de Talamanca. *Amer. Mus. Nov.*, No. 2471: 1-40.
- NICHOLS, J. D., W. CONLEY, B. BATT, AND A. R. TIPTON. 1976. Temporally dynamic reproductive strategies and the concept of r- and K-selection. *Amer. Natur.*, 110: 995-1005.
- PIANKA, E. R. 1970. On r and K selection. *Amer. Natur.*, 104: 592-597.
- PIANKA, E. R. 1972. r and K selection or b and d selection? *Amer. Natur.*, 106: 581-588.
- RAND, A. S. 1961. Ecology, behavior, and morphology of anoline lizards in Puerto Rico. Ph.D. Thesis, Harvard Univ., Cambridge, Mass.
- RAND, A. S. 1967. Ecology and social organization in the iguanid lizard *Anolis lineatopus*. *Proc. U.S. Natl. Mus.*, 122: 1-79.
- RAND, A. S. AND R. M. ANDREWS. 1975. Adult color dimorphism and juvenile pattern in *Anolis cuvieri*. *J. Herpetol.*, 9: 257-260.
- RAND, A. S. AND S. S. HUMPHREY. 1968. Interspecific competition in the tropical rain forest: ecological distribution among lizards at Belem, Para. *Proc. U.S. Natl. Mus.*, 125: 1-17.
- RAND, A. S. AND E. E. WILLIAMS. 1969. The anoles of La Palma: aspects of their ecological relationships. *Breviora*, 327: 1-19.
- RECHER, H. F. 1970. Population density and seasonal changes of the avifauna in a tropical forest before and after gamma irradiation. In Odum, H. T. (ed.), *A Tropical Rain Forest*, Springfield, VA, US AEC: E69-E86.
- RUIBAL, R. AND R. PHILIBOSIAN. 1970. Eurythermy and niche expansion in lizards. *Copeia*, 1970: 645-653.
- RUIBAL, R. AND R. PHILIBOSIAN. 1974a. The population ecology of the lizard *Anolis acutus*. *Ecology*, 55: 525-537.
- RUIBAL, R. AND R. PHILIBOSIAN. 1974b. Aggression in the lizard *Anolis acutus*. *Copeia*, 1974: 349-357.
- SAGE, R. D. 1974. The structure of lizard faunas: comparative biologies of lizards in two Argentina deserts. Ph.D. Thesis, Univ. of Texas, Austin.
- SCHOENER, T. W. 1967. The ecological significance of sexual dimorphism in size in the lizard *Anolis conspersus*. *Science*, 155: 474-477.
- SCHOENER, T. W. 1968. The *Anolis* lizards of Bimini: resource partitioning in a complex fauna. *Ecology*, 49: 704-726.
- SCHOENER, T. W. 1969a. Models of optimal size for solitary predators. *Amer. Natur.*, 103: 277-313.
- SCHOENER, T. W. 1969b. Size patterns in West Indian *Anolis* lizards. I. Size and species diversity. *Syst. Zool.*, 18: 386-401.

- SCHOENER, T. W. 1977. Competition and the niche. *In* Gans, C. and D. W. Tinkle (Eds.), *Biology of the Reptilia*, New York, Academic Press: pp. 35-136.
- SCHOENER, T. W. AND G. C. GORMAN. 1968. Some niche differences in three Lesser Antillean lizards of the genus *Anolis*. *Ecology*, 49: 819-830.
- SCHOENER, T. W. AND D. H. JANZEN. 1968. Notes on environmental determinants of tropical versus temperate insect size patterns. *Amer. Natur.*, 102: 207-224.
- SCHOENER, T. W. AND A. SCHOENER. 1971a. Structural habitats of West Indian *Anolis* lizards. II. Puerto Rico Uplands. *Breviora*, 375: 1-39.
- SCHOENER, T. W. AND A. SCHOENER. 1971b. Structural habitats of West Indian *Anolis* lizards. I. Lowland Jamaica. *Breviora*, 368: 1-53.
- SCHOENER, T. W. AND A. SCHOENER. 1978. Inverse correlation of survival in lizards with island size and avifaunal richness. *Nature*, 274: 685-687.
- SCHWARTZ, A. AND R. THOMAS. 1975. Checklist of West Indian amphibians and reptiles. Carnegie Museum of Natural History, Special Publication No. 1, 216 pp.
- SCOTT, N. J., D. E. WILSON, C. JONES, AND R. M. ANDREWS. 1976. The choice of perch dimensions by *Anolis* lizards. *J. Herpetol.* 10: 75-84.
- SEBER, G. A. F. 1973. The estimation of animal abundance and related parameters. London, Griffin. 506 pp.
- SEXTON, O. J. 1967. Population changes in a tropical lizard *Anolis limifrons* on Barro Colorado Island, Panama Canal Zone. *Copeia*, 1967: 219-222.
- SEXTON, O. J., J. BAUMAN, AND E. ORTLEB. 1972. Seasonal food habits of *Anolis limifrons*. *Ecology*, 53: 182-186.
- SEXTON, O. J. AND K. M. BROWN. 1977. The reproductive cycle of an iguanid lizard *Anolis sagrei*, from Belize. *J. Natur. Hist.*, 11: 241-250.
- SEXTON, O. J., E. P. ORTLEB, L. M. HATHAWAY, R. E. BALLINGER, AND P. LICHT. 1971. Reproductive cycles of three species of anoline lizards from the Isthmus of Panama. *Ecology*, 52: 201-215.
- SIEGAL, S. 1956. Nonparametric statistics. McGraw-Hill. 312 pp.
- SLOBODKIN, L. G., F. E. SMITH, AND N. G. HAIRSTON. 1967. Regulation in terrestrial ecosystems, and the implied balance of nature. *Amer. Natur.*, 101: 109-124.
- SORIA, J., O. ESQUIVEL, L. A. PAREDES, AND R. DIAZ-ROMEY. 1969. Finca La Lola. *Cacao*, 14: 1-42.
- SORIANO-RESSY, M., A. P. DESMARAIS, AND J. W. PEREZ. 1970. A comparison of environments of rain forests in Dominica, British West Indies, and Puerto Rico. *In* Odum, H. T. (Ed.), *A Tropical Rain Forest*, Springfield, VA, US AEC: B329-B346.
- SOUTHWOOD, T. R. E. 1966. *Ecological Methods With Particular Reference to the Study of Insect Populations*. London, Methuen. 391 pp.
- STAMPS, J. A. 1975. Courtship patterns, estrus periods and reproductive condition in a lizard, *Anolis aeneus*. *Physiology and Behavior*, 14: 531-535.
- STAMPS, J. A. 1976. Egg retention, rainfall and egg laying in a tropical lizard, *Anolis aeneus*. *Copeia*, 1976: 759-764.
- TALBOT, J. J. 1977. Habitat selection in two tropical anoline lizards. *Herpetologica*, 33: 114-123.

- TINKLE, D. H., H. M. WILBUR, AND S. G. TILLEY. 1970. Evolutionary strategies in lizard reproduction. *Evol.*, 24: 55-74.
- TOFT, C. A. 1976. Partitioning of food in a community of tropical frogs. Ph.D. Thesis, Princeton Univ., New Jersey.
- TRIVERS, R. L. 1972. Parental investment and sexual selection. In Campbell, B. (Ed.), *Sexual Selection and the Descent of Man*, Chicago, Aldine Publishing Co.: pp. 136-179.
- TRIVERS, R. L. 1976. Sexual selection and resource-accruing abilities in *Anolis garmani*. *Evol.*, 30: 253-269.
- TURNER, F. B. AND C. S. GIST. 1970. Observations of lizards and tree frogs in an irradiated Puerto Rican forest. In Odum, H. T. (ed.), *A Tropical Rain Forest*, Springfield, VA, US AEC: E25-49.
- VANZOLINI, P. E. AND E. E. WILLIAMS. 1970. South American anoles: The geographic differentiation and evolution of the *Anolis chrysolepis* species group (Sauria, Iguanidae). *Arquivos de Zoologia*, S. Paulo, 19: 125-298.
- WIEGERT, R. G. AND D. F. OWEN. 1971. Trophic structure, available resources, and population density in terrestrial vs. aquatic ecosystems. *J. Theoret. Biol.*, 30: 69-81.
- WILLIAMS, E. E. 1972. The origin of faunas. Evolution of lizard congeners in a complex island fauna: A trial analysis. *Evol. Biol.*, 6: 47-90.

B R E V I O R A

MAR 18 1985
HARVARD
UNIVERSITY

Museum of Comparative Zoology

US ISSN 0006-9698

CAMBRIDGE, MASS.

JULY 31, 1979

NUMBER 455

**A NEW PALEOCENE PALAEANODONT AND THE
ORIGIN OF THE METACHEIROMYIDAE (MAMMALIA)**

KENNETH D. ROSE¹

ABSTRACT. A new genus and species of palaeanodont, *Propalaeanodon schaffi*, from the late Paleocene (late Tiffanian) Polecat Bench Formation of the Bighorn Basin, Wyoming, is the oldest and most primitive known metacheiromyid. *Propalaeanodon* foreshadows *Palaeanodon* in the reduction of its postcanine teeth. It resembles epoicotheriids in the presence of postcanines along the entire length of the horizontal ramus, and resembles the slightly older *Amelotabes*, in particular, in the retention of seven postcanines. *Propalaeanodon* thus provides additional evidence for uniting the Metacheiromyidae and the Epoicotheriidae in the suborder Palaeanodonta, and it strongly suggests that the Metacheiromyidae evolved from the Epoicotheriidae, probably during the Tiffanian.

Two humeri from the same stratigraphic level as the holotype are tentatively referred to *Propalaeanodon*. They possess specializations typical of Eocene palaeanodonts, including a prominent deltopectoral crest and a large supinator crest. These features indicate that *Propalaeanodon* was, already in the late Tiffanian, a highly specialized fossorial animal.

The systematic position of the Palaeanodonta is uncertain. They have been considered to be related to the orders Xenarthra or Pholidota or both by all recent students and, indeed, they are the only known fossil group that evolved specializations similar to those in these two orders. Although it is possible that palaeanodonts were an early radiation of probably myrmecophagous, fossorial mammals merely convergent to these orders, the evidence now available enhances the probability that the three groups share a special relationship.

INTRODUCTION

The edentate-like members of the rare mammalian suborder Palaeanodonta have long been considered early relatives of the Xenarthra, the Pholidota, or both. As now known, they are restricted to the earlier Tertiary of western North America. Palaeanodonts are classified in two families, the Epoicotheriidae and the

¹Museum of Paleontology, University of Michigan, Ann Arbor, Michigan 48109.

Metacheiromyidae. To date, nine genera and fifteen species have been recognized, several of which were described quite recently (West, 1973; Rose et al., 1977; Rose, 1978). They range in age from Tiffanian (late Paleocene) to Orellan (middle Oligocene).

Recently the dentary of a primitive new palaeonodont was discovered by Charles Schaff, Museum of Comparative Zoology, in late Tiffanian rocks in the northern Bighorn Basin, Wyoming. The oldest and most primitive metacheiromyid, it sheds new light on the origin of this peculiar family and further documents the unity of the Palaeonodonta. Two palaeonodont humeri, collected much earlier from the same stratigraphic interval, probably represent the same new species.

Abbreviations of institutional names cited herein are as follows:

AC Pratt Museum, Amherst College, Amherst, Massachusetts.

AMNH American Museum of Natural History, New York.

MCZ Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts.

PU Princeton University Museum, Princeton, New Jersey.

UM University of Michigan Museum of Paleontology, Ann Arbor, Michigan.

SYSTEMATIC PALEONTOLOGY

Suborder PALAEANODONTA Matthew, 1918

Family METACHEIROMYIDAE Wortman, 1903

PROPALAEANODON, gen. nov.

Type species. Propalaeonodon schaffi, sp. nov.

Included species. Type only.

Distribution. Late Paleocene (late Tiffanian) of the Bighorn Basin, Wyoming.

Diagnosis. Only metacheiromyid with seven postcanine teeth that extend to the back of the horizontal ramus. All postcanines single-rooted except P_4 , which is two-rooted. Postcanines small, peglike, separated by short diastemata, and with rounded crowns, in these features resembling *Palaeonodon*; but crowns higher and more pointed than in *Palaeonodon*. Dentary slender, slightly smaller than in *Palaeonodon ignavus*; deepest beneath first postcanine (P_1), shallowest beneath last postcanine (M_3), as in *Palaeonodon*. Medial buttress prominent but less so than in *Palaeonodon*.

Etymology. In allusion to its resemblance to, lower stratigraphic level than, and probable relationship to *Palaeanodon*.

PROPALAEANODON SCHAFFI, sp. nov.

Fig. 1A and B, 2A, and 3B

Holotype: MCZ 20122, left dentary with P_{2-3} , alveoli for P_1 , P_4 , M_{1-3} , and posterior margin of canine alveolus.

Hypodigm. Holotype and tentatively PU nos. 13928, 13929.

Horizon and Locality. Latest Paleocene (late Tiffanian), "Silver Coulee beds" of Polecat Bench Formation (of Jepsen, 1940), Jepsen Valley Quarry, NW $\frac{1}{4}$, Sec. 27, T.57N., R.100W., Park County, Wyoming.




Figure 1. Lateral (A) and dorsal (B) views of holotype of *Propalaeanodon schaffi*, MCZ 20122, left dentary with second and third postcanines (P_{2-3}). Dorsal view of left dentary of *Palaeanodon* sp. (C), showing extended edentulous part of ramus behind fifth postcanine; reconstructed from UM 63611 and 66243.




Figure 2. Medial views of left dentaries of palaeonodonts. (A) *Propalaeonodon schaffi*, holotype, MCZ 20122. (B) *Palaeonodon* sp., reconstruction based on UM 63611 and 66243. (C) *Amelotabes simpsoni*, holotype, PU 14855 (right dentary, reversed).

Diagnosis. Only known species of the genus; measurements in Table 1.

Etymology. For Charles R. Schaff, expert collector and preparator, who discovered the holotype.

Description. Only two teeth are preserved in the holotype, although the dentary is nearly complete and provides much information about the mandibular morphology of *Propalaeonodon schaffi*. The horizontal ramus is intact as far forward as the posterior margin of the canine alveolus. This alveolar margin reveals that the canine was large, as in other palaeonodonts, and its

root robust, a feature also indicated by thickening of the dentary anteriorly.

The dentary contained seven postcanine teeth, presumably four premolars and three molars, which were greatly modified from the primitive eutherian condition. Seven is a greater number of teeth than in any other metacheiromyid, indeed more than in any other palaeanodont except *Amelotabes*, which also has seven (Rose, 1978). *Alocodontulum*, in which the lower dentition is unknown, has seven upper postcanines (Rose et al., 1977, 1978), and thus probably had seven lowers as well. The postcanines of *Propalaeonodon* occur along the entire length of the horizontal ramus, as in




Figure 3. Left humeri of palaeanodonts. Scale is 10 mm. (A) Early Eocene metacheiromyid *Palaeonodon ignavus*, based on AMNH 15088 and 16832; after Matthew (1918), but with corrected orientation of proximal end. (B) ?*Propalaeonodon schaffi*, PU 13928. (C) Early Eocene epoicotheriid *Pentapassalus woodi*, part of holotype, AC 2766. Note that this species is referable to *Pentapassalus* (Rose, 1978), not to *Palaeonodon* (Guthrie, 1967; Emry, 1970). The supinator crests in (A) and (C) are incomplete and have not been reconstructed because of the variable development of this crest in different taxa of palaeanodonts; they were in any case larger than depicted.

TABLE 1

Dimensions of the holotype of *Propalaeonodon schaffi*, MCZ 20122.
(Measurements are in mm to nearest .05 mm)

	<i>Length</i>	<i>Breadth</i>
P ₁ alveolus	1.35	0.85
P ₂ alveolus	damaged	
P ₂ crown	1.25	1.00
P ₃ alveolus	2.40	1.15
P ₃ crown	1.80	1.10
P ₄ alveolus	2.80	1.10
M ₁ alveolus	2.00	1.15
M ₂ alveolus	2.00	1.15
M ₃ alveolus	1.60	1.00
P ₁ -M ₃	17.10 (approx.)	
Depth of mandible, linguallly at P ₁ : 6.10 (approx.)		
Depth of mandible, linguallly at M ₃ : 4.80		

epoicotheriids but not in later metacheiromyids, in which the back of the horizontal ramus is edentulous. All teeth in *Propalaeonodon* are single-rooted except P₄, which has two roots (probably a primitive feature). Judging from the alveoli, M₁ and M₂, and possibly P₃, had single roots that were faintly bilobate. The roots of all postcanines were tapered and closed, as in *Palaeonodon*. The epoicotheriids *Amelotabes* (Fig. 2C) and *Tubulodon* are similar in having a two-rooted P₄, but in these taxa several other postcanines are also two-rooted. In *Pentapassalus* M₁ has a bilobate root but P₄ is single-rooted (Gazin, 1952). The postcanines in *Palaeonodon* are all single-rooted (Fig. 1C, 2B).

The crowns of P₂ and P₃ are preserved. P₂ has become dislodged from its alveolus as a result of damage to the front of the jaw and appears higher than P₃. Both teeth are peglike with rounded crowns apparently devoid of enamel. In these characters they closely resemble the postcanines of *Palaeonodon* (Fig. 1C, 2B and Rose, 1978), although they are slightly larger, particularly longer, higher crowned, and less rounded. As in *Palaeonodon* they show no cusp pattern on the crowns, nor is there any evidence of direct occlusion with the upper teeth. Occlusal wear is discernible in molars and even in premolars of some epoicotheriids. The alveolus of P₄ indicates

that it was the largest cheek tooth, a feature in common with *Amelotabes* but in contrast to *Palaeanodon*. There is a slight size reduction from P_4 to M_3 .

All postcanines are separated by short diastemata. At the alveolar margin, the alveoli of P_{2-4} slightly exceed the size of the roots they contain; this probably obtained for the molars as well. These traits are characteristic also of *Palaeanodon*, some advanced epoicotheriids, and many toothed xenarthrans.

The ascending ramus and condyle of *Propalaeanodon* resemble those of *Pentapassalus pearcei* and *Metacheiromys tatusia* (Gazin, 1952; Simpson, 1931). The coronoid process is slightly smaller than in contemporary proteutherians (*sensu* Butler, 1972) and rises at an angle of about 110° to the ventral border of the mandible. The condyle, which is narrower transversely than in most toothed xenarthrans and contemporary proteutherians, is situated relatively low but above the level of the tooth row. Its convex articular surface is directed dorsally and backward, as in *P. pearcei*. The mandibular foramen is also relatively low, being situated on the posterior projection of the tooth row. The internal mandibular groove extends from just anterior to this foramen to a point beneath M_1 .

As in all metacheiromyids, there is a prominent medial buttress, forming anteriorly a low, narrow ridge lingual to the molars. The dorsal surface of the buttress is roughened, particularly behind M_3 . On the external surface of the mandible the masseteric fossa, though rather poorly defined, is deeper than in most other palaeanodonts. The fossa is bounded in front by a crest formed by the thickened anterior border of the coronoid. A more rounded ridge running forward from the condyle forms its lower limit. The horizontal ramus is rather slender and shallow, deeper in front than in back, as in *Palaeanodon*. A small mental foramen is present beneath P_3 .

Two palaeanodont humeri from Princeton Quarry are probably referable to *Propalaeanodon schaffi* (Princeton Quarry is about $\frac{1}{2}$ mile northwest of Jepsen Valley Quarry and at about the same stratigraphic level). They are the only other palaeanodont specimens from the same stratigraphic interval as the holotype. PU 13928, a complete left humerus (Fig. 3B), and PU 13929, a distal end of a left humerus, are of appropriate size for *Propalaeanodon*. They are smaller than the humeri of *Palaeanodon ignavus* (Fig. 3A) and *Pentapassalus woodi* (Fig. 3C and Rose, 1978) and slightly larger

TABLE 2

Dimensions (mm.) of humeri in certain palaeonodons.
(Known humeri of *Palaeonodon* are too incomplete to allow accurate measurements.)

	<i>?Propalaeonodon schaffi</i> , PU 13928	<i>Metacheiromys dasypus</i> , AM 11718	<i>Pentapassalus pearcei</i> (from Gazin, 1952)	<i>Pentapassalus woodi</i> , AC 2766
length	39.3	74.0	37.0	47.1
length of deltopectoral crest (from proximal end of greater tuberosity)	26.3	54.7	28.0	35.5
length of supinator crest (from distal end of humerus)	19.7	38.8	22.0	20.8a
greatest transverse breadth of distal end of humerus	19.8	34.0	21.7	22.6

a = approximate

than the humeri of *Pentapassalus pearcei* (Gazin, 1952). However, all these specimens are essentially similar in structure (see Table 2 for comparative dimensions).

The deltopectoral crest is broad and well developed in PU 13928. It may be relatively slightly longer than in *Palaeonodon*, but this is not certain because the known humeri of the latter are fragmentary and poorly preserved (AMNH nos. 15088, 16831, 16832). The crest is relatively shorter than in *Pentapassalus* and *Metacheiromys* (see Simpson, 1931: 340–342). Its distal end is developed into an anteromedially directed prominence for the insertion of the pectoralis major. Proximolaterally the crest gives rise to a low process for the insertion of the deltoid. A less well-defined crest extends distally from the lesser tuberosity almost to the entepicondylar foramen and bears a small teres tubercle midway along the shaft, on which the teres major inserted, as in *Metacheiromys dasypus* (Simpson, 1931).

The supinator crest is somewhat smaller than in *Pentapassalus pearcei* and *Metacheiromys dasypus* and lacks the hooklike, dorsally projecting flange characteristic of the latter two species.



Rose, K D. 1979. "A new Paleocene Palaeonodont and the origin of the Metacheiromyidae (Mammalia)." *Breviora* 455, 1–14.

View This Item Online: <https://www.biodiversitylibrary.org/item/22496>

Permalink: <https://www.biodiversitylibrary.org/partpdf/231>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.