# THE HABITAT OF ÆNIGMATIAS (DIPTERA: PHORIDÆ)

## By C. T. Brues

Biological Laboratories, Harvard University

The sexes of the interesting genus *Ænigmatias* like those of many Phoridæ are extremely dissimilar owing to the apterous condition and otherwise profoundly modified and degenerate form of the female. Three species, known from the winged male have been found in North The female of one of these is known, defi-America. nitely associated with the male from a pair taken in copula by Malloch and described by him as Ænigmatias flavofemorata. Malloch's types were from Illinois, collected "on a sandy bank along the Illinois Central Railroad between White Heath and the Sangamon River." As the genus is known definitely to be myrmecophilous, at least in Europe, the specimens taken by Malloch were quite probably from the nest of some ant in the vicinity. Previous to this a single female was taken in Arizona by Coquillett and described by him as Æ. schwarzii. Of this he says "it occurred on low vegetation in a local-

ity where no ant nests could be found."

During the summer of 1945, the present writer collected a large number of miscellaneous insects on sheets of "tanglefoot" fly-paper in northwestern Massachusetts while making a census of insects to determine the effect of DDT on the insect fauna of woodlands. Among the more than 300,000 specimens obtained (of which some 51% were Phoridæ) was a single female of Ænigmatias from Petersham, August 4. This, which is apparently the third female to be reported from North America, was trapped on the trunk of a tree several feet above the ground. From this, and from the experience of both Coquillett and Malloch it is apparent that these tiny wingless Enigmatias wander freely from the ant nests where they are domiciled, perhaps following foraging This specimen is not sufficiently well preserved to determine whether it is Æ. flavofemorata, or more probably, *E. eurynota* Brues known from a male collected in 1910 by Mrs. Brues at Forest Hills, Massachusetts.



Brues, Charles T. 1947. "The Habitat of Ænigmatias (Diptera: Phoridæ)." *Psyche* 54(4), 265–265. <a href="https://doi.org/10.1155/1947/20476">https://doi.org/10.1155/1947/20476</a>.

View This Item Online: <a href="https://www.biodiversitylibrary.org/item/206966">https://www.biodiversitylibrary.org/item/206966</a>

**DOI:** https://doi.org/10.1155/1947/20476

Permalink: <a href="https://www.biodiversitylibrary.org/partpdf/235059">https://www.biodiversitylibrary.org/partpdf/235059</a>

### **Holding Institution**

**Smithsonian Libraries and Archives** 

#### Sponsored by

**Biodiversity Heritage Library** 

### **Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <a href="https://www.biodiversitylibrary.org">https://www.biodiversitylibrary.org</a>.