

PROCEEDINGS
OF THE
NEW ENGLAND ZOÖLOGICAL CLUB

DESCRIPTION OF A NEW SUBSPECIES OF
MANACUS CANDEI (PARZUD.).

BY OUTRAM BANGS.

It has been customary to give the range of *Manacus candei*¹ as extending from southern Mexico to Costa Rica, the species having been described originally from specimens from Truxillo, Honduras.

Last winter Mr. W. W. Brown, Jr., secured a large series of specimens at Ceiba, Honduras, (spelled on some maps Laceiba) less than a hundred miles west of the type locality. Both towns are in the hot coastal plain, and the faunal conditions are undoubtedly the same at each place. Therefore Mr. Brown's specimens can be considered as being practically topotypes of true *Manacus candei* (Parzud.). On comparing these skins with a large series from Buena Vista, Vera Cruz, Mexico, a very great difference in color in the adult males is at once obvious, and the Mexican form must be distinguished by name. Besides the pronounced differences in color, the Mexican form averages slightly larger than the more southern true *M. candei*. It may be known by the following brief diagnosis.

¹ *Pipra candei* Parzud., Rev. Zool., 1841, p. 306.



Bangs, Outram. 1903. "Description of a new subspecies of *Manacus candei* (Parzud.)." *Proceedings of the New England Zoo*
..
logical Club 3, 105–106.

View This Item Online: <https://www.biodiversitylibrary.org/item/45429>

Permalink: <https://www.biodiversitylibrary.org/partpdf/240304>

Holding Institution

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Sponsored by

Harvard University, Museum of Comparative Zoology, Ernst Mayr Library

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.