

Hostplant Records and Natural History Notes on Costa Rican Butterflies (Papilionidae, Pieridae & Nymphalidae)

Philip J. DeVries

Department of Zoology, University of Texas, Austin, Texas 78712

Current address: Smithsonian Tropical Research Institute, Box 2072, Balboa, Panama

Abstract. Hostplant records for 209 species of butterflies in the families Papilionidae, Pieridae and Nymphalidae are reported from various habitats throughout Costa Rica to provide a basis for understanding fine-scale hostplant relationships of neotropical butterflies at the population level. Notes on egg, clutch-size, larval behavior, larval feeding, oviposition behavior, and hostplant microhabitat are given. New and unusual hostplant families and errors in the hostplant literature are discussed, and the possibilities of unrecognized sibling species masquerading under a single species name are indicated.

Introduction

To students of butterflies it has become evident that hostplant relationships play an increasingly important role in butterfly biology. Hostplant data have been used in developing classifications and systematics at the family, subfamily, genus, and species levels (Müller, 1886; Singer et al., 1983; DeVries et al., 1985), and in some cases the hostplant has been used as a character in naming populations (e.g. Murphy & Ehrlich, 1984). Hostplant data are of paramount importance for elucidating broad-scale evolutionary patterns of host use among systematic lineages (Brues, 1924; Ehrlich & Raven, 1965; Benson et al., 1976), for understanding butterfly ecology at the community (Gilbert, 1984) and population (Ehrlich, 1984) levels, for the dynamics of host-breadth and host shifts (Singer, 1983, 1984), and for the development of mimicry theory (Gilbert, 1983) and ecological chemistry (Brower, 1984). In short, hostplant relationships are a basis upon which we rest many of our ideas about butterflies.

Taken as a whole, it appears from surveys of literature records that hostplant relationships are well understood for the majority of the systematic lineages of butterflies (Ehrlich & Raven, 1965; P. Ackery, in prep.), particularly at higher systematic levels. However, this is not true with respect to specific butterfly faunas at the genus and species levels. Perhaps the only places on earth where such relationships are well known,

but not completely known at these levels are in the U.S.A., England, parts of northern Europe and Japan. In these areas fine-scale hostplant data are available for entire faunas which can be used to develop systematic and ecological studies with a critical degree of resolution. In all other areas of the world such data are not available, and this is especially so in the tropics. Although the tropics contain the greatest diversity of butterfly species, compared to temperate regions, hostplant relationships of butterflies are poorly known, and least known of all in the neotropics. If we are to gain an understanding of the butterflies of the neotropics comparable to our knowledge of the temperate regions, it is important to have a more complete picture of hostplant relationships than currently exists. This is a tall order considering that the neotropics contain a butterfly diversity greater than any other region on earth (DeVries, 1986). One very useful method is to develop hostplant lists for butterfly faunas of specific areas.

During a long term and continuing study of the butterfly fauna of Costa Rica I have had the opportunity to rear a number of species, as well as the good fortune to work with colleagues who, although involved in studies of their own, have provided me with hostplant records to augment my work. I here summarize the hostplant records for 209 species of Costa Rican butterflies in the families Papilionidae, Pieridae and Nymphalidae reared between 1976 and 1982 by my colleagues and myself. The purpose of this paper is to provide a starting point for eventually understanding fine-scale hostplant relationships at the population level for neotropical butterflies. Natural history information is provided for both butterflies and hostplants in coded form along with each hostplant record. In the remainder of this paper I highlight some new and unusual hostplant records, point to instances where there may be several species currently placed under one name, and draw attention to what I believe to be errors in previous literature. Additional hostplant records of Costa Rican butterflies and an analysis of the ecological patterns of hostplant use which are evident in the present paper will be discussed in a future publication (DeVries, in prep.).

Materials and Methods

Except where indicated, all hostplant records presented (Appendix I) have resulted from a larva feeding on a plant and developing into an adult. All records are from field collected eggs or larvae, except where otherwise noted. Oviposition records are indicated as such, the butterflies have been positively identified with a voucher specimen taken at the time of oviposition, or in a few cases, determined on the wing with the use of binoculars. Therefore, the vast majority of records reported here have voucher specimens in the collection of the author or the person responsible for the record.

Butterfly Nomenclature. The nomenclature follows DeVries (1986) which is modified and refined from DeVries (1983). Additional systematic references useful

for Costa Rican butterflies may be found in Ackery and Vane-Wright (1985), DeVries et al. (1985), Higgins (1981), Jenkins (1983) and Singer et al. (1983).

Plant Determinations. The hostplants in Appendix (I) are for the most part positively determined to genus (many to species), but in a few instances it was only possible to determine the plants to family due to undescribed taxa or insufficient plant material. The following people have lent their expertise in determining the hostplants: H. Bold, W. Burger, I. A. Chacon, R. Foster, L. D. Gomez, W. Haber, B. Hammil, L. Gilbert, G. Hartshorn, D. H. Janzen, S. Knapp, R. Marquis, R. Pohl, L. Poveda and F. G. Stiles. Useful references for identification of plants of Costa Rica include: Allen (1956), Burger (1971, 1977), Croat (1978), Daniels and Stiles (1979), Janzen and Leisner (1980) and Stiles (1980).

Coded Information in Appendix I: Explanations for the coded sequences in Appendix I are found in Tables I, II and III. This information is found in a series of columns located to the right of each butterfly taxon and corresponds to: hostplant identity, locality of record and who reared the butterfly, egg clutch-size, oviposition behavior, larval behavior, larval feeding and binomic notes concerned with the ovipositing butterfly and microhabit of the hostplant respectively. Further details on localities, habitats, and butterfly biology in Costa Rica are found in DeVries (1986).

All hostplant records reported here are from Costa Rica except those coded *AA and *KS. These come from Barro Colorado Island, Panama and augment Costa Rican records. Except for those hostplant records marked with an asterisk and a set of initials preceding the locality code, all records are those of DeVries. The initials refer to records of the following people: AA = A. Aiello, IC = I. A. Chacon, RC = R. Cubero, SK = S. Koptur, LEG = L. E. Gilbert, WH = W. Haber, DJ = D. H. Janzen, JM = J. Mallet, RM = R. Marquis, KS = K. Steiner, MCS = M. C. Singer, FGS = F. G. Stiles, and TR = T. Ray.

Comments

The hostplant records in Appendix I completely corroborate the pattern of relationship between hostplants and butterflies as described by Ehrlich and Raven (1965). There are, however, a few records in Appendix I which deserve special comment because they either represent hostplant families not reported in Ehrlich and Raven (1965), or they indicate a need for further study within the neotropical butterfly fauna as a whole. These records are briefly discussed here by butterfly family.

Papilionidae: The record of *Papilio cleotas* feeding on Moraceae requires confirmation since no papilionid species is known to feed on this plant family anywhere in the world. Hostplant records in the literature frequently report that *Papilio cresphontes* and *P. thoas* both utilize Rutaceae and Piperaceae as larval hostplants. Based on field experience and the data presented here I believe that such records are misidentifications of the butterfly species; the species are extremely similar in appearance. In Costa Rica *P. cresphontes* larvae feed only on Rutaceae,

and conversely, *P. thoas* larvae feed only on Piperaceae.

Pieridae: The records of *Catasticta* and *Pereute* feeding on Loranthaceae and the behavior and morphology of larvae and pupae suggest a close affinity to the extensive Old World genus *Delias*. The records also further amplify those reported in Ehrlich and Raven (1965). However, I believe the reference of *Pereute* feeding on the Lauraceae (Jørgensen, 1932) is in error (DeVries, 1982). I also believe the record of *Perrhybris lypera* feeding on Lauraceae is a plant misidentification (see Young, 1980, 1982; DeVries, 1982). The records of *Aphrissa statira* feeding on Bignoniaceae are very unusual for the Pieridae, and suggest either a chemical convergence between certain Bignoniaceae and Caesalpiniaceae or that there is more than a single species under the name *A. statira*.

Nymphalidae: The records of *Agrias amydon* on Erythroxylaceae represent an unusual hostplant family for the subfamily Charaxinae. It is further notable because of commercial interest in both the butterfly and chemistry of secondary chemicals of the hostplant genus. The record of *Zaretis itys* feeding on Piperaceae is unusual since its usual hostplants are Flacourtiaceae in the same habitat. Several interesting records are reported from the Nymphalinae. In my experience, the larvae of *Eunica monima* feed only on Burseraceae. I believe that the original record of this species feeding on Rutaceae (Dyar, 1912) represents a misidentification of the hostplant which subsequently became embedded in the literature. Muyshondt (1975) stated that Müller (1886) was in error for reporting *Diaethria marchalli* to feed on *Trema* (Ulmaceae) and said that in El Salvador he only found this species feeding upon Sapindaceae. In Costa Rica, *Trema* is the only hostplant for *D. marchalli* of which I am aware, suggesting that either the butterfly switches hostplant families in areas north of Costa Rica, or that there are two species of butterfly involved here. Data reported here suggest either that certain species of *Adelpha* are highly polyphagous, or that there are cryptic species involved under the same name (e.g. *A. boreas*, *A. celerio*, *A. heraclea*). *Adelpha melanthe*, on the other hand, changes hostplant family from east to west in Costa Rica, but all are closely allied within the Urticales. In the subfamily Satyrinae the record of *Megeuptychia antonoe* feeding on Cyclanthaceae is unusual, but Cyclanthaceae has recently been demonstrated to be a host of *Caligo* in Costa Rica as well (I. A. Chacon, pers. comm.). The record of *Euptychia insolata* feeding on Neckeraceae is the first record for any butterfly to utilize a member of the Bryophyta as a larval hostplant, although some members of the genus are known to feed on Lycopsida (Singer et al., 1983). Lastly, the records of *Pierella* and *Cissia confusa* on Poaceae, Marantaceae, Palmae and Heliconiaceae indicate a high degree of polyphagy, mostly within the same habitat.

Acknowledgments. I would first like to acknowledge a debt of gratitude to my

colleagues who were named previously, for allowing me to publish their hostplant records and who helped determine the hostplants. A special thanks to L. D. Gomez, D. H. Janzen, L. E. Gilbert and Servicios de Parques Nacionales de Costa Rica for logistical support. For babysitting caterpillars in the field during my absence and for assistance in the field I thank: I. A. Chacon, J. Denslow, A. Ehrlich, P. R. Ehrlich, L. E. Gilbert, W. Muriel Hallwachs, R. Hesterberg, D. H. Janzen, J. Longino, J. Mallet, R. Marquis, M. C. Singer, F. G. Stiles and C. Todzia. This manuscript benefited from comments by: P. R. Ackery, L. E. Gilbert, D. J. Harvey, D. H. Janzen, R. Marquis, R. Mattoni, C. D. Thomas and R. I. Vane-Wright. I dedicate this paper to the late A. Pepper. The publication of this paper was made possible from grants to the author made available from the Smithsonian Tropical Research Institute and the Dept. of Zoology, University of Texas.

Literature Cited

- ACKERY, P. R. & R. I. VANE-WRIGHT, 1984. Milkweed butterflies: their cladistics and biology. *Bull. Br. Mus. (Nat. Hist.)*.
- ALLEN, P. H., 1956. *The Rain Forests of Golfo Dulce*. University of Florida Press, Gainesville, Florida.
- BENSON, W. W., K. S. BROWN & L. E. GILBERT, 1976. Coevolution of plants and herbivores: passion flower butterflies. *Evolution* 29:659-680.
- BROWER, L. P., 1984. Chemical defences in butterflies. *Symposia of Roy. Ent. Soc. Lond.* 11:109-134.
- BRUES, C. T., 1924. The specificity of food-plants in the evolution of phytophagous insects. *Am. Nat.* 58:127-144.
- BURGER, W., 1971. Flora Costaricensis. *Fieldiana Botany* 35:1-227.
- _____, 1977. Flora Costaricensis. *Fieldiana Botany* 40:1-291.
- CROAT, T. B., 1978. *Flora of Barro Colorado Island*. Stanford Univ. Press, Stanford Calif.
- DANIELS, G. S. & F. G. STILES, 1979. The *Heliconia* taxa of Costa Rica: keys and descriptions. *Brenesia* 15: suppl.:1-150.
- DEVRIES, P. J., 1982. The case of *Perrhybris lypera* (Pieridae) and the Lauraceae: hostplant record or assumption? *J. Lep. Soc.* 36:229-230.
- _____, 1983. Checklist of butterflies. IN: *Costa Rican Natural History*. D. H. Janzen (ed.). pp. 654-678, Univ. Chicago Press, Chicago, Illinois.
- _____, 1986. *The Butterflies of Costa Rica and their Natural History*. Princeton Univ. Press, Princeton, N.J. (in press).
- DEVRIES, P. J., I. J. KITCHING & R. I. VANE-WRIGHT, 1985. The systematic position of *Antirrhina* and *Caerois*, with comments on the classification of the Nymphalidae (Lepidoptera). *Syst. Ent.* 10:11-32.
- DYAR, H. G., 1912. Descriptions of the larvae of some Lepidoptera from Mexico. *Proc. Ent. Soc. Wash.* 14:54-58.
- EHRLICH, P. R., 1984. The structure and dynamics of butterfly populations. *Symposia Roy. Ent. Soc. Lond* 11:25-40.
- EHRLICH, P. R. & P. H. RAVEN, 1965. Butterflies and plants: a study in coevolution. *Evolution* 18:586-608.
- GILBERT, L. E., 1983. Coevolution and mimicry. IN: *Coevolution*. D. Futuyma & M. Slatkin (eds.) pp. 263-281. Sinaeur Press, Mass.
- _____, 1984. The biology of butterfly communities. *Symposia Roy. Ent. Soc.*

- Lond. 11:41-54.
- HABER, W., 1978. Evolutionary ecology of tropical mimetic butterflies (Lepidoptera: Ithomiinae). Ph.D. thesis, Univ. of Minnesota.
- HARTSHORN, G. S., 1983. Plants. IN: *Costa Rican Natural History*. D. H. Janzen (ed.). pp. 118-183. Univ. Chicago Press, Chicago, Illinois.
- HIGGINS, L. G., 1981. A revision of the genus *Phyciodes* Hübner and related genera with a review of the classification of the Melitaeinae (Lepidoptera: Nymphalidae). Bull. Brit. Mus. Nat. Hist. (Ent.) 43:77-243.
- JANZEN, D. H. & R. LEISNER, 1980. Annotated check-list of plants of lowland Guanacaste Province, Costa Rica, exclusive of grasses and non-vascular cryptogams. *Brenesia* 18:15-90.
- JENKINS, D. W., 1983. Neotropical Nymphalidae I. Revision of Hamadryas. Bull. Allyn Mus. 81:1-146.
- JÖRGENSEN, P., 1932. Lepidopterologisches aus Südamerika. Deutsch. Ent. Zeitschr. Iris, Dresden. 46:37-66.
- MÜLLER, W., 1886. Südamerikanische nymphalidenraupen: versuch eines natürlichen systems der nymphaliden. Zoologische Jarbücher, Jena. 1:417-678.
- MURPHY, D. D. & P. R. EHRLICH, 1984. Biosystematics of the *Euphydryas* of the Central great basin with description of a new subspecies. J. Res. Lep. 22:254-261.
- MUYSHONDT, A., 1975. Notes on the life cycle and natural history of the butterflies of El Salvador. VI,A. *Diaethria astala* Guerin (Nymphalidae-Calli-corinae). J. New York. Ent. Soc. 83:10-18.
- SINGER, M. C., 1983. Determinants of multiple host use by a phytophagous insect population. *Evolution* 37:389-403.
- _____, 1984. Butterfly-hostplant relationships: host quality, adult choice and larval resources. *Symposia Roy. Ent. Soc. Lond.* 11:81-88.
- SINGER, M. C., P. J. DeVRIES & P. R. EHRLICH, 1983. The *Cissia confusa* species-group in Costa Rica and Trinidad (Lepidoptera: Satyrinae). Zool. J. Linn. Soc. 79:101-119.
- STILES, F. G., 1980. Further data on the genus *Heliconia* (Musaceae) in northern Costa Rica. *Brenesia* 18:147-154.
- YOUNG, A. M., 1980. Notes on the behavioral ecology of *Perrhybris lypera* (Pieridae) in northeastern Costa Rica. J. Lep. Soc. 34:36-47.
- _____, 1982. *Perrhybris lypera* (Pieridae) feeding on Lauraceae; a response to DeVries. J. Lep. Soc. 36:230-232.

Table 1. Numerical Codes for Hostplant Families in Appendix I.

1 Acanthaceae	17 Erythroxylaceae	33 Poaceae
2 Amaranthaceae	18 Euphorbiaceae	34 Poaceae:Bambusoidea
3 Annonaceae	19 Fabaceae	35 Rubiaceae
4 Aristolochiaceae	20 Flacourtiaceae	36 Rutaceae
5 Asclepidiaceae	21 Heliconiaceae	37 Sapindaceae
6 Bignoniaceae	22 Hernandiaceae	38 Selaginellaceae
7 Brassicaceae	23 Lauraceae	39 Simaroubaceae
8 Bromeliaceae	24 Loranthaceae	40 Solanaceae
9 Burseraceae	25 Marantaceae	41 Tiliaceae
10 Caesalpiniaceae	26 Melastomataceae	42 Tropaeolaceae
11 Capparidaceae	27 Mimosaceae	43 Turneraceae
12 Caricaceae	28 Moraceae	44 Ulmaceae
13 Compositae	29 Neckeraceae	45 Umbelliferae (= Apiaceae)
14 Cyclanthaceae	30 Palmae (= Arecaceae)	46 Urticaceae
15 Cyperaceae	31 Passifloraceae	47 Verbenaceae
16 Ericaceae	32 Piperaceae	

Table 2. Codes for Rearing Localities Cited in Appendix I.

AT = Atenas (Al)	LS = Finca la Selva (HER)
BLH = Bajo la Hondura (SJ)	MV = Monte Verde (PUN)
BAR = Barranca (PUN)	PATT = Patarra (SJ)
BCI = Barro Colorado Island (PAN)	RCN = Rincón de Osa (PUN)
CDM = Cerro de la Muerte (SJ)	SV = San Vito de Java (PUN)
CURR = Curridibat (CAR)	SRNP = Parque Santa Rosa (GUAN)
COP = Copey (SJ)	SJ = San José (SJ)
CAN = Cañas (GUAN)	TUR = Turrialba (CAR)
CAR = Parque Braulio Carrillo (SJ)	UCR = Universidad de Costa Rica (SJ)
CVDO = Parque Corcovado (PUN)	VTUR = Volcán Turrialba (CAR)
EST = Estrella de Cartago (CAR)	VBVA = Volcán Barva (HER)
ELROD = Finca el Rodeo (SJ)	VSM = Volcán Santa María (GUAN)
GOL = Golfito (PUN)	VDS = Virgen del Socorro (AL)
HER = Heredia (HER)	VM = Volcán Miravalles (AL)
LALT = Finca las Alturas (PUN)	

Costa Rican Province Codes are: (AL) = Alajuela, (CAR) = Cartago, (GUAN) = Guanacaste, (HER) = Heredia, (PUN) = Puntarenas, (SJ) = San José. The code (PAN) = the country Panamá.

Table 3. Codes Used for Eggs, Larvae and Hostplant in Appendix I.

Eggs and Oviposition

S = eggs laid singly

C = eggs laid in clusters

1 = female may oviposit several times on same plant.

2 = female usually oviposits only once on a plant.

3 = female always oviposits several times on same plant.

Larval Behavior

S = solitary

G = gregarious

1 = larvae feed on new, then old leaves.

2 = larvae will feed on all leaves.

3 = larvae feed only on young leaves.

4 = larvae feed only on older leaves.

5 = larvae feed on plant parts other than leaves.

Hostplant Microhabitat

1 = female oviposits on plants occurring in open areas.

2 = female oviposits on plants occurring in forest.

3 = female oviposits on plants occurring at forest or riparian edges.

4 = female oviposits on plants occurring in forest light gaps.

Appendix I. Reading from left to right the information in this appendix is divided into five columns corresponding to: 1) butterfly taxa, 2) hostplant, locality and authorships of rearing, 3) clutch-size of eggs and frequency of oviposition, 4) larval behavior and feeding and 5) micro-habitat of hostplant and natural history notes. Explanations for the coded sequences are found in the text and in Tables 1, 2 and 3.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
---------	----------------------	-------------------	--------------------------------------

FAMILY: PAPILIONIDAE

GENUS: <i>Battus</i>			
<i>polydamas</i> L.	<i>Aristolochia</i> (4) <i>grandiflora</i> SJ <i>anguicida</i> SRNP undetermined LS	C G1 C G1	All hostplants are 3 and 4. Egg clusters range from 5 to 10.
<i>belus</i>	<i>Aristolochia</i> (4) <i>grandiflora</i> SJ, TUR. undetermined LS. undetermined CVDO undetermined SV	S1 S1 S1 S1 S1 S1	Hostplants are 3, 4 and riparian edges.
<i>varus</i> Kol.			
<i>crassus</i> Cr.	<i>Aristolochia</i> (4) <i>veraguensis</i> CVDO undetermined LS	C G1 C G1	Hostplants are 2, 3, 4 and riparian edges. Egg clusters range from 20 to 40 eggs.
GENUS: <i>Parides</i>			
<i>photinus</i> Dbdly.	<i>Aristolochia</i> (4) <i>pilosa</i> ELROD, AT.	C G2	Hostplants are 3 and 4 provided they receive direct sunshine. Plants are usually juvenile growing along the ground.
<i>childrenae</i> Gray	<i>Aristolochia</i> (4) <i>torduzii</i> LS. undetermined *RM LS	S1 S1	Hostplants are all 2, either young or old, provided they have vigorous new growth.
<i>sesostris</i>			
<i>zestos</i> Gray	<i>Aristolochia</i> (4) undetermined VM	?	This rearing was of six mature larvae all on the same juvenile plant, eating all leaves.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
GENUS: Papilio			
<i>cresphontes</i> Cr.	<i>Citrus</i> (36) several species SV, AT, GOL, PAT.	S1 S1	All <i>Citrus</i> = 1, <i>Zanthoxylum</i> = 2, 3 on medium sized trees in the forest. Janzen's record is from a late instar larva.
	<i>Zanthoxylum</i> (36)	S1 S1	
	<i>setulosum</i> SRNP, CAN, CVDO		
	<i>Esenbeckia</i> (36)		
	<i>litoralis</i> *DJ (79-SRNP-106)	S1 S1	
<i>thoas</i>	<i>Piper</i> (32) <i>tuberculatum</i> SRNP, AT		All plants are either 1 or 4 depending on the species and the habitat. All eggs are S2; all larvae are S1.
	<i>sancti-felicitis</i> LS		
	<i>marginatum</i> *DJ (81-SRNP-1046)		
	<i>amalago</i> *DJ (81-SRNP-827)		
	<i>auritum</i> LS		
	<i>multiplinervium</i> *RM LS		
	<i>reticulatum</i> *JM CVDO		
	<i>friedrichsthalii</i> *JM CVDO		
	undetermined VDS		
<i>polxenes</i>	<i>Apium</i> (45)		
<i>stabilis</i> R & J	<i>leptophyllum</i> SJ,VDS,VB, PAT,COP, TUR.	S1 S2	Both hostplants are 1, females oviposit on young and mature plants.
	<i>Foeniculum</i> (45)		
	<i>vulgare</i> MV.	S1 S2	
<i>cleotas</i>	<i>Sorocea</i> (28)	?	
<i>archytas</i> Hop.	<i>trophoides</i> *WH VM	?	Haber reared this species from a solitary mature larva feeding on mature leaves. This represents a very unusual record for the Papilionidae.
<i>victorinus</i>	<i>Persea</i> (23)	?	
<i>vulneratus</i> Butler	<i>americana</i> *WH MV	?	Haber reared this species from a solitary late

instar larva feeding on all leaves of an open grown tree.

<i>Hernandia</i> (22)	S2	All of the hostplants are 4, but plants in light gaps are in deep shade. The females oviposit on saplings about 1 meter tall.
<i>didymantha</i> LS	S3	
undetermined CAR	?2	
undetermined *RM LS	S?	
<i>Cassimiroa</i> (36)	C	<i>Cassimiroa</i> and <i>Citrus</i> are 1 and from previous experience in Santa Rosa I assume that <i>Zanthoxylum</i> is either 1 or 4. Females oviposit on young or mature plants.
<i>edulis</i> SJ	G1	
<i>Citrus</i> (36)	C	
various species CAR,CVDO,TUR,LS, PAT	G1	
<i>Zanthoxylum</i> (36)	C?	
<i>setulosum</i> *DJ (79-SRNP-163)	G1	

FAMILY: PIERIDAE

GENUS: *Dismorphia*

<i>amphionia</i>				
<i>praxinoe</i> Dbldy.				
	S1	S2	Hostplants are all 2, 3, or 4 in shady areas.	
	S1	S2	Females oviposit on saplings or medium sized	
	S1	S2	trees in the understory. Larvae usually eat old	
	S1	S2	leaves but all are acceptable.	
<i>Inga</i> (27)				
<i>undetermined</i> CAR	S1	S?	I have observed oviposition but have not reared	
<i>undetermined</i> LS	S1	S?	larvae. The female oviposited four eggs on the	
<i>sapindoides</i> *SK TUR	S1	S?	same sapling in shady understory of cloud	
<i>densiflora</i> *SK TUR	S1	S?	forest habitat.	
<i>Inga</i> (27)				
<i>undetermined</i> VB	S1	S3	Hostplants are 2, 3 and 4 depending on the	
<i>eunoae</i>			habitat. All plants are saplings.	
<i>desine</i> Hew.				
<i>Inga</i> (27)				
<i>zathoe</i>				
<i>pallidula</i> But. & Dru.	S1	S3	Hostplants are 2, 3 and 4 depending on the	
	S1	S3	habitat.	
	S1	S3		

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>crisia</i> <i>lubina</i> But.	<i>Inga</i> (27) undetermined EST undetermined *SK MV	S1 S1 S1	All hostplants are 2, 3 and 4 and range from understory saplings to mature canopy trees. Females will lay from 5 to 15 eggs on the same plant depending on the size of the plant.
<i>zaela</i> <i>oreas</i> Sal.	<i>Inga</i> (27) undetermined CAR	S1 S2	Hostplants are 3 and 4 and all are saplings in the understory or in shady light gaps.
GENUS: Lieinx			
<i>nemesis</i> G & S	<i>Inga</i> (27) <i>mortonia</i> *SK MV	S1 S1 S1 S1 S1	Hostplants are 3 and 4. This species can apparently feed on a variety of species of <i>Inga</i> but is restricted entirely to new growth leaves.
<i>longispica</i> *SK MV		S3	
<i>brenesii</i> *SK MV		S3	
<i>densiflora</i> *SK MV		S3	
GENUS: Anteos			
<i>clorinde</i> Godart	<i>Cassia</i> (10) <i>emarginata</i> SRNP	S1 S1	Hostplants are 1 and may be young or mature.
<i>mareula</i> Fab.	<i>Cassia</i> (10) <i>emarginata</i> SRNP	S1	Hostplants are 1 and may be young or mature.
GENUS: Phoebis			
<i>argante</i> Fab.	<i>Cassia</i> (10) <i>biflora</i> SRNP <i>fruticosa</i> LS	S1 S1 S1	All hostplants are 1 except <i>Pentaclethra</i> which may be 4. All range from saplings to mature trees and may be seedlings in <i>Pentaclethra</i> .
	<i>Pentaclethra</i> (27)	S1	
	<i>macroloba</i> LS	S1	

<i>philea</i> L.	<i>Inga</i> (27) <i>vera</i> *DJ (79-SRNP-387)	?	S
	<i>ruiziana</i> *JM CVDO	?	S
	<i>Cassia</i> (10) <i>grandis</i> SJ	S1	S1
	<i>alata</i> SJ	S1	All plants are 1 or 3 and range in size from young to mature forest trees.
	<i>leptocarpa</i> *DJ (79-SRNP-398)	?	S?
	<i>hayesiana</i> *DJ (82-SRNP-4)	?	S?
	<i>fruticosa</i> *RM LS	?	S?
	<i>Cassia</i> (10) <i>fruticosa</i> *RM LS	S1	Marquis records this species from numerous larvae feeding only on new leaves of an isolated tree in an open area.
	<i>sennae</i> Cr.	?	S?
	<i>Cassia</i> (10) <i>biflora</i> SRNP <i>obtusifolia</i> *DJ (79-SRNP 407)	?	All hostplants are 1 or 3 and usually sapling trees.
GENUS: <i>Aphrissa</i>			
<i>statira</i> Cr.	<i>Callichlamys</i> (6) <i>latifolia</i> *DJ (81-SRNP-72)	?	Janzen reared this species from solitary late instar larvae only on new leaves. These records are the first for the Pieridae feeding on the Bignoniaceae.
<i>albula</i> Cr.	“Woody vine” (6) undetermined *DJ (79-SRNP-10918)	?	S3
GENUS: <i>Eurema</i>			
<i>mexicana</i> Feld.	<i>Diphysa</i> (19) <i>robinoides</i> HER	S1	Hostplant is 1 or 3 in bright sunshine. Plants range from saplings to mature trees.
	<i>Cassia</i> (10) <i>fruticosa</i> LS	S1	Hostplant is 1 and ranges from saplings to mature trees. Mature larvae are able to feed on old leaves.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>dina</i> West.	<i>Picramnia</i> (39) <i>alleni</i> ELROD <i>quaternaria</i> SRNP	S1 S3 S1 S3	Hostplants are 2, 3 or 4 provided there is some shade. Plants range from saplings to mature trees.
<i>daira</i> Feld.	<i>Aeschnophlebia</i> (19) undetermined SRNP, CAN.	S1	Hostplant 1, usually in recently disturbed pastures and roadsides.
GENUS: <i>Catasticta</i>			
<i>teutila</i> Dbdly.	<i>Dendrophthora</i> (24) <i>costaricensis</i> VTUR, CDM	C	The hostplant is an epiphytic parasite. There has been confusion in the literature, because the host tree rather than the parasite was often taken to be the hostplant of the butterfly. Females oviposit on mature hostplants occurring on mature emergent canopy trees or on large remnant trees in pastures.
<i>theresa</i> Butler	<i>Antidaphne</i> (24) <i>viscidoides</i> COP	C	The hostplant is an epiphytic parasite of <i>Alnus</i> (Betulaceae) that occurs along riparian edges. Egg clusters range from 10 to 50 eggs.
<i>nimbice</i> bryson G & S	<i>Struthanthus</i> (24) undetermined *IC PAT	?	Chacon reared this species from a mass of early instar larvae feeding on all leaves. The hostplant is an epiphytic parasite of forest trees.
GENUS: <i>Pereute</i>			
<i>cheops</i> Staud.	<i>Antidaphne</i> (24) <i>viscidoides</i> COP	C	This is a single cluster of eggs after I observed oviposition. Larvae have not been reared past first instar.

<i>charops</i> Boisd.	C	I have repeatedly watched, with binoculars, several females oviposit large clusters of eggs on what was certainly Loranthaceae. In addition I have seen large groups of larvae feeding on leaves of the same epiphytic parasite. These larvae descended the tree trunk and pupated on the tree trunk. The hostplants were growing on large emergent canopy trees along riparian edges.
GENUS: Hesperocharis		
<i>crocea</i> Bates	?	Chacon reared this species from a small aggregation of late instar larvae feeding on all leaves. The hostplant is an epiphytic parasite.
Struthanthus (24)	?	
undetermined *IC SJ	S1	Hostplants are 3 and 4 and are mature woody shrubs.
GENUS: Itaballia		
<i>demophile</i> J & T	?	
<i>Caparis</i> (11) <i>indica</i> SRNP, BAR <i>frondosa</i> *WH CAN	S1	Hostplants are 3 and 4 and are mature woody shrubs.
<i>Podandrogyne</i> (11) <i>pulcherrima</i> BLH, CAR, COP, SV	S1	Hostplant is 2 or 3 usually along riparian edges. Females oviposit on young and mature plants. Larvae also feed on flower buds.
<i>caesia</i> <i>tenuicornis</i> B & D		
GENUS: Pieriballia		
<i>mandella</i> <i>noctipennis</i> B & D	?	Haber reared this species from a solitary late instar larva feeding on all leaves. The plant occurred as a small understory tree.
<i>Caparis</i> (11) <i>pseudocacao</i> *WH MV		

Egg & Larval Data

Hostplant & Locality

Species

GENUS: Perrhybris

pyrrha Fab.

Capparis (11)
isthmensis CVDO

C C G2 Hostplant is 2 and 4, either young or mature tree.

lypera Kollar

Capparis (11)
pittieri LS
pittieri *WH LS
undetermined *LEG LS

C C G2 Hostplants are 2, 3 or 4, ranging from seedling to sapling plants. Egg clusters range from 10 to 50 per leaf.

GENUS: Ascia

monuste L.

Lepidium (7)
undetermined SJ
Crataeva (11)
undetermined *LEG CVDO

S1 S1 Hostplant is always 1 and in areas of recent disturbance. Both young and mature plants are acceptable to ovipositing females. The *Crataeva* record is an oviposition as viewed through a Questar telescope.

josepha G & S

Capparis (11)
indica SRNP, CAN
incana *DJ (81-SRNP-1086)

S1 S1 Hostplants are 3 and 4 and may be sapling or mature trees.

GENUS: Leptophobia

aripa Boisd.

Nasturtium (7)
officinale SV, EST, CAR.
Tropaeolum (42)
maritizianum VDS, CAR

C C G2 Both hostplants are riparian edge species and usually grow in bright sunshine. Plants may be young or mature.

FAMILY: NYMPHALIDAE
SUBFAMILY: CHARAXINAE

Bionomic Notes of Host and Butterfly

GENUS: *Archaeoprepona*

<i>demophon</i> L.	<i>Nectandra</i> or <i>Ocotea</i> (23) undetermined LS	S1	S1	Hostplants are 3 or 4 and range from young to mature trees. Larvae rest on frass chains.
	<i>Ocotea</i> (23) <i>veraguensis</i> *DJ (81-SRNP-218)	?	S1	
	<i>Persea</i> (23) <i>americana</i> VB	S1	S1	Hostplants may be 1, 2, 3 or 4 depending on the habitat and range from seedlings to mature forest trees. Larvae rest on frass chains.
	<i>Phoebe</i> (23) undetermined *WH MV	?	S?	
			?	Records are from late instar larvae which fed on all leaves of a forest edge tree.

GENUS: *Prepona*

<i>laertes</i>	<i>Inga</i> (27)	S2	S2	
<i>octavia</i> Fruh.	<i>ruiziana</i> *JM CVDO	?	?	

GENUS: *Agrias*

<i>amydon</i>	<i>Erythroxylum</i> (17)	S2	S2	
<i>philatelica</i> DeVr.	<i>hawanense</i> * DJ (80-SRNP-209)	?	?	Both records are from solitary late instar larvae feeding on all leaves. The hostplant is an understory shrub along edges and in gaps.
	undetermined *TR LS	?	?	

GENUS: *Siderone*

<i>marthesia</i> Cr.	<i>Casearia</i> (20)	S2	S2	
	<i>sylvestris</i> *DJ (81-SRNP-339)	?	?	The hostplants are 2, 3 or 4 and range from saplings to mature trees. <i>Zuelania</i> records only on saplings.
	<i>Zuelania</i> (20)	?	?	
	<i>quidonia</i> *DJ (81-SRNP-1045)	?	?	

GENUS: *Zaretis*

<i>itys</i> Cr.	<i>Ryania</i> (20)	S1	S1	All hostplants are 2, 3 and 4 and range from saplings to mature forest trees, except Piperaceae, which are herbaceous. This species
	<i>speciosa</i> LS			
	<i>Laetia</i> (20)			
	<i>procera</i> LS	S1	S1	

Egg & Larval Data

Bionomic Notes of Host and Butterfly

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>Cassearia</i> (20)			
<i>nitida</i> LS	S1	S1	feeds on a wide variety of hosts at La Selva, which also may be found to be the case in the Pacific drainage. Of interest is that Atlantic and Pacific drainage adults look very different; a detailed comparison of the early stages should provide an insight to whether or not the name <i>elllops</i> should be applied to the Pacific drainage insects.
<i>arborea</i> LS	S1	S1	
<i>corymbosa</i> *DJ (81-SRNP-272)	?	S1	
<i>arguta</i> *DJ (81-SRNP-232)	?	S1	
undetermined *JM CVDO	?	S	
<i>Piper</i> (32)	?	S1	
<i>arieianum</i> *RM LS			
 GENUS: <i>Consul</i>			
<i>fabius</i> Dbdly.			
<i>Piper</i> (32)			
<i>sancti-felicis</i>	S1	S2	Hostplants are 2, 3 and 4. Early instars rest on frass chains, later instars rest in a tube.
<i>tuberculatum</i>	S1	S2	
undescribed species	?	S2	
<i>auritum</i> *JM CVDO	?	S2	
<i>amalago</i> *DJ (81-SRNP-600)	?	S2	
<i>jansoni</i> Salvin			
<i>Piper</i> (32)			
<i>reticulatum</i> *JM CVDO	?	S2	All records of this species are from solitary late instar larvae from hostplants growing along forest or riparian edges.
undetermined CAR	?	S2	
<i>reticulatum</i> *RM LS	?	S2	
 GENUS: <i>Hypna</i>			
<i>clytemnestra</i> Cr.			
<i>Croton</i> (18)			
<i>schiedeanus</i> *RM LS	?	S2	Marquis reports this from a solitary late instar larva feeding on a sapling plant. The larva rests on frass chains.
 GENUS: <i>Memphis</i>			
<i>eurypyre</i>			
<i>confusa</i> Hall			
<i>Croton</i> (18)			
<i>jalapensis</i> SJ, ELROD	S1	S1	Hostplant is 3 or riparian edges, ranging from

saplings to mature trees.

Croton (18)
jalapensis ELROD, AT

S1 S1 Hostplant is 3, ranging from saplings to mature trees. Early instars rest on frass chains, later roll a tube.

Accalypha (18)
garnieri SRNP

S1 S1 Hostplant is 3 or 4, ranging from saplings to mature shrubs. Early instars make frass chains, later they roll a tube.

Croton (18)
undetermined SJ, VB
undetermined AT
schiedeanus LS

S1 S2 Hostplants are 2, 3 or 4 and range from saplings to mature forest trees. Early instars make frass chains, later they roll a tube.

Croton (18)
jalapensis AT, SJ
schiedeanus *JM CVDO

S1 ?
? S2 Hostplants are 1, 2, 3 or 4 and range from saplings to mature forest trees. Early instars make frass chains, later they roll a tube.

Piper (32)
undetermined EST, CAR

? S2 This species is recorded from solitary late instar larvae feeding on all leaves. The hostplant is a semi-woody vine growing at the bases of large trees in the forest.

Croton (18)
schiedeanus LS
schiedeanus *JM CVDO

S1 ?
? S2 Hostplants are 3 or 4 but most commonly along riparian edges. Early instars make frass chains, later they make tubes.

Croton (18)
bilbergianus *AA (79-70, 79-101)

S S3 Aiello reared these from Barro Colorado, Panama. There the hostplant occurs as a small tree in light gaps.

Piper (32)
undescribed species LS
xanthostachyum *RM LS

? S2 Hostplants are 2 or 4; one is an epiphytic vine, the other terrestrial. Early instars make frass chains, later they roll tubes.

glycerium Dbldy.

ateda Guer.-Men.

arginussa
eubaena Boisd.

pithysa Felder

beatrix Druce

artacaena Hew.

oenomaia Boisd.

cleomestra Hew.

Egg & Larval Data

Bionomic Notes of Host and Butterfly

Species	Hostplant & Locality	Data	Bionomic Notes of Host and Butterfly
<i>morvus</i> <i>bouisduwalli</i> Comst.	<i>Persea</i> (23) undetermined *IC LS	?	S2 Chacon records this species from a solitary late instar larva. The host was a mature tree along forest edge. Mature larvae roll tubes.
<i>forneri</i> G & S	<i>Ocotea</i> (23) <i>veraguensis</i> *DJ (81-SRP-939)	?	S1 Hostplant is 2, 3 or 4. Early instars make frass chains, last instars roll tubes.
FAMILY: NYMPHALIDAE			
SUBFAMILY: NYMPHALINAE			
GENUS: Marpesia			
<i>petreus</i> Bates	<i>Ficus</i> (28) undetermined SRNP <i>goldmani</i> *DJ (81-SRNP-158a)	S1 ?	S1 Hostplants are 1, 2, 3 and 4 and usually large trees. S1
<i>chiron</i> Fab.	<i>Brosimum</i> (28) <i>alicastrum</i> *DJ (80-SRNP-113) <i>lactescens</i> *RM LS	?	S2 Both records are from late instar larvae on mature forest trees. S2
<i>merops</i> Boisd.	<i>Brosimum</i> (28) <i>lactescens</i> LS	?	?
GENUS: Colobura			
<i>dirce</i> L.	<i>Cecropia</i> (28) various species: SRNP, MV, SV, TUR, SJ, LALT, VSM, CVDO, AT		All of my records have eggs laid in loose clusters of 10 to 30 eggs on saplings to mature trees. These have resulted in small clusters of

larvae that are semi-gregarious. J. Mallet reared "dirce" from La Selva and found gregarious masses of larvae composed of more than 30 individuals from emergent trees. There are striking differences in larval coloration suggesting that perhaps there are two species (*dirce* and *dirceoides*) as indicated in Sepp (1828).

GENUS: *Historis**odius* Fab.

Cecropia (28)
peltata SRNP, AT, VSM
undetermined LS, CAR, TUR

S1 S2 Hostplants are usually 3 or 4 depending on the habitat, ranging in size from saplings to mature trees. Early instars make frass chains, later they rest on the apical meristem.

acheronta Fab.

Cecropia (28)
peltata SRNP, CVDO

S1 S1 Hostplants are usually saplings along edges. Larval behavior as for *odius*.

GENUS: *Smyrna**blomfieldia*
datis Fruh.

Ureia (46)
baccifera SRNP

? S2 I have reared this from a solitary late instar larva. In SRNP the hostplant occurs as a riparian plant.

GENUS: *Pycina**zamba* zelys G & S

Ureia (46)
undetermined MV
undetermined CAR

S1 ? These records are from a number of oviposition records from mature plants along forest edge. I have not reared the larvae.

GENUS: *Tigridia**acea* L.

Cecropia (28)
undetermined LS, CAR

Females oviposit singly but tend to lay in loose

Bionomic Notes of Host and Butterfly**Data****Hostplant & Locality****Species**

undetermined *JM
Pauromia (28)
aspera LS
 undetermined CAR

GENUS: Hamadryas

februa Hubner

Dalechampia (18)
scandens SRNP, AT

S1

clusters of 3 to 8 eggs on the same leaf. Larvae are solitary or semi-gregarious and feed only on young leaves. Hostplants are always saplings and seedlings, and occur as 2, 3 or 4.

amphinome
mexicana Lucas

Dalechampia (18)
scandens SRNP

C

Hostplant is 3 or 4; eggs in long chains.

guatemalena Bates

Dalechampia (18)
scandens SRNP

S1

Hostplant is 3 or 4 or vines ascending into the forest canopy.

laodamia
saurites Fruh.

Dalechampia (18)
 undescribed species *JM LS

?

Mallet reports this species from a solitary late instar larva. Hostplant was along forest edge ascending into the canopy.

iptheme Bates

Dalechampia (18)
cissifolia *AA (78-66, 78-79)

S

Aiello reports this from Panama where the hostplant occurs in open areas and along forest edges.

GENUS: Ectima

rectifascia But. & Dru.

Dalechampia (18)
 undescribed species LS

S1

Hostplants are 3 and 4 and are all juvenile plants.

GENUS: Eunica

monima

Bursera (9)

<i>modesta</i> Bates	S1	S3	Hostplant can be 1, 2, 3 or 4 and during outbreak years the larvae can defoliate entire trees.
<i>mira</i> G & S			
<i>Mabea</i> (18)		S5	Steiner reports that in Panama larvae are only found on male portions of the inflorescence and not on leaves; plants are riparian. At La Selva I found that eggs are laid on fruiting trees and that first instars will feed on very new leaves; the plants occur in deep forest.
<i>occidentalis</i> *KS-BCI		?	
<i>occidentalis</i> LS	S1	S3-5	
<i>Mabea</i> (18)		?	Steiner reports this species as in <i>mira</i> .
<i>occidentalis</i> *KS-BCI			
<i>mygdonia</i> Godart			
<i>Mabea</i> (18)		S5	
<i>occidentalis</i> *KS-BCI		?	
GENUS: Dynamine			
<i>salpensa</i> Bates			
<i>Dalechampia</i> (18)		S1	
undescribed species LS		S5	Hostplant is 3 or 4. Females oviposit only on developing flowers, larvae feed only on flower bracts and developing ovaries. Larvae will not eat leaves.
<i>mylitta</i> Cr.			
<i>Dalechampia</i> (18)		S1	
<i>scandens</i> SRNP, VSM		S3-5	Hostplant is 3 or 4; larvae feed only on new leaves or flower parts.
<i>glauce</i> Bates			
<i>Dalechampia</i> (18)		S1	
<i>cissifolia</i> AT		S3-5	Hostplant is 3 or riparian edge; larvae only feed on new leaves or flower parts.
<i>hopi</i>			
<i>albicola</i> Rober		S1	?
<i>laothoe</i>			Mallet records this as an oviposition record on flower buds.
<i>libera</i> Fab.			
GENUS: Temenis			
<i>Seriania</i> (37)			
<i>paucidentata</i> CVDO	S1	S2	Hostplants are 2, 3 and 4 and range from juvenile to mature plants. Females oviposit
undetermined LS	S1	S2	on damaged portions of leaves. Early instars
<i>atrolineata</i> *DHJ (81-SRNP-1092)	?	S2	make frass chains, later rest on leaf edges.
<i>Paulinlia</i> (37)			
undetermined *RM LS			

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
GENUS: Epiphile			
<i>adrasta</i> Hew.	<i>Serjania</i> (37) undetermined MV <i>Cardiospermum</i> (35) undetermined SJ	S1 S2	Hostplants are 2 or 4 or as canopy vines and range from juvenile to mature plants. Early instars make frass chains, later instars rest on upperside of leaves.
<i>oreas</i> <i>plutios</i> G & S	<i>Serjania</i> (37) undetermined CAR	S1 S2	Hostplants are 2, 3 or 4 or canopy vines. Early instars make frass chains; later instars rest on frass chains or upperside of leaves.
GENUS: Catonephele			
<i>numilia</i> esite Feld.	<i>Alchornea</i> (18) <i>costariicensis</i> LS	S1 S2	Hostplant is 3 and 4. Females oviposit on sapling sized plants. Early instars make frass chains, later instars rest on the uppersides of leaves.
<i>orites</i> Stichel	<i>Alchornea</i> (18) <i>costariensis</i> LS	S1 S2	Hostplants are 2, 3 and 4. Females oviposit on seedlings and saplings, usually within the seed shadow of a mature tree. Early instars make frass chains, later instars rest on upperside of leaves.
<i>chromis</i> <i>godmani</i> Stichel	<i>Alchornea</i> (18) <i>poasana</i> EST	S1 S?	This is an oviposition record where the female laid single eggs on sapling sized plants along forest edge. Larvae were not reared to adulthood.
<i>nyctimus</i> Wstwd.	<i>Veconcibea</i> (18) <i>pleistemonia</i> *IC LS	? S2	Chacon records this from a solitary larva on a sapling sized tree along forest edge.

GENUS: *Nessaea*

- aglaura* Dbdly. & Hew.
Alchornea (18)
costaricensis LS
Plukenetia (18)
volubilis *RM LS

leaves.

GENUS: *Diaeathria*

- euepepla* G & S
Serjania (37)
 undetermined LA LT

marchalli Guer.-Men.

- Trema* (44)
micrantha LS, CAR, VSM, SV

GENUS: *Callicore*

- pitheas* Latr.

- Serjania* (37)
mexicana *AA (78-92) BCI

- lyca*
aerias G & S

Hostplants are 2, 3 and 4; all seedlings. Early instars make frass chains, later instars rest on the upperside of leaves. Marquis records this species from a solitary late-instar larva feeding on all leaves.

Hostplants are 2, 3 and 4, and range from juveniles to canopy vines. Early instars make frass chains, later rest on leaves.

Hostplants are 2, 3, 4 and range from saplings to mature trees. Early instars make frass chains, later they rest on the underside of leaves.

GENUS: *Pyrrophora*

- edocla* Dbdly. & Hew.

- Serjania* (37)
 undetermined SV

crameri Aur.

- Paulinia* (37)
 undetermined CVDO

Hostplants are 2, 3 and 4; all seedlings. Early instars make frass chains, later instars rest on the upperside of leaves. Marquis records this species from a solitary late-instar larva feeding on all leaves.

Hostplants are 2, 3 and 4, and range from juveniles to canopy vines. Early instars make frass chains, later rest on leaves.

Hostplants are 2, 3, 4 and range from saplings to mature trees. Early instars make frass chains, later they rest on the underside of leaves.

Aiello states that early instar larvae skeletonize the young leaves, later instars are on all leaves. The hostplant is 3 and 4.

Hostplants are 3 and 4, usually canopy or sub-canopy vines.

I reared this species from late-instar larvae on large woody vines occurring along forest and riparian edges. Larvae make frass chains.

Hostplant is 2, 3 and 4. Females oviposit only on newly emerging leaves. Larvae make frass chains.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
GENUS: Adelpha			
<i>melanthe</i> Bates	<i>Trema</i> (44) <i>micrantha</i> SRNP, VSM <i>Ureia</i> (46) undetermined LS undetermined CAR, VDS	S1 S1	S2-4 All hostplants are 1, 2, 3 or 4 depending on the habitat and range from young to mature plants. Early instars make frass chains, later instars rest on leaves. Although all leaves are acceptable to larvae, they are usually found on old leaves.
	<i>Myrocarpa</i> (46) <i>longipes</i> LS <i>Cecropia</i> (28) undetermined MV, SV		
<i>celerio</i> Butler	<i>Ureia</i> (46) undetermined LS <i>Miconia</i> (26) <i>argentea</i> *JM CVDO	S1 S1 ?	S2 Ureia is 3 and 4 in swampy areas that receive bright sunshine. Early instars make frass chains, later instars rest on undersides of leaves.
<i>iphicla</i>	<i>Calicophyllum</i> (35) <i>candidissimum</i> *DJ (79-SRNP-135)	?	
<i>iphicoleola</i> Bates		S2	Janzen records this species from several late instar larvae feeding on all leaves. The host-plant is 3 and 4.
<i>bassiloides</i> Bates	<i>Ixora</i> (35) <i>nicaraguensis</i> *RM LS <i>nicaraguensis</i> *JM CVDO <i>Alibertia</i> (35) <i>edulis</i> *DJ SRNP “unknown genus” (35)	?	S All records are from late instar larvae feeding on all leaves. Hostplants are found as 3 and 4.
<i>tracta</i> Butler	undetermined *WH MV undetermined *IC PAT	?	S2 Both records are from late instar larvae feeding on all leaves of an unidentified woody shrub along forest edges.
<i>boeotia</i>	<i>Luehea</i> (41) <i>seemannii</i> *JM CVDO	?	?
<i>boeotia</i> Felder		?	No information as to larval feeding. The host

commonly occurs as all categories in Corcovado but no information on where early stages are likely to occur.

<i>heraclea</i> Felder	<i>Vitex</i> (47) <i>cooperi</i> LS	S1	<i>Vitex</i> is in my experience the common host at La Selva and is 3 and 4, usually from saplings to mature trees. <i>Piper</i> record is from a late instar larva feeding on all leaves in light gap. Larvae rest on frass chains. These records represent quite a diverse set of hostplants from the same locality.
	<i>Piper</i> (32) <i>arieianum</i> *RM LS	?	S1
		S1	I have not succeeded in rearing this species to adulthood. Early instars on <i>Chomelia</i> feed on new leaves and make frass chains. <i>Chomelia</i> occurs as 3 and 4. Marquis records this species from a solitary late instar larva; <i>Satyria</i> is an epiphyte. This is an unusual record for the family Nymphalidae.
		S1	S? I have not succeeded in rearing this species to adulthood. Early instars on <i>Chomelia</i> feed on new leaves and make frass chains. <i>Chomelia</i> occurs as 3 and 4. Marquis records this species from a solitary late instar larva; <i>Satyria</i> is an epiphyte. This is an unusual record for the family Nymphalidae.
	<i>Chomelia</i> (35) <i>bispinosa</i> VDS	S1	
	<i>Satyria</i> (16) undetermined *RM LS	?	S?
		S1	Hostplants are along roadsides, riparian edges in bright sunshine. Early instars make frass chains, later instars rest on leaves.
	<i>Sabicea</i> (35) <i>aspera</i> CAR	S2	Hostplants are 2, 3 and 4 depending on the habitat and range from seedlings to saplings. Early instars make frass chains, later they rest on underside of leaves.
	"Unknown genus" (35) undetermined SV, CAR	S2	
	<i>zalmona</i> <i>sophax</i> G & S	S4	<i>Pentagonia</i> is 2, 3, 4 and is either seedling or sapling sized; <i>Psychotria</i> is sapling-2. Females oviposit on damaged portions of old leaves.
	<i>leucophaelma</i> <i>mephestophelis</i> But.	?	Larvae feed on old leaves in all plants and make frass chains.
	<i>Pentagonia</i> (35) <i>macrophylla</i> LS	S4	
	<i>Psychotria</i> (35) undetermined *RM LS	S?	
	<i>Calycophyllum</i> (35) <i>candidissimum</i> *DJ (81-SRNP-740)	?	

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>clytherea</i> <i>marcia</i> Fruh.	<i>Sabicea</i> (35) <i>villosa</i> LS, CAR, CVDO	S1	S1 Hostplants are 3 and 4 in bright sunshine. All instars rest on frass chains. Mallet reared this species from the same hostplant in Corcovado.
<i>fessonnia</i> Hew.	<i>Randia</i> (35) <i>echinocarpa</i> *DJ (79-SRNP-216) <i>Karstenii</i> *DJ (81-SRNP-725)	? ?	S2 These records are from solitary late instar larvae. Hostplants are 3 and 4.
<i>demialba</i> Butler	<i>Rondeletia</i> (35) undescribed species	*WH MV	? S3 This record is from a solitary late instar larva; hostplant was 3.
GENUS: <i>Hypanartia</i>			
<i>lethe</i> Fab.	<i>Phenax</i> (46) undetermined SJ	S2	S1 Hostplants are 1 and 3. <i>Phenax</i> is an herbaceous plant in open areas, <i>Trema</i> is a secondary succession tree. On <i>Phenax</i> the larva forms a cup from one leaf with silk and rests and eventually pupates in this silk lined cup.
	<i>Trema</i> (44) <i>micrantha</i> *DJ (81-SRNP-296)	?	S
GENUS: <i>Siproeta</i>	<i>Blechum</i> (1) <i>pyramidatum</i> *DJ (81-SRNP-71)	?	S2 Hostplants are all 1.
<i>stelenes</i>	undetermined CAR		
GENUS: <i>Chlosyne</i>	<i>Melanthera</i> (13) <i>aspera</i> SRNP	?	S2 Hostplant is 1.

<i>gaudialis</i> Bates	<i>Justicia</i> (1) undetermined	?	S2	Hostplant is 3, 4 ranging from seedling to mature plants.
<i>nerva bonplandi</i> Latr.	<i>Amaranthus</i> (2) undetermined LS	C?	G2	Hostplant is 2, 3 and 4 from seedling to mature plants.
<i>lacinia</i> Geyer	<i>Clibadium</i> (13) undetermined *RM LS	C?	G2	Reported from a mass of late instar larvae on an open area plant.
<i>melanarge</i> Bates	<i>Aphelandra</i> (1) <i>depihana</i> SRNP	C?	G2	Hostplant is 2, 3 and 4 ranging from juvenile to mature plants.
GENUS: <i>Eresia</i>				
<i>coela</i> Druce	<i>Justicia</i> (1) undetermined LS	C	G	Hostplants are 2, 3 and 4. <i>Justicia</i> are usually seedlings. I have not reared this species to adulthood on <i>Justicia</i> .
	<i>Herpetacanthus</i> (1) ADD <i>panamensis</i> *LEG LS	?	G	
GENUS: <i>Castilia</i>				
<i>ofella</i> Hew.	<i>Justicia</i> (1) <i>comaeta</i> LS	C	G2	Hostplants are 2, 3, 4 ranging from juvenile to mature plants. Eggs in clusters of 20 to 40.
	<i>Aphelandra</i> (1) <i>storkesi</i> *JM CVDO	?	G	
<i>myia</i> Hew.	<i>Justicia</i> (1) undetermined LS	C	G?	This record is from a single batch of 10 larvae feeding on a mature plant along forest edge.
<i>eranites</i> Hew.	<i>Justicia</i> (1) undetermined LS	C	G	I record this from several oviposition records all on seedling plants in open areas. First instar larvae feed on new leaves, later instars feed on all leaves. I have not reared this species to adulthood.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
GENUS: Anthanassa			
<i>ptolyca</i> Bates	<i>Justicia</i> (1) undetermined **AA (80-65) BCI	?	Aiello records this species from several late instar larvae feeding on all leaves. The host occurs along roadsides as an herbaceous weed.
GENUS: Tegosa			
<i>anieta</i> Hew.	<i>Vernonia</i> (13) undetermined VDS	C	Hostplants are 3 or riparian edges and range from juvenile to mature plants.
GENUS: Euptoeita			
<i>hegesia</i> <i>hoffmani</i> Comstock	<i>Turnera</i> (43) <i>ulmifolia</i> SRNP, CVDO	S2 S1	Hostplants are 1, either juvenile or mature plants.
FAMILY: NYMPHALIDAE			
SUBFAMILY: HELICONIINAE			
The bulk of the pertinent Costa Rican hostplant records has been reported by Benson, et. al. (1978) and the reader is referred to that paper. The records presented here complete their picture.			
GENUS: Dryadula			
<i>phaetusa</i> L.	<i>Passiflora</i> (31) <i>talamancensis</i> *JM CVDO	S	Mallet records this species from late instar larvae on host growing in open areas of bright sunshine.
GENUS: Eueides			
<i>vibilia</i>	<i>Passiflora</i> (31)		

<i>vialis</i> Stichel	<i>pittieri</i> *JM CVDO	?	G4	This species is reported from a solitary cluster of larvae on a host in the forest.
<i>procula</i> <i>vulgaris</i> But. & Dru.	<i>Erblichia</i> (43) <i>odorata</i> *DJ (79-SRNP-408)	?	G	Janzen reports that larvae make holes in the leaf as early instars and later feed at the leaf margins. The hostplant is a forest tree (see Janzen, 1983).

SUBFAMILY: ACRAEINAE

GENUS: Actinote

<i>leucomelas</i> Bates	<i>Mikania</i> (13) undetermined	SV, MV, CAR	C	Hostplant is 1, 2, 3 and 4.
<i>lapitha</i> Staud.	<i>Mikania</i> (13) <i>riparia</i> CVDO		?	Hostplant is 1 as a scandent semi-woody vine.

SUBFAMILY: DANAINAE

GENUS: Anetia

<i>thirza</i>	"genus?" (5)	COP	S1	This represents an oviposition record.
<i>insignis</i> Salvin	undetermined			

GENUS: Lycorea

<i>cleobaea</i>	<i>Carica</i> (12) <i>papaya</i> LS, SJ, ELROD	S1	S2	Hostplants are 2, 3, 4 and range from juvenile to mature forest trees. There is variation among hosts as to what plant fraction the larva will eat. Larvae feeding on Caricaceas cut the main and accessory veins of the leaf and then feed on the leaf portion distal to stem; presumably to avoid secondary compounds.
<i>atergatis</i> Dist.	<i>Jacarattia</i> (12) <i>dolichaula</i> LS	S1	S2	
	<i>Ficus</i> (26)	?	S3	
	undetermined *WH MV	?	S3	
	undetermined *DJ (79-SRNP-113)	?		
	<i>Matalea</i> (5)	?	S	
	<i>quirosi</i> *DJ (79-SRNP-131)	?	S	
	<i>Asclepias</i> (5)	?	S	
	<i>curassavica</i> *WH MV	?		

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>lliione</i> <i>albescens</i> Distant	<i>Ficus</i> (28) <i>tuerckheimii</i> *WH MV	?	S3 Haber reports this from a solitary late instar larva feeding on the stump sprouts of the host-plant growing in an open pasture.

SUBFAMILY: ITHOMIINAE

The bulk of Costa Rican hostplant records for the Ithomiinae may be found in Haber (1978). The following records are additions or new records to his list or records gathered by other workers. In addition in the present list I have included data pertinent to eggs, larvae and adults.

GENUS: Melinaea

<i>lilis</i>	<i>Markea</i> (40)	S1	Hostplant is an epiphyte, females oviposit on plants occurring as 2, 3, 4 and range from seedling to mature plants. Early instars cut leaf veins before feeding on tissues distal to the stem.
<i>imitata</i> Bates	<i>neurantha</i> LS, CAR	S1	In the insectary, females of Costa Rica stock preferentially oviposited on this plant (Mexican origin) although <i>Markea</i> and <i>Solandra</i> were present, and both are used by <i>lilis imitata</i> in the insectary. All hostplants are epiphytic.
<i>scy lax</i> Bates	<i>Juamuloa</i> (40) <i>mexicana</i> *LEG	S1	
<i>isthmia</i> Bates	<i>Solanum</i> (40) <i>lanceaefolium</i> *LEG LS <i>lanceaefolium</i> *RM LS <i>torvum</i> *FGS UCR	C	G2 Hostplants are 3 and 4 and range from seedlings to mature vines.
<i>menapis</i>	<i>Solanum</i> (40) <i>torvum</i> CAR	C	G2
<i>saturata</i> G & S	<i>torvum</i> *FGS UCR	C	G2 Hostplants are 3 or riparian edges.

GENUS: *Ithomia*

<i>patilla</i> Hew.	<i>Witheringia</i> (40) <i>solanacea</i> *FGS UCR undetermined species *LEG SV	S1 S2	Hostplants are 3 and 4.
<i>diasa</i>	<i>Lycianthes</i> (40) <i>heteroclita</i> *LEG LS	S1 S2	Hostplants are 3 and 4.
<i>hippocrennis</i> But. & Dru. <i>heraldica</i> Bates	<i>Acnistus</i> (40) <i>arborescens</i> SV, SJ, UCR, PAT <i>Witheringia</i> (40) undetermined species *LEG SV, LS	S1 S2	Hostplants are 1, 3 and 4 and may be young or mature plants.
<i>celemia</i>	<i>Capsicum</i> ? (40) undetermined species *LEG RCN	S1 S2	Hostplant occurs as an understory forest shrub and is acceptable as young and mature plants. Probably a <i>Witheringia</i> (S. Knap, pers. comm.).
<i>xenos</i> Bates	<i>Acnistus</i> (40) <i>arborescens</i> *FGS UCR	S1 S2	Hostplants are 2, 3 and 4.
GENUS: <i>Ceratinia</i>			
<i>tutia</i>	<i>Solanum</i> (40) <i>nudum</i> *LEG RCN	S1 S2	Hostplants are forest understory shrubs and usually occur in heavy second growth vegetation.
<i>dorilla</i> Bates	<i>nudum</i> *JM CVDO	S1 S2	
GENUS: <i>Direnna</i>			
<i>klugi</i> Geyer	<i>Solanum</i> (40) *all FGS <i>umbellatum</i> UCR	S1 S2	Stiles records all of these species from the same locality. All hostplants are 2, 3, 4 provided open sunshine during a part of the day.
	<i>torvum</i> UCR	S1 S2	
	<i>lanceolatum</i> UCR	S1 S2	
	<i>ochraceo-ferrugineum</i>	S1 S2	

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>euchytma</i> Feld.	<i>Solanum</i> (40) <i>ochraceo-ferrugineum</i> *LEG LS	S1 S3	Hostplants are 3 and riparian edges.
GENUS: Godyris			
<i>zygia</i> G & S	<i>Cestrum</i> (40) <i>nocturnum</i> *LEG SV	S1 S2	Hostplants are understory shrubs and range from juvenile to mature plants.
GENUS: Hyalyris			
<i>excelsa</i>	<i>Solanum</i> (40) <i>lanceolatum</i> SJ, CAR, VDS	S1 S1	Hostplants are 2, 3 or 4 and usually seedling or sapling plants.
<i>decumena</i> G & S			
GENUS: Hypothryis			
<i>lycaste</i>	<i>Solanum</i> (40) <i>umbellatum</i> *FGS UCR	S1 S2	No data on hostplant habitat.
<i>callispila</i> Bates	<i>ochraceo-ferrugineum</i>		
<i>euclea</i>	<i>Solanum</i> (40) <i>umbellatum</i> *FGS UCR	S1 S?	No data on hostplant habitat.
<i>valore</i> Haensch			
GENUS: Episcada			
<i>salvinia</i>	<i>Solanum</i> (40)		
<i>opleri</i> Lamas	“new species” *LEG SV	C G2	Hostplants are all second growth shrubs.
	<i>nudum</i> IFGS UCR	C G2	
GENUS: Pteronymia			
<i>agalla</i>	<i>Solanum</i> (40)		
<i>obscurata</i> Fab.	<i>brenesii</i> *LEG SV, RCN	S1 S?	No data on hostplant habitat.

<i>artena</i> Hew.			
<i>Lycianthes</i> (40)			
undetermined species *LEG	S?	Gilbert reports this from several eggs on old leaves. No data on hostplant habitat. Mallet	
<i>Solanum</i> (40)	S2	reports this from <i>Solanum</i> which are all second	
<i>confirme</i> "new species" *JM	S2	growth or forest understory shrubs.	
<i>Cyphomandra</i> (40)			
<i>costaricensis</i> *LEG SV	S1	Hostplant is 3 and 4. Females oviposit on mature plants.	
<i>lonera</i> But. & Dru.			

SUBFAMILY: MORPHINAE

GENUS: Antirrhrea

pterocapha G & S

Calypterygme (30)
undetermined CAR

?

Hostplants are solitary, growing in dense shade, usually at the base of a large tree. Larvae feed on leaves that are covered in epiphylls.

miltiades Fab.

Geonoma (30)
longivaginata *RM LS

?

S4 Marquis reports this from a solitary late instar larva from a plant growing in deep shade.

GENUS: Caerois

gerdrutus Stich.

Socratea (30)
undetermined *IC LS

?

S2 Chacon reports this from a solitary late instar larvae feeding on a large tree growing in a swamp.

GENUS: Morpho

amatontae Dey.

Pterocarpus (19)
officinalis CVDO

?

S2 I report this from a solitary late instar larva feeding on a sapling in a swamp. The larva fed nocturnally.

granadensis
polybapta Butler

Machaerium (19)
seemani

S? S2 I report this from solitary larvae feeding on

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>peleides</i> Esper	seedling and sapling sized plants. Females lay on this plant when confined.		
<i>Machaerium</i> (19)			
<i>seemanii</i> LS, CVDO, CAR, VSM	S	S2	Hostplants are all 2, 3, 4 and range from seedlings to large forest trees. Larvae are nocturnal feeders and usually crawl off the host leaves while not feeding.
undetermined LSALT	S?	S2	
undetermined VDS	S?	S2	
<i>biovulatum</i> *DJ (81-SRNP-711)	S?	S2	
<i>Mucuna</i> (19)	S2	S2	
<i>mutisiana</i> **confined female MV, SV			
<i>Dalbergia</i> (19)	S?	S2	
<i>retusa</i> *DJ (81-SRNP-226)	?	?	
<i>Pterocarpus</i> (19)			
<i>rohrrii</i> **AA (80-16) BCI			
SUBFAMILY: BRASSOLINAE			
GENUS: <i>Dynastor</i>			
<i>darius</i>			
<i>stygianus</i> Butler			
<i>Aechmea</i> (8)			
<i>magdalena</i> LS	S1	S2	Hostplants are 2, 3, 4 and in swampy areas if terrestrial. The undetermined record is an epiphyte. Larvae rest in the middle of the leaf blade when not feeding.
undetermined LS	?	S2	
<i>Bromelia</i> (8)			
<i>pinguin</i> *DJ (82-SRNP-1308)	?	S2	
GENUS: <i>Opsiphanes</i>			
<i>tamarindii</i>			
<i>sikyon</i> Fruh.			
<i>cassina</i>			
<i>fabricii</i> Boisd.			
<i>Acrocomia</i> (30)	S1	S2	Hostplants can be 1, 2, 3 or 4.
<i>vinifera</i> SRNP			
<i>Cocos</i> (30)	S1	S2	Hostplants are 1 or 3 and range from juvenile to large trees.
<i>nucifera</i> CVDO		S2	

<i>staudingeri</i> G & S	<i>Chusquea</i> (34) undetermined *IC PAT	?	S2	Chacon reports this species from late instar larvae which feed nocturnally.
<i>bogotanus</i> Distant	"palm" (30) undetermined *RC VDS inhabiting T8 Colombia (30)	?	S2	Cubero reports this from larvae feeding on juvenile plants growing in the forest.
GENUS: <i>Eryphanis</i>				
<i>polyxena</i>	"exotic bamboo" (34) *RC	S	S2	Cubero reports this from a confined female. Larvae feed nocturnally.
<i>lycomedon</i> Felder				
GENUS: <i>Caligo</i>				
**NOTE: It is common in the hostplant literature to record <i>Caligo</i> from <i>Musa</i> . However this certainly represents a radiation onto an introduced plant; <i>Musa</i> is native to the Old World. In my rearing studies of <i>Caligo</i> I have found that any species of <i>Heliconia</i> will serve as a hostplant and a stimulus to oviposit, both in the field and in the insectary.				
<i>memnon</i> Felder	<i>Heliconia</i> (21) various species SJ, SV, AT, CVDO	See Comments	G2	Hostplants are 2, 3 or 4 and range from juvenile to mature plants. Females oviposit small clusters of eggs (2-15) and then will lay several clusters on the same plant. All leaves are acceptable to all instars.
<i>eurylochus</i>	<i>Heliconia</i> (21) various species CAR, CVDO, LS, SV, LALT	C1	G2	Hostplants are 2, 3, 4 and range from juvenile to mature plants. Larvae are gregarious and aggregations may contain all instars without cannibalism.
<i>sulanus</i> Fruh.				
<i>atreus</i> Kollar	<i>Heliconia</i> (21) various species CAR, LS, CVDO <i>Asterogyne</i> (30) <i>marshiana</i> *IC LS	C2	G2	Hostplants are 2, 3, 4 and range from juvenile to mature plants. Chacon reports this record from a solitary late instar larva on old leaves. Usually larvae are in small aggregations where all instars may be present without cannibalism.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>illioneus</i> Kollar	<i>Heliconia</i> (21) various species LS	C2	Hostplants are 2, 3, 4 and may range from juvenile to mature plants. Larvae are gregarious and may be present in all instars without cannibalism.
GENUS: <i>Dulcedo</i>			
<i>polita</i> Hew.	<i>Geonoma</i> (30) undetermined LS	S1	Hostplants are all seedlings occurring in well shaded swampy areas.
	<i>Euterpe</i> (30) undetermined LS	S1	
GENUS: <i>Pierella</i>			
<i>luna</i> Fab.	<i>Heliconia</i> (21) <i>latispatha</i> CVDO	S2	Hostplants are all seedlings occurring in well shaded light gaps.
	<i>Calathea</i> (25) <i>marantifolia</i> *JM CVDO	?	
<i>helvetica</i>	<i>Heliconia</i> (21) various species LS, TUR	S2	Hostplants are all 2, 3, 4 and are seedlings except for <i>Panicum</i> .
<i>incanescens</i> G & S	<i>Astrophyne</i> (30) undetermined LS	S2	
	<i>Calathea</i> (25) undetermined LS	S2	
	<i>Panicum</i> (33) undetermined *MCS CVDO	S2	
GENUS: <i>Cyllopsis</i>			
<i>philodice</i> G & S			

undetermined COP, CDM	S1	S3	Hostplants occur in dense stands in open areas at high elevations.
<i>Chusquea</i> (34) undetermined COP, PAT	C?	*G2	Hostplants occur as dense stands along riparian edges or forest edges. Early instars are gregarious, later instars appear to be solitary (I have only lab reared them in later stages). Eggs are laid on young plants with abundant new growth.
<i>Cyperus</i> (15) <i>lazulae</i> BLH, CAR	S2	S1	Hostplants occur in open areas usually associated with swampy or riparian situations.
GENUS: <i>Oressinoma</i>			
<i>typhla</i> West. & Hew.			
<i>Olyra</i> (33) <i>latifolia</i> *DJ (81-SRNP-946)		?	Janzen reports this species from solitary larvae feeding on hostplants growing along forest edges.
GENUS: <i>Taygetis</i>			
<i>andromeda</i> Cr.			
<i>Neckeropsis</i> (29) <i>undulata</i> *MCS CVDO	S1	S2	Hostplant is an epiphytic moss on tree trunks in well shaded rainforest habitats. This is a highly unusual host record for any butterfly family.
GENUS: <i>Euptychia</i>			
<i>insolata</i> But. & Dru.			
<i>Selaginella</i> (38) <i>horizontalis</i> *MCS LS, CVDO, SV undetermined species ELRD undetermined CAR	S1	S2	Hostplants are 2, 3 and 4.
<i>westwoodi</i> Butler			
<i>Selaginella</i> (38) <i>arthriticum</i> *MCS LS, SV, CVDO, AT	S1	S2	Hostplants are 2, 3 and 4.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>molinae</i> Hub.	<i>Selaginella</i> (38) <i>arthriticum</i> *MCS SV undetermined CAR	S1 S2 S1 S2	Hostplants are 2, 3 and 4.
GENUS: <i>Chloreuptychia</i>			
<i>arnaeae</i> Fab.			
	<i>Eleusine</i> (33) undetermined *MCS SV, CVDO	S1 S2	Hostplants occur in isolated clumps in light gaps and at bases of trees.
	<i>Opismenus</i> (33) undetermined *MCS AT	S1 S2	
	<i>Ichnanthus</i> (33) undetermined *MCS LS undetermined CAR	S1 S2 S1 S2	
GENUS: <i>Cissia</i>			
<i>usitata</i> Butler	<i>Eleusine</i> (33) <i>indica</i> *MCS LS, CVDO	S2 S2	Hostplants occur along sunny edges and light gaps.
<i>themis</i> Butler	"various grasses" (33) undetermined *MCS AT	S2 S2	Hostplant occurs in small clumps in open forest.
<i>confusa</i> Staud.			
	<i>Panicum</i> (33) undetermined *MCS CVDO undetermined CAR	S2 S2 S2 S2	All hostplants occur as isolated clumps (grass) or seedlings (palms and <i>Calathea</i>) on the shady forest floor.
	<i>Euterpe</i> (30) undetermined *MCS LS CVDO	S2 S2	
	<i>Geonoma</i> (30) <i>congesta</i> LS, CAR	S2 S2	
	<i>Calathea</i> (26) undetermined LS	S2 S2	

labe Butler
"grasses" (33)
undetermined *MCS LS

palladia Butler
"various grasses" (33)
greenhouse data *MCS LS

Panicum (33)
undetermined LS, CVDO, SV
"grass" (33)
undetermined CAR

gomezi S, D & E
"grass" (33)
undetermined *MCS CVDO

Panicum (33)
maximum SRNP, AT
"various grasses" (33)
undetermined *MCS LS, CVDO, SV

"various grasses" (33)
undetermined *MCS AT, CVDO, SV

renata Cr.

hesione Sulzer
"various species" *MCS LS, CVDO,
SV

metallica Boisd.

hermes Fab.

The eggs are deposited singly off the hostplants, usually on nearby dead vegetation. The host occurs in lush, tall clumps in large light gaps.

In captivity, eggs are laid singly. All leaves of various grasses were acceptable to all instars.

Hostplants occur in isolated clumps in shady forest.

Hostplants occur in isolated clumps in shady forest.

Hostplants are 3 and 4 provided there is direct sunshine at some time of the day.

This is recorded from a solitary late instar larva feeding on a grass growing along forest edge.

?

Female oviposits off the hostplant on nearby vegetation and litter. Hostplants occur in very open forest or edges.

Hostplants occur as isolated clumps in open areas or along forest edges.

"grass" (33)

Hostplants occur in isolated clumps in the forest or along forest edges.

Eleusine (33)
undetermined *MCS LS, CVDO, SV

"grass" (33)

Hostplant almost always in open sunny areas, usually second growth, occasionally within the forest shade.

Species	Hostplant & Locality	Egg & Larval Data	Bionomic Notes of Host and Butterfly
<i>calixta</i> Butler	<i>Laparus</i> (15) <i>lazulae</i> BLH, CAR “grass” (33) undetermined CAR	S2 S2 S2	Hostplants occur in wet open areas along roadsides and trails, or riparian edges.
<i>polyphemus</i> Butler	<i>Chusquea</i> (34) undetermined *IC PAT	S2 S2	Chacon reports this species from late instar larvae feeding on a dense stand of hostplant along forest edges. The larvae feed nocturnally.
GENUS: <i>Megeuptychia</i>			
<i>antonoe</i> Cr.	<i>Calathea</i> (25) <i>lutea</i> *JM CVDO <i>Cyclanthus</i> (14) undetermined LS	C	Mallet reports this species from seedlings occurring in deep forest or along forest edges. My record is an oviposition as seen with binoculars while I was in the forest canopy.
GENUS: <i>Dioriste</i>			
<i>cothonides</i> GRS.-SMT.	<i>Chusquea</i> (34) undetermined COP, PAT	C	Hostplants occur as dense thickets along forest and riparian edges. Early instars are gregarious, later instars are solitary. The larvae feed nocturnally.
GENUS: <i>Eretris</i>			
<i>suzannae</i> DeVr.	<i>Chusquea</i> (34) undetermined BLH	S1	Hostplants occur as dense thickets along forest and riparian edges. The egg is deposited on the terminal spine of the leaf.

GENUS: **Pedaliodes***perperna* Hew.*Rhipidocladum* (34)
maxonii ESTS1 Hostplants occur in isolated, juvenile clumps on
the forest floor in old light gaps.*ereiba*
cremera G & S*Chusquea* (34)
undetermined COPS1 ? This is reported as an oviposition record only;
larvae were not reared past first instar. Females
lay eggs singly in the axils of the whorled leaves
near the main stem.GENUS: **Catargynnis***rogersi* G & S*Chusquea* (34)
undetermined BLH, COP, CARS2 ? Hostplants occur in dense stands in shady areas
in the forest or edges. Females lay eggs in leaf
axils; larvae not reared.



DeVries, Philip J. 1986. "Hostplant records and natural history notes on Costa Rican butterflies (Papilionidae, Pieridae & Nymphalidae)." *The Journal of Research on the Lepidoptera* 24(4), 290–333. <https://doi.org/10.5962/p.266791>.

View This Item Online: <https://www.biodiversitylibrary.org/item/224814>

DOI: <https://doi.org/10.5962/p.266791>

Permalink: <https://www.biodiversitylibrary.org/partpdf/266791>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: The Lepidoptera Research Foundation, Inc.

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <https://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.