

## SHORT COMMUNICATION

### Endosymbiotic Rickettsiales (Alphaproteobacteria) from the spider genus *Amaurobioides* (Araneae: Anyphaenidae)

F.S. Ceccarelli<sup>1</sup>, C.R. Haddad<sup>2</sup> and M.J. Ramírez<sup>1</sup>: <sup>1</sup>División de Aracnología, Museo Argentino de Ciencias Naturales, Av. Angel Gallardo 470, C1405DJR - Buenos Aires, Argentina; E-mail: saracecca@hotmail.com; <sup>2</sup>Department of Zoology & Entomology, University of the Free State, P. O. Box 339, Bloemfontein 9300, South Africa

**Abstract.** Endosymbiotic bacteria are commonly found in terrestrial arthropods and their effects have been studied extensively. Here we present the first recorded case of endosymbiotic bacteria found in the spider family Anyphaenidae. A fragment of the cytochrome oxidase c subunit I “barcoding” region belonging to unidentified Rickettsiales, presumably belonging to the genus *Rickettsia*, was sequenced from six individuals of *Amaurobioides africana* Hewitt, 1917.

**Keywords:** Bacteria, barcoding, intertidal

The gram-negative proteobacteria are one of the most widespread and diverse groups of bacteria, including medically important pathogens, free-living nitrogen-fixing organisms, and the order Rickettsiales, which contains obligate intracellular bacteria such as *Wolbachia* and *Rickettsia* that can be found living inside the cells of terrestrial arthropods (Ferla et al. 2013). These two genera have been the subject of numerous studies in arachnids, as they represent important pathogens in some cases (e.g., Paddock et al. 2010), while in others they are endosymbionts that manipulate their host's physiology, behavior and/or bias the host's sex ratio to favor their transmission (Rowley et al. 2004; Goodacre et al. 2006; Duron et al. 2008; Gunnarsson et al. 2009; Wang et al. 2010; Vanthournout et al. 2011; Goodacre & Martin 2012, 2013). *Rickettsia*-infected spiders have also been shown to display increased long-distance dispersal tendencies (Goodacre et al. 2009). Bacterial endosymbionts are usually transferred vertically in spiders, although there is evidence for horizontal transfer in closely related taxa (Baldo et al. 2008).

Early methods of detection of bacterial endosymbionts in insects and spiders relied on staining techniques (Cowdry 1923). With the advent of PCR-sequencing techniques, molecular detection of specific endosymbionts was made possible with relative ease, in the case of spiders resulting in targeted studies (e.g., Baldo et al. 2008; Jin et al. 2013) or sometimes as a byproduct of a study with a different aim (e.g., Řezáč et al. 2014 in *Dysdera microdonta* Gasparo, 2014). With bacteria-specific primers, amplification of endosymbiont DNA from potential host tissue provides positive results only for infected hosts. On the other hand, more “universal” primers may find the annealing sites in both host and symbiont, if said sites are conserved enough for the genomic region. One such case appears to be the “barcoding” fragment of the Cytochrome Oxidase C subunit I (COI), a protein-coding gene that appears to have its origins deep within the origins of life on the planet (Castresana et al. 1994). The fact that COI has highly conserved regions means that certain primers can be used across a wide range of organisms to amplify the same gene region. This is advantageous if the tissue used for DNA extraction exclusively belongs to one species. However, in the case of organisms hosting endosymbionts, this could be seen as a complication, although for large-scale barcoding studies, it has been shown to be manageable (Smith et al. 2012).

As part of a larger study on the intertidal anyphaenid genus *Amaurobioides* O. Pickard-Cambridge, 1883 (Araneae: Anyphaenidae) (Ceccarelli et al. in prep.), DNA was extracted from leg tissue of 19 individuals of *A. africana* Hewitt, 1917 and approximately 630 base-pairs of the COI gene fragment were amplified and sequenced using the primers LCOI 1490 (Folmer et al. 1994) and HCOoutout

(Prendini et al. 2005). For six out of the 19 individuals, the sequenced COI region did not belong to the targeted host species, but to an unknown species of the order Rickettsiales, presumed to be an intracellular symbiont. There was no variation in the nucleotides of the six sequences obtained, indicating that the *A. africana* individuals in this study were all infected with the same bacterial species. The sampling localities of the six infected specimens are shown in Fig. 1a and the COI sequences have been deposited in GenBank (accession numbers KU600819–KU600824).

The identification of the Rickettsiales COI sequences amplified in this study was based on comparisons to sequences available in the public databases *International Barcode of Life Database* (BOLD systems; <http://www.boldsystems.org/>) and *GenBank* (<http://www.ncbi.nlm.nih.gov/genbank/>). The information available from BOLD was minimal (the level of identification provided was to the order Rickettsiales) and there were cases of misidentification in GenBank, as the *BLAST*-queried sequences from this study were 99% identical to COI sequences labelled as “Hymenoptera sp.”, while the correct identification as proteobacteria started at 90% identity.

At this point, questions relating to why non-bacterial primers preferentially amplified COI regions of endosymbionts rather than host DNA in this study—even resulting in clean sequences (rather than a mix of host and symbiont amplicon)—remain largely unanswered. A visual inspection of the priming sites revealed that 9 Rickettsiales COI sequences downloaded from GenBank had 80–90% identity in the last 10 base-pairs towards the 3' end of the forward and reverse primers. This base-pair identity, coupled with the possibility of a very high number of endosymbiotic Rickettsiales, may have been enough to give initial preference and later exclusivity to the bacterial over the host DNA for primer annealing and DNA amplification during PCR. As mentioned earlier, the presence of endosymbionts is not thought to interfere with DNA barcoding of arthropods when using universal primers (Smith et al. 2012). However, the possibility still exists that COI sequences of endosymbionts are obtained when in fact the target organism is the host, as shown in this study, along with other isolated cases (e.g., Řezáč et al. 2014) and the presence of misidentified Rickettsiales COI sequences in GenBank (where the target organism was the host and thus the identification was placed as Hymenoptera sp.).

Of the COI sequences in GenBank from Rickettsiales (all belonging to the genus *Rickettsia*) with an identity score >75% for the sequences from this study, ten were selected for Bayesian phylogenetic analyses, along with a sequence from a closely related genus (*Orientia*, based on Weinert et al. 2009), four sequences of *Wolbachia* and a sequence belonging to *Anaplasma*, to root the tree. The COI sequences were






Figure 1.—a. Map showing localities where *Rickettsia*-infected specimens of *Amaurobioides africana* were collected; b. Bayesian phylogenetic tree of COI sequences for selected *Rickettsia* species and specimens from closely related genera, obtained from the NCBI database. Nodal support in Bayesian posterior probability (PP) represented by filled ( $0.95 < PP \leq 1$ ) and empty ( $0.9 < PP \leq 1$ ) circles. GenBank accession numbers are shown after taxon names in brackets. Terminal taxa labelled as NEW SEQUENCE are from this study. Colored boxes around taxon names represent host classes (blue = Insecta; red and yellow = Arachnida; arachnid orders: red = Acari; yellow = Araneae); c. General habitus of *A. africana*; d–e. Rock faces in the intertidal zone at De Hoop Nature Reserve, showing abandoned retreats (arrows) of *A. africana* (d), and at Jeffrey's Bay, showing sealed retreats of *A. africana* (e). Photos: C.R. Haddad.

aligned using TranslatorX (Abascal et al. 2010) and a partitioning strategy along with nucleotide substitution models for each partition (TrNef+I, TrN+G and TrN+I+G for COI 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> codon positions, respectively) chosen by PartitionFinder v.1.1.1 (Lanfear et al. 2012). A Bayesian phylogenetic tree was obtained by forming a consensus of 20,000 trees (minus 10% burn-in) from 20 million generations of Markov Chain Monte Carlo simulations performed in MrBayes v. 3.2.3 (Ronquist et al. 2012). Based on the phylogenetic tree obtained (Fig. 1b), the sequences from this study belong to an

unidentified *Rickettsia* species, closely related to a *Rickettsia* species infecting the spider *Dysdera microdonta*. Apart from being confident that *A. africana* can harbor the endosymbiont *Rickettsia*, a more in-depth study is required to fully understand the distribution, ecology and biology of the endosymbionts detected in this study.

Of particular interest in this relationship is the biology of the host spiders, which are exclusively found in the intertidal zone of rocky shores in marine habitats (Fig. 1c). The spiders regularly construct their silken retreats in rock faces (Fig. 1d, e), which they seal with silk



during high tide to avoid immersion in salt water, emerging at low tide to forage (Lamoral 1968). Therefore, transmission of the symbionts through the water medium in which the spiders occur seems unlikely. Apart from the most likely transmission pathway of the endosymbionts in *A. africana* being vertical transmission, the possibility of horizontal transmission should not be ruled out at this stage; a plausible additional explanation may be the transmission of the endosymbionts during ingestion of prey tissues, such as isopods, amphipods and dipterans that occur in the intertidal zone (Lamoral 1968). Further, the possibility that the same endosymbionts may infect various other spiders and pseudoscorpions occurring in the intertidal zone in South Africa (Lamoral 1968; Haddad & Dippenaar-Schoeman 2009; Larsen 2012; Owen et al. 2014), and the platygastriid wasp egg parasitoid of the only other truly exclusive intertidal spider in South Africa, *Desis formidabilis* (O. Pickard-Cambridge, 1890) (Desidae), viz. *Echthrodosis lamorali* Masner, 1968 (Owen et al. 2014), requires further investigation. Nevertheless, this study represents the first record of Rickettsiales in anyphaenids and is a contribution towards a broader understanding of proteobacteria in spiders.

#### ACKNOWLEDGMENTS

Candice Owen (Rhodes University, Grahamstown) and Dawn Larsen (Iziko South African Museum, Cape Town) are thanked for the loan of recently collected specimens that contributed towards this study. This study was supported from grant PICT 2011-1007 from ANPCyT and a postdoctoral fellowship from CONICET.

#### LITERATURE CITED

- Abascal, F., R. Zardoya & M.J. Telford. 2010. TranslatorX: multiple alignment of nucleotide sequences guided by amino acid translations. *Nucleic Acids Research* 38:W7–W13.
- Baldo, L., N.A. Ayoub, C.Y. Hayashi, J.A. Russell, J.K. Stahlhut & J.H. Werren. 2008. Insight into the routes of *Wolbachia* invasion: high levels of horizontal transfer in the spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA diversity. *Molecular Ecology* 17:557–569.
- Castresana, J., M. Lubben, M. Saraste & D.G. Higgins. 1994. Evolution of cytochrome oxidase, an enzyme older than atmospheric oxygen. *The EMBO Journal* 13:2516–2525.
- Cowdry, E.V. 1923. The distribution of *Rickettsia* in the tissues of insects and arachnids. *Journal of Experimental Medicine* 37:431–456.
- Duron, O., G.D.D. Hurst, E.A. Horne, J.A. Josling & J. Engelstädter. 2008. High incidence of the maternally inherited bacterium *Cardinium* in spiders. *Molecular Ecology* 17:1427–1437.
- Ferla, M.P., J.C. Thrash, S.J. Giovannoni & W.M. Patrick. 2013. New rRNA Gene-based phylogenies of the alphaproteobacteria provide perspective on major groups, mitochondrial ancestry and phylogenetic instability. *PLoS ONE* 8(12):e83383.
- Folmer, O., M.B. Black, W. Hoch, R.A. Lutz & R.C. Vrijehock. 1994. DNA primers for amplification of mitochondrial Cytochrome c Oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3:294–299.
- Goodacre, S.L. & O.Y. Martin. 2012. Modification of insect and arachnid behaviours by vertically transmitted endosymbionts: infections as drivers of behavioural change and evolutionary novelty. *Insects* 3:246–261.
- Goodacre, S.L. & O.Y. Martin. 2013. Endosymbiont infections in spiders. Pp. 93–105. *In Spider Ecophysiology*. (W. Nentwig, ed.). Springer-Verlag, Heidelberg.
- Goodacre, S.L., O.Y. Martin, D. Bonte, L. Hutchings, C. Woolley, K. Ibrahim et al. 2009. Microbial modification of host long-distance dispersal capacity. *BMC Biology* 7:32–39.
- Goodacre, S.L., O.Y. Martin, C.F.G. Thomas & G.M. Hewitt. 2006. *Wolbachia* and other endosymbiont infections in spiders. *Molecular Ecology* 15:517–527.
- Gunnarsson, B., S.L. Goodacre & G.M. Hewitt. 2009. Sex ratio, mating behaviour and *Wolbachia* infections in a sheetweb spider. *Biological Journal of the Linnean Society* 98:181–186.
- Haddad, C.R. & A.S. Dippenaar-Schoeman. 2009. A checklist of the non-Acarine Arachnids (Chelicerata: Arachnida) of the De Hoop Nature Reserve, Western Cape Province, South Africa. *Koedoe* 51(149):1–9.
- Jin, Y., C. Deng, H. Qiao, Y. Yun & Y. Peng. 2013. Molecular detection and phylogenetic relationships of three symbiotic bacteria in spiders (Araneae) from China. *Entomological News* 123:225–236.
- Lamoral, B.H. 1968. On the ecology and habitat adaptations of two intertidal spiders, *Desis formidabilis* (O.P. Cambridge) and *Amaurobioides africanus* Hewitt, at “The Island” (Kommetjie, Cape Peninsula), with notes on the occurrence of two other spiders. *Annals of the Natal Museum* 20:151–193.
- Lanfear, R., B. Calcott, S.Y.W. Ho & S. Guindon. 2012. Partition-Finder: Combined selection of partitioning schemes and substitution models for phylogenetic analyses. *Molecular Biology and Evolution* 29:1695–1701.
- Larsen, N. 2012. Predatory behaviour of Afrotropical Palpimanidae. *Newsletter of the African Arachnological Society* 25:9.
- Owen, C.A., J.A. Coetzee & S. van Noort. 2014. Distributional range of the South African maritime spider-egg parasitoid wasp, *Echthrodosis lamorali* (Hymenoptera: Platygasteridae: Scelioninae). *African Invertebrates* 55:323–332.
- Paddock, C.D., P.-E. Fournier, J.W. Sumner, J. Goddard, Y. Elshenawy, M.G. Metcalfe et al. 2010. Isolation of *Rickettsia parkeri* and identification of a novel spotted fever group *Rickettsia* sp. from Gulf Coast ticks (*Amblyomma maculatum*) in the United States. *Applied and Environmental Microbiology* 76:2689–2696.
- Prendini, L., P. Weygoldt & W.C. Wheeler. 2005. Systematics of the *Damon variegatus* group of Africa whip spiders (Chelicerata: Amblypygi): evidence from behavior, morphology and DNA. *Organisms, Diversity and Evolution* 5:203–236.
- Rezáč, M., F. Gasparo, J. Král & P. Heneberg. 2014. Integrative taxonomy and evolutionary history of a newly revealed spider *Dysdera ninnii* complex (Araneae: Dysderidae). *Zoological Journal of the Linnean Society* 172:451–474.
- Ronquist, F., M. Teslenko, F. van der Mark, D.L. Ayres, A. Darling, S. Höhna et al. 2012. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. *Systematic Biology* 61:539–542.
- Rowley, S.M., R.J. Raven & E.A. McGraw. 2004. *Wolbachia pipientis* in Australian spiders. *Current Microbiology* 49:208–214.
- Smith, M.A., C. Bertrand, K. Crosby, E.S. Eveleigh, J. Fernandez-Triana et al. 2012. *Wolbachia* and DNA barcoding insects: patterns, potential, and problems. *PLoS ONE* 7:e36514.
- Vanthournout, B., J. Swaegers & F. Hendrickx. 2011. Spiders do not escape reproductive manipulations by *Wolbachia*. *BMC Evolutionary Biology* 11:15–12.
- Wang, Z.-Y., C. Deng, Y.-L. Yun, C. Jian & Y. Peng. 2010. Molecular detection and the phylogenetics of *Wolbachia* in Chinese spiders (Araneae). *Journal of Arachnology* 38:237–241.
- Weinert, L.A., J.H. Werren, A. Aebi, G.N. Stone & F.M. Jiggins. 2009. Evolution and diversity of *Rickettsia* bacteria. *BMC Biology* 7:6–20.

Manuscript received 15 December 2015, revised 25 January 2016.





Ceccarelli, Fadia Sara, Haddad, C R , and  
Rami

rez,  
Marti

n J. 2016. "Endosymbiotic Rickettsiales (Alphaproteobacteria) from the spider  
genus Amaurobioides (Araneae: Anyphaenidae)." *The Journal of arachnology*  
44(2), 251–253. <https://doi.org/10.1636/joa-s-15-009>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/249644>

**DOI:** <https://doi.org/10.1636/joa-s-15-009>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/267191>

**Holding Institution**

Smithsonian Libraries

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: American Arachnological Society

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's



Ceccarelli, Fadia Sara, Haddad, Charles R, and  
Rami

rez,  
Marti

n J. 2016. "Endosymbiotic Rickettsiales (Alphaproteobacteria) from the spider  
genus Amaurobioides (Araneae: Anyphaenidae)." *The Journal of arachnology*  
44(2), 251–253. <https://doi.org/10.1636/joa-s-15-009>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/249644>

**DOI:** <https://doi.org/10.1636/joa-s-15-009>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/267191>

**Holding Institution**

Smithsonian Libraries

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: American Arachnological Society

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's



Ceccarelli, Fadia Sara, Haddad, Charles R, and  
Rami

rez,  
Marti

n J. 2016. "Endosymbiotic Rickettsiales (Alphaproteobacteria) from the spider  
genus Amaurobioides (Araneae: Anyphaenidae)." *The Journal of arachnology*  
44(2), 251–253. <https://doi.org/10.1636/joa-s-15-009>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/249644>

**DOI:** <https://doi.org/10.1636/joa-s-15-009>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/267191>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: American Arachnological Society

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's



Ceccarelli, Fadia Sara, Haddad, Charles R, and  
Rami

rez,  
Marti

n J. 2016. "Endosymbiotic Rickettsiales (Alphaproteobacteria) from the spider  
genus Amaurobioides (Araneae: Anyphaenidae)." *The Journal of arachnology*  
44(2), 251–253. <https://doi.org/10.1636/joa-s-15-009>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/249644>

**DOI:** <https://doi.org/10.1636/joa-s-15-009>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/267191>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Biodiversity Heritage Library

**Copyright & Reuse**

Copyright Status: In Copyright. Digitized with the permission of the rights holder

Rights Holder: American Arachnological Society

License: <https://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://www.biodiversitylibrary.org/permissions/>

This document was created from content at the **Biodiversity Heritage Library**, the world's

largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.