

Plants of Alpine and Subalpine Regions
Mt Albert, 1905, '06.

Equisetum arvense

Allen's Ravine up to 3000 ft.

E. pratense.

Allen's Ravine up to 1800 ft.

E. sylvaticum

Allen's Ravine, in wet places to the crest.

Selaginella selaginoides

Wet hornblende or serpentine slopes and in bogs.

Lycopodium Selago

Hornblende and serpentine slopes -

L. Selago, var. appressum

Hornblende crests.

L. lucidulum.

Woods of north slope, to timber-line.

L. lucidulum, var. parophyllum.

Camp at timber-line, north slope.

L. annotinum.

Slopes -

L. annotinum, var. pringlei -

Crests (hornblende)

L. sitchense

Scrub spruce, etc. of Flagstaff Peak, and at head of
Asbestos Brook.

L. alpinum.

Damp north slope, on hornblende.

Polypodium

Phegopteris polypodioides

Damp slopes, alpine meadows, etc.

P. Dryopteris

" " " " "

Adiantum pedatum, var. aleuticum

Serpentine from the Crests to Lac ^{au} Diab^{le}.

Locally on upper hornblende in ravines where seemingly washed down from Scip -

Pteris aquilina

Among low spruces along Snow Brook.

Cryptogramma densa

In dry serpentine, southern slope of Devil's Gulch -

Asplenium viride

1 small plant in wet " " " " " "

A. Filix-foemina

Abundant on damp slopes.

Polystichum scapulinum

Southerly slopes, on serpentine (usually dry) Asbestos Brook and Devil's Gulch.

P. Braconii

Allens Ravine below timberline.

Aspidium fragrans

on a boulder in the woods, Macoun's Brook.

A. spinulosum, v. dilatatum

Damp slopes.

Cystopteris bulbifera

Macoun's Brook.

Cystopteris fragilis.

Macoun's Brook. Allen's Ravine below 2500 ft.

C. montana

Macoun's Brook. Allen's Ravine below 1500 ft.

Woodсия ilvensis

Hornblende slope north of Flagstaff Peak.
Allen's Ravine.

Osmonda Claytoniana

Hornblende slopes.

Oxoclea Stuebelii.

Allen's Ravine, below 2000 feet.

Larix laricina

Scattered and dwarfed on serpentine to the summit.

Picea canadensis

Chiefly on Hornblende.

P. mariana.

Abundant but dwarfed on Serp., rare on Hornbl.

Abies balsamea.

General, but dwarf.

Thuja occidentalis.

Allen's Ravine below 1600 feet.

Juniperus communis, v. montana.

General, most abundant on serpentine.

Hieracium adonata

Bay south of Flagstaff Peak.

H. alpinum

Hornblende Crests & crevices.

Oryzopsis asperifolia

General in damp open places.

Milium effusum.

Macoun's Ravine. Allen's Ravine up to 3000 ft.

Phleum alpinum.

Common on Hornblende slopes.

Cinna latifolia

Common on slopes and meadows.

Agrostis borealis.

Common on slopes.

A. hymenalis, v. geminata.

Snow Brook, Devil's Gulch.

Calamagrostis canadensis.

Common

C. Langedorfii.

Hornblende slopes, near Crest.

C. neglecta, v. borealis.

On the serpentine by small pond above Allen's Ravine, also on serp. small pond at head of Ruisseau à Plaque à Malade.

Deschampsia caespitosa, v. alpina.

Common on seep.

D. flexuosa

Common

D. atropurpurea

Boys and meadows. Hambl. slopes, following
down brooks of north side to 2000 feet.

Melica stuiata.

Common

M. stuiata, forma albicans.

Common.

Trisetum subspicatum, v. molle.

Allen's Ravine.

T. melicoides.

Macoun's Brook.

Danthonia spicata.

Devil's Gulch. Snow Brook.

D. intermedia.

Bag, head of Asbestos Brook.

Asbestos Brook, Snow Brook, and Devil's Gulch.
Serpentine meadow E. of East Peak.

Poa alpina

Macoun's Brook. Allen's Ravine.

Poa —

At Allen's Ravine

Poa glauca

Macoun's Brook, Allen's Ravine below 1800 ft.

Festuca altaica

Common on serpentine.

Agropyrum biflorum

Bag south west of Flagstaff Peak.

A. novae-angliae?

Bag s.w. of Flagstaff Peak. Devil's Gulch. Snow Brook.

A. caninum

Allen's Ravine below 2000 ft.

Scirpus caespitosus

Abundant.

Eriophorum Chamissonis

Bag s.w. of Flagstaff Peak. Pond-holes in serpentine.

E. callitrix

Bag s.w. of Flagstaff Peak.

E. polystachion

Bag s.w. of Flagstaff Peak.

Carex aenea

Allen's Ravine, below 2000 ft.

C. tribuloides, var. reducta

Allen's Ravine, below 1800 ft.

Carex Diocypna, var. Bolanderi
Macoun's Brook.

C. trisperma

Bog s.w. of Flagstaff Peak.

C. lagopina

Springy spots and crevices of Hornblende, north slope.

C. canescens

Pond at head of Ruisseau à Plaque à Malade.

C. brunnescens.

Common

C. stellulata

Frequent in boggy places

C. stipata

C. xirpoides

Common

C. deflexa

Allen's Ravine

C. pedunculata.

Macoun's Brook.

C. vaginata

Bog s.w. of Flagstaff Peak.

C. laxiflora, v. leptonervia

Hornblende slopes.

C. arctata

Hornblende slopes.

C. capillaris

C. capillaris, var. elongata

Macoun's Brook.

Carex paupercula

Common by ponds on the Serpentine.

C. paupercula, var. irrigua.

Bogs and pond-shores.

C.



Fernald, Merritt Lyndon. 1905. "Plants of Alpine and Subalpine Regions, Mt. Albert, 1905, '06." *Diary of trip to Canada, 1904 June 29-August 19 and Plants of Alpine and Subalpine Regions, Mt. Albert, 1905-1906, 1904-1906*

View This Item Online: <https://www.biodiversitylibrary.org/item/237981>

Permalink: <https://www.biodiversitylibrary.org/partpdf/267421>

Holding Institution

Harvard University Botany Libraries

Sponsored by

CLIR

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.