

BASELINE VEGETATION ANALYSIS FOR A REMNANT BLACKLAND PRAIRIE IN WALKER COUNTY, TEXAS

Eric Keith

24 Summer Place
Huntsville TX, 77340
ek7275@cox.net

Nancy Hyde

958 N. Longfellow St.
Arlington VA 22205
nshesq@comcast.net

ABSTRACT

Baseline vegetation data on a remnant blackland prairie in Walker County, Texas was collected using permanent vegetation plots as a basis for monitoring future vegetation changes. This remnant prairie is privately owned and is part of working cattle and hay production ranch named Lone Oak Ranch. Historically, the prairie has been managed to preserve this rare and globally imperiled habitat.

KEYWORDS: Floristics, Walker County, Texas, vegetation

INTRODUCTION

Lone Oak Ranch includes one of the largest and most pristine and/or well recovered blackland prairies known in the Pineywoods of East Texas. The habitat concerned is defined as the Little Bluestem (*Schizachyrium scoparium*)-Indiangrass (*Sorghastrum nutans*) Community Series (Texas Natural Heritage Program 1993). This plant association can be more narrowly defined in eastern Texas as the Little Bluestem (*Schizachyrium scoparium*) – Missouri Coneflower (*Rudbeckia missouriensis*) – Narrowleaf Gumweed (*Grindelia lanceolata*) – Cusp Gayfeather (*Liatris mucronata*) Prairie or West Gulf Coastal Plain Fleming Calcareous Prairie (NatureServe 2005); it is ranked as a G1 community, meaning that it is considered “critically imperiled globally because of extreme rarity or because of some factor(s) making it especially vulnerable to extinction with typically 5 or fewer occurrences or very few remaining

Species	#1	#2	#3
<i>Berchemia scandens</i>		2067	
<i>Bumelia lanuginosa</i>		300	467
<i>Celtis laevigata</i>		267	
<i>Cornus drummondii</i>		7267	
<i>Crataegus crus-galli</i>		733	
<i>Crataegus spathulata</i>		200	
<i>Diospyros virginiana</i>		200	
<i>Forestiera ligustrina</i>		2267	
<i>Gleditsia triacanthos</i>	67	133	
<i>Ilex decidua</i>		267	
<i>Ilex vomitoria</i>		133	
<i>Juniperus virginiana</i>	200	67	
<i>Ligustrum sinense</i>		133	
<i>Lonicera japonica</i>		200	
<i>Pinus taeda</i>		67	
<i>Prunus angustifolia</i>		67	
<i>Quercus virginiana</i>		267	
<i>Rhamnus caroliniana</i>		533	
<i>Rubus trivialis</i>	733	2933	533
<i>Symphoricarpos orbiculatus</i>		7733	733
<i>Toxicodendron radicans</i>		333	
<i>Vitis aestivalis</i>		67	
All species	1000	30400	1733

Table 1. Woody species recorded in each plot in stems/ha. Woody species were counted in a 30-m X 5-m plot.

individuals (<1,000) or acres (<2,000) or linear miles (<10)” (Nature Conservancy 2000, NatureServe 2005).

This plant association is naturally rare in the West Gulf Coastal Plain, occurring only sporadically where areas of Vertisol soils are present over the Fleming geologic formation (Nature Conservancy 2000, NatureServe 2005). Very few high-quality examples are known, and most existing examples have been heavily degraded by past land uses, and many more have been completely converted to other land uses (Nature Conservancy 2000, NatureServe 2005, personal

observation). Primary threats to this community include land development, fire suppression, livestock grazing, oil and gas exploration, soil harvesting, and impacts from road development and vehicle use. The historical fire frequency of this grassland is estimated to range from 5 to 20 years (NatureServe 2005). Woody species such as Eastern red cedar (*Juniperus virginiana*), gum bumelia (*Bumelia lanuginosa*), Alabama supplejack (*Berchemia scandens*), rough-leaf dogwood (*Cornus drummondii*), saw greenbrier (*Smilax bona-nox*), coral berry (*Symphiorcarpos orbiculatus*), southern dewberry (*Rubus trivialis*) and honey locust (*Gleditsia triacanthos*) invade this community and change the physiognomy as a result of fire suppression (NatureServe 2005, personal observation).

Lone Oak Prairie is composed of approximately 120 acres of remnant and/or restored blackland prairie. Historical ground disturbance such as farming and intensive grazing occurred throughout the property. However, these historically disturbed areas are now composed primarily of native herbaceous species typical for blackland prairies such Indiangrass, little bluestem, lovegrass (*Eragrostis intermedia*), meadow dropseed (*Sporobolus compositus*), and Florida paspalum (*Paspalum floridanum*). The most heavily disturbed portions of the prairie have a significant component of the invasive species, King Ranch (KR) bluestem (*Bothriochloa ischaemum*). However, the portion of the prairie located in the southeastern corner of the property appears to have avoided or successfully recovered from most of the historical ground disturbance and is in near pristine condition. In addition, KR bluestem is less prominent in this portion of the prairie.

Lone Oak Prairie is particularly susceptible to encroaching development. Interstate Highway 45 and heavily-traveled Highway 30 to Bryan-College Station intersect less than a mile to the east of the property. The resulting development (including gas stations, restaurants, hotels, and a Wal-Mart) is typical of the structures found at such major intersections. A newly constructed local roadway (parallel to I-45) and the associated structures (including office buildings and a shopping mall) are located less than ½ mile to the east between I-45 and the property and visible from the southeastern corner of the prairie.

A subdivision and other residences are located less than $\frac{1}{2}$ mile to the west. Some of the residences to the west are also visible from the property. The surrounding forests are a matrix of pine-hardwood and bottomland hardwood forests with a dense shrub layer and sparse herbaceous understory.

MATERIALS AND METHODS

Plot establishment and measurement follow National Park Service protocol as outlined in the Western Region Fire Monitoring Handbook (1992) prepared by the Western Region Prescribed and Natural Fire Monitoring Task Force. Using this methodology, three brush plots were established in three different areas of the prairie. Each brush plot is a 30-meter transect line running in a north-south direction. Both ends of each transect were marked with rebar stakes and t-posts. To determine absolute cover of all species, all vegetation, contacting a 2-m tall rod placed perpendicular to the ground, was recorded every 30-cm along the 30-m transect for a total of 100 contacts. All woody species were recorded in a 30-m X 5-m plot along the transect. Herbaceous plant density was recorded using three 1-m x 1-m plots every ten meters along the transect. Plot 1 was established in a historically disturbed prairie where native herbaceous vegetation is becoming re-established. Plot 2 was established along the eastern edge of the prairie where woody species are beginning to encroach onto the grassland. Plot 3 was established in a near pristine portion of the prairie that is composed primarily of native herbaceous species. Nomenclature for species recorded in plots generally follows Diggs et al. (1999) and Turner et al. (2003). Vegetation was analyzed and quantitatively described following protocols outlined in the Fire Monitoring Handbook (NPS, 1992) and NPS National Fire Monitoring Handbook Software, Version 3.10.2.1 (Syndoriak, 1991).

RESULTS AND DISCUSSION

Data collected in this study provides baseline data in three portions of a remnant and/or recovering blackland prairie. This data will be used to monitor long term vegetation changes that occur from management activities such as prescribed burning, woody species

encroachment, climate changes, etc. Table 1 illustrates all woody species recorded; Table 2 illustrates absolute cover for all species recorded along the 30-m² transect; and Table 3 illustrates herbaceous species recorded in three 1-m² transects.

ACKNOWLEDGMENTS

We thank the Spencer Family Partners (Cheryl Spencer Patton, Carl Casey Spencer, and Joy Spencer Jechow), owners of the Lone Oak Prairie, for their support of this ongoing project.

LITERATURE CITED

- Diggs, G., B. Lipscomb and R. O'Kennon.** 1999. Shinnery's and Mahler's Illustrated Flora of North Central Texas. Sida, Bot. Misc. 16. Botanical Research Institute of Texas, Fort Worth.
- National Park Service.** 1992. Western Region Fire Monitoring Handbook, 3rd ed. NPS, Western Region, San Francisco.
- The Nature Conservancy.** 2000. The West Gulf Coastal Plain Ecoregional Conservation Plan. West Gulf Coastal Plain Ecoregional Planning Team, The Nature Conservancy, Nacogdoches, TX, USA.
- NatureServe Explorer:** An online encyclopedia of life [web application]. 2005. Version 1.6 . Arlington, Virginia, USA: NatureServe. Available: <http://www.natureserve.org/explorer>. (Accessed: November 12, 2005).
- Sydoriak, W.** 1991. FMH Software Manual (companion to Western Region Fire Monitoring Handbook). NPS, Western Region, San Francisco.
- Texas Natural Heritage Program.** 1993. Plant Communities of Texas (Series Level). Unpublished document. 26 pp.
- Turner, B.L., H. Nichols, G. Denny, O. Doron.** 2003. Atlas of the Vascular Plants of Texas. Sida, Bot. Misc. 24. Botanical Research Institute of Texas, Fort Worth.

Species (% Cover)	#1	#2	#3
<i>Agalinis heterophylla</i>			1.0
<i>Ambrosia psilostachya</i>	3.0	3.0	
<i>Ambrosia trifida</i>		2.0	
<i>Amphiachyris dranunculoides</i>	1.0	2.0	1.0
<i>Andropogon glomeratus</i>	5.0	7.0	7.0
<i>Aristida longespica</i>	1.0	2.0	1.0
<i>Aristida oligantha</i>	23.0	9.0	
<i>Bothriochloa ischaemum</i>			1.0
<i>Bothriochloa laguroides</i>	13.0	12.0	16.0
<i>Bromus japonica</i>		2.0	
<i>Carex cherokeensis</i>	9.0	2.0	5.0
<i>Carex microdonta</i>	3.0	2.0	1.0
<i>Cornus drummondii</i>		1.0	
<i>Croton monathogynus</i>	1.0	2.0	
<i>Cuscuta indecora</i>		5.0	2.0
<i>Dalea multiflora</i>	1.0	2.0	
<i>Desmanthus illinoensis</i>		3.0	
<i>Dichanthelium scribnerianum</i>	7.0	8.0	1.0
<i>Eragrostis intermedia</i>	5.0	6.0	
<i>Euphorbia bicolor</i>	11.0	7.0	
<i>Grindelia lanceolata</i>	10.0	2.0	2.0
<i>Hedyotis nigricans</i>		1.0	2.0
<i>Indigofera miniata</i>		1.0	
<i>Iva annua</i>	18.0	8.0	14.0
<i>Liatris mucronata</i>			1.0
<i>Muhlenbergia capillaris</i>		1.0	
<i>Neptunia lutea</i>	1.0		
<i>Oenothera speciosa</i>	1.0	1.0	
<i>Oxalis stricta</i>	1.0		
<i>Paspalum floridanum</i>		9.0	2.0
<i>Paspalum setaceum</i>	1.0		
<i>Rubus trivialis</i>		7.0	
<i>Rudbeckia missouriensis</i>	8.0	10.0	9.0

<i>Salvia lyrata</i>			1.0
<i>Schizachyrium scoparium</i>	5.0	30.0	51.0
<i>Setaria parviflora</i>	3.0	9.0	6.0
<i>Solidago canadensis</i>	2.0	1.0	4.0
<i>Sorghastrum nutans</i>	6.0	2.0	19.0
<i>Sporobolus clandestinus</i>	1.0	1.0	
<i>Sporobolus compositus</i>	26.0	17.0	24.0
<i>Symphyotrichum ericoides</i>	5.0		4.0
<i>Symphoricarpos orbiculatus</i>		7.0	
<i>Verbena halei</i>	1.0		
<i>Verbena xutha</i>	1.0		
Substrate	4.0	1.0	1.0

Appendix 1. All species recorded in cover transects in each plot according to absolute cover. Cover measurements were recorded every 0.3-m along a 30-m transect. Substrate includes leaf litter, duff, rocks, and bare soil.



Keith, Eric L and Hyde, Nancy. 2006. "Baseline vegetation analysis for a remnant blackland prairie in Walker County, Texas." *Phytologia* 88, 186–192.
<https://doi.org/10.5962/bhl.part.27429>.

View This Item Online: <https://www.biodiversitylibrary.org/item/48984>

DOI: <https://doi.org/10.5962/bhl.part.27429>

Permalink: <https://www.biodiversitylibrary.org/partpdf/27429>

Holding Institution

New York Botanical Garden, LuEsther T. Mertz Library

Sponsored by

The LuEsther T Mertz Library, the New York Botanical Garden

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Phytologia

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.