GORGONIA SANGUINOLENTA.

CHARACTER GENERICUS.

Animal crescens plantæ facie. Stirps cornea, ossea, vel lapidea. Cortex mollis, osculis polypiseris.

CHARACTER SPECIFICUS, &c.

GORGONIA depressa ramosissima dichotomopinnata, corticis slavi poris distichis purpureis.

Pall. el. 200ph. p. 175.


GORGONIA SANGUINOLENTA.

Lin. Syst. Nat. Gmel. p. 3087.

Luculente adeo et accurate explicuit hanc speciem celeberrimus Pallas in opere cui titulus Elenchus Zoophytorum ut ipsissima auctoris verba sine ulla immutatione transcribere non dubitaverim.

"Magnitudo pedalis vel ultra, multa tamen specimina palmo non altiora occurrunt. Truncus in eodem constanter plamo ramosus; Rami pinnati, ramulis passim sine ordine dichotomis, subdivisis, sepe

fæpe fastigiatis. Stirps tota depresso-plana. Lignum corneum, colore et substantia. Cortex calcareus, citrinus, extus lævis. Pori crebri, ramulis prominuli, in marginibus ramorum duplici utrinque serie dispositi, sparsive, halone ruberrime roseo inclusi. Pumila specimina pinnata, procera ramofiora sunt."





AN

PURPLE-SIDED GORGONIA.

GENERIC CHARACTER.

Animal growing in the form of a plant.

Stem of a horny, offeous, or even lapideous fubftance.

Bark foft, marked with numerous polype-cells.

SPECIFIC CHARACTER, &c.

Flat, branched, pinnated GORGONIA, with yellow bark and double-rowed purple pores.

Purple-spotted yellow GORGONIA.

The description of this Gorgonia, by Dr. Pallas, is so accurate, that it is unnecessary to add any thing to the account given by that celebrated Naturalist, who informs us that it sometimes grows to the height of a foot or more; though specimens often occur of much inferior size. The trunk is always branched a flat direction, with the branches pinnated; the smaller ones occasionally subdivided and dichotomous. The body or wood is of a horny substance and color; the bark yellow and smooth; the pores numerous, and projecting in the form of tubercles, disposed

disposed in a double series on the margins of the branches, each being surrounded by a very bright rose-coloured or purple border. The smaller specimens are commonly pinnate; the larger ones more ramissed. This elegant coral is a native of the Atlantic and of the American seas.

Purple-formed relies 100 200 ONLA.

Biggins of beneather that the best of all the control of

with serous, and projecting in the lorin of subscribes,



Shaw, George. 1796. "Purple-Sided Gorgonia, Gorgonia sanguinolenta [Pl. 264]." *The Naturalist's Miscellany* 8(LXXXVIII), https://doi.org/10.5962/p.310807.

View This Item Online: https://www.biodiversitylibrary.org/item/276320

DOI: https://doi.org/10.5962/p.310807

Permalink: https://www.biodiversitylibrary.org/partpdf/310807

Holding Institution

Museums Victoria

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.