

**ANTHRENUS MUSEORUM (COLEOPTERA:
DERMESTIDAE), AN EGG PREDATOR OF LYMANTRIA
DISPAR (LEPIDOPTERA: LYMANTRIIDAE)
IN CONNECTICUT AND A REVIEW OF DERMESTIDS
AS GYPSY MOTH EGG PREDATORS¹**

Paul W. Schaefer², R. S. Beal, Jr.³

ABSTRACT: Larvae of *Anthrenus museorum* were collected in egg masses of *Lymantria dispar* in Tolland Co., Connecticut, in March 1982. Field evidence and subsequent laboratory rearing confirmed predation on viable moth eggs. In Japan and South Korea, several new collection records and a possible new species of *Trogoderma* from Japan are reported. The first recovery of *Cryptorhopalum ruficorne* in Delaware is included. Dermestids in the role as predators on gypsy moth eggs worldwide are compiled in a table and briefly reviewed. Any impact appears to be directly density dependent and at high moth populations the impact may be significant. This form of egg predation has not been thoroughly studied.

Dermestid beetles (Coleoptera: Dermestidae) are known principally as pests of stored products, carpets, dried animal tissues (e.g. skins, furs, feathers, insect collections), and a wide variety of other organic matter (Griswold 1941; Hinton 1945). A catalogue of the Dermestidae of the world, including distributions, was compiled by Mroczkowski (1968).

The association of dermestid beetles with gypsy moth, *Lymantria dispar* (L.) (Lepidoptera: Lymantriidae), eggs was recognized long ago (Forbush & Fernald 1896; Burgess 1899) but the significance and impact of this association remain elusive. There have been many anecdotal accounts of finding larval dermestids feeding in gypsy moth egg clusters but little has been done to clarify the impact of this predation, with the possible exception of the study by Nonveiller (1959). Usually larvae are found to have hollowed out a portion of an egg mass. Few have determined to what extent this form of egg predation has contributed to gypsy moth egg mortality. In the process of tunneling within an egg mass, disruption of the protective setae covering the egg mass occurs. It has been suggested that this is important since it enhances parasitism by egg parasitoids, e.g., *Ooencyrtus kuvanae* (Howard) (Hymenoptera: Encyrtidae) and *Anastatus japonicus* Ashm. (Hymenoptera: Eupelmidae), by making more eggs in a cluster accessible to these parasitoids (Mason & Ticehurst 1984). Under field conditions, Nonveiller (1959) concluded that dermestids had a significant impact on gypsy moth egg survival under outbreak conditions, sometimes ac-

¹ Received July 24, 1995. Accepted September 30, 1995.

² USDA, Beneficial Insects Introduction Research Lab., 501 S. Chapel Street, Newark, Delaware 19713.

³ 1094 Pine Country Court, Prescott, Arizona 86303.

counting for up to 50% mortality of viable eggs; many dermestids were also found in old egg clusters where they were clearly feeding as necrophagans and not predators. At lower population levels the impact is greatly diminished (Nonveiller 1959). It appears that mortality of the gypsy moth egg stage due to dermestid predation is directly density dependent; this has yet to be experimentally confirmed.

FIELD COLLECTION

North America: The senior author collected a number of dermestid larvae in egg masses of gypsy moth at Willington, Tolland Co., Connecticut, on March 16, 1982. Egg masses with dermestid larvae were found under the eaves and roof overhang on a building adjacent to gypsy moth infested forests. At least one predatory larva was found in a gypsy moth egg mass on a *Quercus rubra* L. trunk adjacent to the building. As egg masses were scraped off the building, tunneling and apparent destruction of viable eggs were evident. Collected dermestid larvae were returned to the laboratory and allowed to complete development in the presence of the remains of the field collected egg masses or on laboratory reared eggs. In total, 14 adult dermestids were reared from the Willington material.

These adults, subsequently identified as *Anthrenus museorum* (L.)⁴ (Coleoptera: Dermestidae), were introduced into pint paper cartons and provided with laboratory reared gypsy moth egg masses. During exposure to fresh gypsy moth egg masses, viable eggs were deposited; subsequently a new generation of dermestids was reared exclusively on gypsy moth eggs. Development of the laboratory reared generation occurred at ambient laboratory room temperatures (ca. 18-26° C.) and emergence of the adults occurred in March, 1983. Thus partial development of the parental generation and complete development of the progeny occurred within one year. This record of *A. museorum* as a predator on viable gypsy moth eggs in Connecticut is the first North American record. It is also the first confirmed occurrence of the species in Connecticut, previous authors often mistaking *A. castaneae* Melsheimer for this species.

Gypsy moth egg masses collected in Newark, Delaware, on April 4, 1983, were isolated in pint paper cartons. The following month, three adults of the dermestid *Cryptorhopalum ruficorne* LeConte emerged, a first record of this association in Delaware, although *C. ruficorne* was reported as a predator of gypsy moth eggs in nearby states (Mason & Ticehurst 1984; Beal 1985).

Japan and South Korea: During the collection of gypsy moth egg masses in our previously reported study (1975-78) of egg parasitism (Schaefer *et al.* 1988), a collection of dermestids was obtained concurrently. Dermestid larvae were isolated and reared on available field collected gypsy moth eggs. Emerging adults were killed, pinned and labeled. Although some of those records

⁴ First identified in 1983 by J. M. Kingsolver, USDA, SEL, Beltsville, MD (now retired).

were published previously (Schaefer, 1980; 1981), all of our dermestid records are included in Table 1.

Deposition of Specimens: A synoptic collection of available dermestid material is deposited in the U. S. National Museum, Wash. D.C., The Carnegie Museum, Pittsburgh, Penna., and in the Gillette Entomological Museum at Colorado State University.

BRIEF REVIEW OF DERMESTIDS AS GYPSY MOTH EGG PREDATORS

In North America, there are relatively few references to the presence of Dermestidae in gypsy moth egg masses (Table 1). Burgess (1899) reported finding *A. verbasci* (L.) larvae feeding on gypsy moth egg masses in Massachusetts in October, and that adults appeared the following May. Hoebeke *et al.* (1985) illustrated and provided a key to identify eastern North American species of *Anthrenus*, including both *A. verbasci* and *A. museorum*, and Griswold (1941) studied the biology of *A. verbasci* in detail. Campbell (1967) stated that unidentified dermestid larvae were occasionally seen preying on gypsy moth eggs in northeastern New York. In central Pennsylvania, Brown and Cameron (1982) listed larvae of *Dermestes lardarius* L. as a predator associated with gypsy moth eggs. Also in Pennsylvania and in West Virginia, larvae of *C. ruficorne* were found attacking gypsy moth eggs (Mason & Ticehurst 1984) and, as mentioned above, P.W.S. confirmed *C. ruficorne* in this same role in Delaware. All these North American records and all other known records worldwide are included in Table 1, which is an updated version of the basic table of predators first compiled by Brown and Cameron (1982).

In other areas of the world, especially in Eurasia, there are considerably more records of dermestid associations with gypsy moth eggs. The most common genera are *Anthrenus*, *Attagenus*, *Dermestes*, *Megatoma* and *Trogoderma*. Of apparent lesser frequency are species in the genera *Globicornus*, *Ctesias*, and *Zhantievus*. Nonveiller (1959) found six new dermestid species feeding in gypsy moth egg masses and of these, *Megatoma pici* was the most abundant. Interestingly, Nonveiller found that larvae of *M. pici* were fairly mobile and larvae would readily move between different egg masses. All of these dermestid beetles appear highly opportunistic; no doubt any insect egg mass similar to those produced by gypsy moths will be suitable for attack. For example, in India, *Lymantria obfuscata* (L.), often referred to as the Indian gypsy moth, has the same egg laying behavior. Its eggs are reportedly attacked by "*Anthrenus* sp. prob. *museorum*", and an *Orphinus* sp. (Dharmadhikari *et al.* 1985). Howard (1897) and Burgess (1899) record dermestids also attacking eggs of white-marked tussock moth, *Orgyia leucostigma* (J. E. Smith) (Lymantriidae), which are deposited in a hardened foam-like material.

Table 1: World list of Dermestidae reported as predators (or apparent predators) of gypsy moth eggs, *Lymantria dispar* (L.) (Lepidoptera: Lymantriidae).

Species (Alphabetically listed)	Location	Reference/Collector/Year ^a
EUROPE AND ASIA		
<i>Anthrenus museorum</i> (L.)	Japan	Schaefer 1980
	South Korea	Schaefer (1976)
	Japan (Kyushu & Honshu)	Schaefer (1976)
<i>verbasci</i> (L.)	European SSR	Karnozhikii 1957
	Japan (Kyushu & Honshu)	Schaefer (1976)
	Japan	Schaefer 1980
	Ukraine	Kotenko 1982
<i>Attagenus unicolor unicolor</i> Brahm (?) (recorded as <i>A. piceus</i> Olivier)	Russia	Nonveiller 1959
<i>unicolor japonicus</i> Reitter sp.	Japan (Kyushu)	Schaefer (1976)
	Japan	Schaefer 1980
	Europe	Thompson & Simmonds 1964 ^b
<i>Ctesias serra</i> F.	Yugoslavia	Nonveiller 1959
<i>Dermestes ater</i> De Geer	Europe	Forbush & Fernald 1896
	Azerbaijan	Aliev <i>et al.</i> 1974
<i>bicolor</i> F.	Ukraine	Kotenko 1982
<i>erichsoni</i> Ganglbauer	Yugoslavia	Nonveiller 1959
	Bulgaria	Schedl 1936; Karnozhikii 1957; Nonveiller 1959
		Stefanov & Keremidchiev 1961 ^b
	Romania	Pirvescu 1978; Teodorescu 1980
	Russia	Shapiro 1956; Vorontsov 1950
		Nonveiller 1959
	Ukraine	Kotenko 1982
<i>lardarius</i> L.	Russia Far East	Kolomiets 1987
	France	Picard 1921; Nonveiller 1959
	Bulgaria	Schedl 1936; Karnozhikii 1957; Nonveiller 1959;
		Stefanov & Keremidchiev 1961
	Italy	Prota 1966
	Azerbaijan	Aliev <i>et al.</i> 1974
	Europe	Burgess & Crossman 1929 ^b
	Russia	Shapiro 1956; Vorontsov 1950
		Nonveiller 1959
	Ukraine	Kotenko 1982
<i>undulatus</i> Brahm	Italy (Sardinia)	Luciano & Prota 1983
	Russia	Shapiro 1956; Vorontsov 1950; Nonveiller 1959
	Ukraine	Kotenko 1982
sp.	Japan	Schaefer 1980
spp.	Yugoslavia	Nonveiller 1959
	Ukraine	Zelinskaya 1981

<i>Globicornus nigripes</i> F.	Yugoslavia	Nonveiller 1959
<i>Megatoma conpersa</i> Solskij	Russia Far East	Kolomiets 1987
<i>pici</i> Kalik	Yugoslavia	Nonveiller 1959 ^b ; 1976
<i>pubescens</i> Zetterstedt	Yugoslavia	Nonveiller 1959
<i>undata</i> L.	Yugoslavia	Nonveiller 1959
	Romania	Pirvescu 1978
<i>Orphinus</i> sp.	Japan	Schaefer (1976)
<i>Trogoderma</i> sp.	Ukraine	Kotenko 1982
<i>varium</i> (Matsumura & Yokoyama)	Japan (Kyushu & Shikoku)	Schaefer (1976)
sp. possibly undescribed ^c	Japan (Shikoku)	Schaefer (1976)
Unidentified	Japan	Howard 1910

AFRICA

<i>Anthrenus verbasci</i> (L.)	Morocco	DeLepiney 1930 ^b
<i>vladimiri</i> Menier & Villemant	Morocco	Menier & Villemant 1993
<i>Dermestes lardarius</i> L.	No. Africa	Thompson & Simmonds 1964 ^b
<i>Trogoderma versicolor</i> Creitz	Morocco	DeLepiney 1927 ^b , 1929, 1930 ^b , 1933
		Nonvieller 1959
<i>versicolor</i> var.		
<i>meridionale</i> Kraatz	Morocco	Hérard & Fraval 1980
sp.	Morocco	DeLepiney 1927 ^b
<i>Zhantievus lymantriae</i> Beal	Morocco	Beal 1992

NORTH AMERICA

<i>Anthrenus museorum</i> (L.)	Connecticut	Schaefer (1982) ^d
<i>verbasci</i> (L.)	Massachusetts	Burgess 1899; Howard 1910
<i>Cryptorhopalum ruficorne</i> LeConte	Pennsylvania & West Virginia	Mason & Ticehurst 1984; Beal 1985
	Delaware	Schaefer (1983) ^e
<i>Dermestes lardarius</i> L.	North America	Griffiths 1976 ^f
	Pennsylvania	Brown & Cameron 1982
<i>Trogoderma</i> prob. <i>ornatum</i> (Say)	Massachusetts	Howard 1910
(listed as <i>T. tarsale</i> Melsheimer)		
Unidentified	New York	Campbell 1967 ^b

^a Years given in parentheses are years of specimen collection. Specimens collected by the senior author in the 1970's were identified by Sadanari Hisamatsu (1977) and/or by John M. Kingsolver (1979). Years not in parentheses refer to date of publication; see Literature Cited.

^b Cited in Griffiths 1976.

^c D. G. H. Halstead, *in litt.*

^d Identified by J. M. Kingsolver (1983) and confirmed by R.S.B., Jr. (1995).

^e Identified by J. M. Kingsolver (1984), examined by R.S.B., Jr. (1985).

^f Listed in Griffiths' (1976) table but text does not substantiate a North American record associated with egg masses, however a record coming from a pupal mass is recorded.

We believe that, in time, many other associations will become known as other species of opportunistic dermestids are found to take advantage of the nutritional resources and protected niches represented by individual masses containing hundreds of gypsy moth eggs. This will be especially true as the invading gypsy moth in North America moves into new geographical areas and comes in contact with other dermestid species for the first time.

ACKNOWLEDGMENTS

We especially thank D. G. H. Halstead of Central Science Laboratory, Slough, Bucks, U.K.; John M. Kingsolver, USDA, Systematic Entomology Lab., Beltsville, MD, and Sadanari Hisamatsu, Ehime University, Matsuyama, Japan, for the identification of dermestids at various times. Thanks also to Joseph M. Tropp for laboratory rearing the dermestids at Newark. Kevin W. Thorpe and Jeffrey C. Miller and two anonymous reviewers offered valuable suggestions on this manuscript.

LITERATURE CITED

- Aliev, A. A., R. E. Effendi & A. M. Mamedov. 1974. (Little known insect enemies of the gypsy moth) Zash. Rast. 5:36. (in Russian) (Rev. App. Entomol. (A) 64:6204).
- Beal, R. S., Jr. 1985. A taxonomic revision of the Nearctic species of *Cryptorhopalum* (Dermestidae: Coleoptera). Trans. Amer. Entomol. Soc. 111:171-221.
- Beal, R. S., Jr. 1992. A new genus and species of Dermestidae (Coleoptera) from Morocco. Coleop. Bull. 46:403-408.
- Brown, M. W. & E. A. Cameron. 1982. Natural enemies of *Lymantria dispar* (Lep.: Lymantriidae) eggs in central Pennsylvania, U.S.A., and a review of the world literature on natural enemies of *L. dispar* eggs. Entomophaga 27:311-322.
- Burgess, A. F. 1899. An egg-eating beetle. pp. 475-477, In Report on the work of exterminating the gypsy moth by the state (Massachusetts) Board of Agriculture, January 1899. Publ. Doc. No. 4, Gypsy moth appendix.
- Burgess, A. F. & S. S. Crossman. 1929. Imported insect enemies of the gypsy moth and the brown-tail moth. USDA Tech. Bull. 86, 148 pp.
- Campbell, R. W. 1967. Analysis of numerical change in gypsy moth populations. For. Sci. Monogr. 15:1-33.
- DeLepiney, J. 1927. Les insectes nuisibles du chêne liège dans la forêt de la Momora (Maroc). Ann. Epiphyt. 13:145-174. (Rev. Appl. Entomol. (A) 16:314).
- DeLepiney, J. 1929. Les insectes nuisibles du chêne liège dans la forêt de la Momora (Maroc), deuxième étude. Ann. Epiphyt. (14:313-321. (Rev. Appl. Entomol. (A) 18:85).
- DeLepiney, J. 1930. Contribution à l'étude du complexe biologique de *Lymantria dispar*. Mem. Soc. Sci. Nat. Maroc. 23:1-100. (Abst. in Rev. Appl. Entomol. (A) 18:569).
- DeLepiney, J. 1933. Le rôle de la direction des eaux et forêts du Maroc et de l'Institut Scientifique Cherifien dans la lutte biologique entreprise contre *Lymantria dispar* à l'aide de *Schedius kuwanai*. Congr. intern. Entomol. Paris 1932 5:807-812. (Cited in Mason & Ticehurst 1984, not seen by the authors).
- Dharmadhikari, P. R., G. Ramaseshiah & P. D. Achan. 1985. Survey of *Lymantria obfuscata* and its natural enemies in India. Entomophaga 30:399-408.
- Forbush, E. H. & C. H. Fernald. 1896. The gypsy moth: *Porthetria dispar* (Linn.). Wright & Potter Printing Co, Boston, 495 pp.
- Griffiths, K. J. 1976. The parasites and predators of the gypsy moth: A review of the world literature with special application to Canada. Canadian For. Ser., Dept. of the Environ., Sault Ste. Marie, Ont., Report 0-X-243, 92 pp.

- Griswold, G. H.** 1941. Studies on the biology of four common carpet beetles. Part 1. The black carpet beetle (*Attagenus piceus* Oliv.), the varied carpet beetle (*Anthrenus verbasci* L.), and the furniture carpet beetle (*Anthrenus vorax* Waterh.). Cornell Univ., Agric. Exp. Stn. Memoir 240, pp. 1-57.
- Hérard, F. & A. Fraval.** 1980. La répartition et les ennemis naturels de *Lymantria dispar* (L.) (Lep.: Lymantriidae) au Maroc, 1973-1975. Acta Oecol. Appl. 1:35-48.
- Hinton, H. E.** 1945. A monograph of the beetles associated with stored products. Vol. 1. British Museum, London. 443 pp.
- Hoebeker, E. R., A. G. Wheeler, Jr. & R. S. Beal, Jr.** 1985. *Anthrenus pimpinellae* F., a palearctic dermestid established in eastern North America (Coleoptera: Dermestidae). J. New York Entomol. Soc. 93:1216-1222.
- Howard, L. O.** 1897. A study in insect parasitism. U. S. Dept. Agric., Bureau Entomol. Tech. Ser. 5, 57 pp.
- Howard, L. O.** 1910. Technical results from the gypsy moth parasite laboratory. I. The parasites reared or supposed to have been reared from the eggs of the gypsy moth. U. S. Dept. Agric., Tech. Ser. Bull. 19, 12 pp.
- Karnozhikii, N.** 1957. (Observations on the gypsy moth, *Ocneria dispar* L., outbreak during 1952-1955 with reference to the possibility of biological control.) Trudove na zoologicheskii institut (Sofia) 4:1-102. (in Bulgarian). (Cited in Nonveiller 1959, not seen by the authors).
- Kolomiets, N. G.** 1987. (Insects — parasites and predators of the gypsy moth (*Lymantria dispar* L. (Lepidoptera)) of the Asiatic part of the USSR). Izvestiya Sibirskogo Otdeleniya Akademii Nauk SSSR, Biologicheskikh Nauk 6:83-89. (in Russian, Engl. summ.) (Rev. Appl. Entomol. (A) 76:2046).
- Kotenko, A. G.** 1982. Hide beetles (Coleoptera: Dermestidae) — natural enemies of the gypsy moth in the southern Ukraine. Vestnik Zoologii 1:41-45. (in Russian) (Biol. News & Info. 4:1769).
- Luciano, P. & R. Protà.** 1983. Planning *Lymantria dispar* L. control in Sardinia. In 10th Intern. Congr. Plant Prot. 1983, Vol. 3. Proc. conference at Brighton, England, 20-25 Nov. 1983. Plant. Prot. for Human Welfare, Croydon, UK; British Crop Prot. Council 1074 pp. (Rev. Appl. Entomol. (A) 72:3255).
- Mason, T. L. & M. Ticehurst.** 1984. Predation of *Cryptorhopalum ruficorne* (Coleoptera: Dermestidae) on egg masses of the gypsy moth, *Lymantria dispar* (Lepidoptera: Lymantriidae). Can. Entomol. 116:1675-1677.
- Menier, J. J. & C. Villemant.** 1993. Description et biologie de *Anthrenus* (*Peacockia* n. s.gen.) *vlaafimiri* n. sp., Prédateur des pontes de *Porthetria dispar* (L.) au Maroc (Coleoptera: Dermestidae; Lepidoptera: Lymantriidae). Revue Française D'Entomologie 15:61-66. (with Engl. summ.)
- Mroczkowski, M.** 1968. Distribution of the *Dermestidae* (Coleoptera) of the world with a catalogue of all known species. Annales Zoologici (Polska Akademia Nauk) 26:15-189.
- Nonveiller, G.** 1959. Predators of gypsy moth eggs found in Yugoslavia during the 1945-1950 gypsy moth plague. Zast. Bilja 10:15-35. (Transl. from French).
- Nonveiller, G.** 1976. Predators of gypsy moth eggs found in Yugoslavia during the 1945-1950 gypsy moth plague. Selected Articles from Plant Protection, 1950-1970. Belgrade, Yugoslavia, Nolit Publ. House, pp. 451-475 (Technical Transl. no. 72-56059).
- Picard, F.** 1921. Le bombyx disparate ou spongieuse. Progrès Agricole et Viticole 79:160-165 (Cited in Nonveiller 1959, not seen by the authors).
- Pirvescu, D.** 1978. Développement des gradations de *Lymantria dispar* dans les peuplements de robinier de l'inspectariat forestier Dolj. Zast. Bilja 29:151-154.
- Protà, R.** 1966. Contributi alla conoscenza dell'entomofauna della quercia da sughero (*Quercus suber* L.). V. Osservazioni condotte in Sardegna su *Ooencyrtus kuwanai* (Howard) (Hymenoptera: Encyrtidae) nuovo per la fauna Italiana. Stazione Sper. Sughero, Tempio Pausania. Memoria 17, 26 pp. (in Italian, Engl. summ.)
- Schaefer, P. W.** 1980. Natural enemies of gypsy moth (*Lymantria dispar*) in Japan and Korea,

- especially new and potentially useful species. 16th Int. Congr. Entomol. Kyoto, Japan. Abst. 9P,2, p. 297.
- Schaefer, P. W.** 1981. Explorations in Japan and Korea by ARS Asian Parasite Laboratory, 1975-1977. pp. 340-348. In Doane, C. C. & M. L. McManus (eds.), *The Gypsy Moth: Research Toward Integrated Pest Management*. U.S. Dept. Agric. Tech. Bull. no. 1584, 757 pp.
- Schaefer, P.W., K. Kanamitsu & H.-L. Lee.** 1988. Egg parasitism in *Lymantria dispar* (Lepidoptera: Lymantria) in Japan and South Korea. Kontyu, Tokyo 56:430-444.
- Schedl, K.** 1936. Der Schwammspinner (*Porthetria dispar* L.) in Euroasien, Africa and Neuengland. Monogr. zur. angew. Entomol. 12:1-242.
- Shapiro, V. A.** 1956. (Main parasites of the gypsy moth (*Porthetria dispar* L.) and prospects for their use.) Zool. Zhurnal 35:251-265 (in Russian) (Cited in Nonveiller 1959, not seen by the authors).
- Stefanov, D. & M. Keremidchiev.** 1961. The possibility of using some predators and parasitic insects in the biological control of the gypsy moth (*Lymantria dispar*) in Bulgaria. Nauk. Trud. Vis. Lesctekj. Inst. 9:157-168. (Rev. Appl. Entomol. (A) 53:170).
- Teodorescu, I.** 1980. (Beneficial insect fauna (predators) in the woods of northern Oltenia). Studii si Cercetari de Biologie, Biologie Animala 32:3-6. (in Romanian, Engl. summ.).
- Thompson, W. R. & F. J. Simmonds.** 1964. A catalogue of the parasites and predators of insect pests. Section 3. Predator host catalogue. Imper. Agric. Bur., Commonw. Inst. Biol. Contr., 204 pp.
- Vorontsov, A. I.** 1950. (Dermestid beetles as predators of the gypsy moth) Zool. Zhurnal 29:406-416 (in Russian) (Cited in Nonveiller 1959, not seen by the authors).
- Zelinskaya, L. M.** 1981. (Factors causing mortality of eggs of the gypsy moth in forest plantations in the south of the Ukrainian SSR.) pp. 52-54. In Aukshtikal' nene, A. M. (ed.) Noveishie dostizheniya lesnoi entomologii (po materialam USh s"ezda VEO, Vil'nyus 9-13 Oktyabraya 1979g.) (in Russian) (Rev. Appl. Entomol. (A) 72:4803.).



Schaefer, Paul W. and Beal, Richard S. 1996. "Anthrenus Museorum (Coleoptera: Dermestidae), An Egg Predator Of Lymantria Dispar (Lepidoptera: Lymantriidae) In Connecticut And A Review Of Dermestids As Gypsy Moth Egg Predators." *Entomological news* 107, 143–150.

View This Item Online: <https://www.biodiversitylibrary.org/item/20638>

Permalink: <https://www.biodiversitylibrary.org/partpdf/31158>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: American Entomological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.