



AILURCEDUS BUCCOIDES,

AILURÆDUS BUCCOIDES.

Barbet-like Cat-bird.

Kitta buccoides, Temm. Pl. Col. 575.—Von Rosenb. J. f. O. 1864, p. 122.

Cissa buccoides, Gray, Gen. B. iii., App. p. 14.

Ptilonorhynchus buccoides, Gray, P. Z. S. 1858, p. 194.—Id. Cat. Mamm. & B. N. Guin. p. 37.—Schl. Mus. P. B., Coraces, p. 118.—Gray, Hand-l. B. i. p. 294.—Schl. N. T. D. iv. p. 49.

Ælurædus buccoides, Elliot, Monogr. Parad. pl. 36.—Sclater, P. Z. S. 1873, p. 697.

ALTHOUGH this bird, aptly named *buccoides* by Temminck on account of its resemblance to the Barbets of the genus *Megalæma*, has been known for so many years, and specimens not unfrequently turn up in collections from New Guinea, there is nothing recorded of its habits or economy. It is confined to the islands of New Guinea and Waigiu, the Leiden Museum possessing a considerable number of specimens collected in these localities by the late Dr. Bernstein. The original specimen was procured at Triton Bay, on the west coast of New Guinea; and other localities mentioned by Professor Schlegel are Sorong, the north coast of Salwatti, and the island of Batanta.

In its general green coloration this bird resembles the other species of the genus; but it may be readily distinguished by the more decided spotting of the breast and by the very distinct whitish streaks which are seen on the nape of the neck; besides which it is by far the smallest species of the whole genus, a group of birds intimately allied to the Bower-birds (*Chlamydodera* and *Ptilonorhynchus*); but up to the present moment we have no information as to whether the members of the genus *Ailurædus* ever construct a bower.

Upper surface of body, the wings included, grass-green, the secondaries slightly tipped with yellow; head olive-brown washed with green; all the feathers of the neck pale yellow, with large terminal spots of black, giving a streaked appearance to this part; a great many of the interscapular plumes barred with yellow, some spotted with black; tail green; sides of face and throat buffy white, the former thickly spotted with black; rest of under surface ochraceous buff washed with green, with a black spot to each of the feathers, disappearing on the under tail-coverts; the spots greenish on the lower abdomen; under wing-coverts salmon-buff, the outermost with brown-mottled bases; bill yellowish white; tarsi and feet lead-colour. Total length 10 inches; culmen 1·0. wing 4·9, tail 3·6, tarsus 1·4.

Apparently there is no difference in the sexes; and the figures in the accompanying Plate are as nearly life-size as possible.



Gould, John and Sharpe, Richard Bowdler. 1875. "Ailuroedus buccoides, Barbet-like Cat-bird [Pl. 41]." *The birds of New Guinea and the adjacent Papuan islands : including many new species recently discovered in Australia* 1(I), –.
<https://doi.org/10.5962/p.322640>.

View This Item Online: <https://www.biodiversitylibrary.org/item/229855>

DOI: <https://doi.org/10.5962/p.322640>

Permalink: <https://www.biodiversitylibrary.org/partpdf/322640>

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.