

A PRELIMINARY INVESTIGATION OF THE CADDIS-FLIES (INSECTA: TRICHOPTERA) OF THE QUEENSLAND WET TROPICS

Ken Walker, Arturs Neboiss, John Dean¹, David Cartwright²

Museum of Victoria, Department of Entomology, 71 Victoria Crescent, Abbotsford, Victoria 3067

Abstract

A checklist of the trichopteran fauna of the Queensland Wet Tropics is presented. The species composition of at least 217 taxa, including 95 new species and four new genera, is summarised and discussed with reference to the overall Australian fauna and available information on the Trichoptera fauna of several eastern Australian areas.

Introduction

The Queensland Wet Tropics (QWT), including the Wet Tropics World Heritage Area, lies along the east coast of Australia between Cooktown in the north and Townsville in the south, extending inland to the Atherton Tableland (Fig. 1.). Proclamation of the Wet Tropics World Heritage Area was recognition of the environmental significance and sensitivity and the cultural importance of the region. The Wet Tropics Management Authority funded several studies of the local fauna and flora and this paper summarises our findings on the Trichoptera fauna.

The study was based on material from 151 sites held in various Australian museum collections (22,549 specimens: 10,527 ♂♂, 9213 ♀♀, 2297 larvae and 512 pupae). Sites were grouped into investigation areas numbered 1-10 (Fig. 1), which enabled a preliminary assessment of regional variations in the Wet Tropics Trichoptera fauna, using the PATN programs Decorana and Twinspan (Belbin 1988). The criteria for grouping sites into investigation areas was latitudinal except for areas 4-7. These areas occupy similar latitudes but represent vastly different habitats. Investigation area 5 represents low coastal areas, while area 7 comprises sites on the Atherton Tableland. Investigation area 4 (Kuranda district) represents a region between the lowlands and the Atherton Tableland, while area 6 (Bellenden Ker and Mt Bartle Frere) represents high altitude sites.

Results

Taxa identified during this study are listed in Table 1. The number of taxa is greater than the actual number of species present, as adults and immatures of some species have been listed as separate taxa due to the lack of association of life stages. The checklist represents, in effect, a taxonomic appraisal of all families found within the QWT. The specific epithet of each new species has been assigned a genitalia preparation number unique to that species (eg. PT-2010 or CT-221) and such numbers will be included in future published descriptions.

Present address:

¹Environment Protection Authority, 27 Francis Street, Melbourne, Vic., 3000

²Werribee Treatment Complex, Private Bag 10, P.O. Werribee, Vic., 3030




Fig. 1. Map of Queensland Wet Tropics, including the World Heritage Area (shaded) and showing collection sites (diamond symbols) and investigation areas (circled and numbered). Note: Star symbols mark major towns.

Table 1 (cont.). Trichoptera species checklist for the Queensland Wet Tropics, with presence or absence within each investigation area noted. (Note: L - denotes taxa recorded as larvae only.)

INVESTIGATION AREAS	1	2	3	4	5	6	7	8	9	10
Hydrobiosidae										
<i>Apsilochorema gisbum</i> (Mosely)	+	+	-	-	-	-	+	+	+	-
<i>Apsilochorema obliquum</i> (Mosely)	-	+	+	-	+	+	+	-	+	+
<i>Ethochorema brunneum</i> (Mosely)	+	+	+	+	+	+	+	+	+	+
<i>Ptychobiosis nigrita</i> (Banks)	-	+	+	-	+	+	+	-	+	+
<i>Ulmerochorema seona</i> (Mosely)	-	-	+	-	-	-	-	-	-	-
<i>Ulmerochorema stigma</i> (Ulmer)	+	+	+	+	+	-	+	+	+	+
<i>Ulmerochorema</i> sp. nov. PT-1036	-	+	+	-	+	+	+	-	+	+
Hydropsychidae										
<i>Aethaloptera sexpunctata</i> (Kolenati)	-	-	-	-	+	+	-	+	-	-
<i>Asmicridea</i> sp.	-	+	-	+	+	+	+	+	+	-
<i>Asmicridea</i> sp. 4 (L)	-	-	+	-	+	-	+	+	-	-
<i>Asmicridea</i> sp. 5 (L)	+	+	-	-	+	-	+	-	-	-
<i>Baliomorpha banksi</i> (Mosely)	+	+	+	+	+	-	+	-	+	+
<i>Cheumatopsyche</i> sp. nov. 11 (L)	-	+	-	-	-	-	-	-	-	-
<i>Cheumatopsyche</i> sp. nov. 12 (L)	+	-	+	-	-	-	+	-	-	-
<i>Cheumatopsyche</i> sp. nov. 13 (L)	+	+	-	-	-	-	-	-	+	-
<i>Cheumatopsyche</i> sp. nov. 14 (L)	-	-	-	+	+	-	+	+	+	+
<i>Cheumatopsyche</i> sp. nov. 15 (L)	+	+	+	+	+	-	+	+	-	+
<i>Cheumatopsyche</i> sp. nov. 16 (L)	+	-	+	+	+	-	+	+	+	-
<i>Cheumatopsyche</i> sp. nov. 17 (L)	-	-	-	-	+	-	-	-	-	-
<i>Cheumatopsyche</i> sp. nov. 19 (L)	-	+	-	-	-	-	-	+	-	-
<i>Cheumatopsyche</i> sp. nov. 22 (L)	-	-	-	-	-	-	+	-	+	-
<i>Cheumatopsyche</i> spp. indet.	+	+	+	+	+	+	+	+	+	+
<i>Diplectronea</i> sp. nov. 7 (L)	-	-	+	+	-	-	+	+	+	-
<i>Diplectronea</i> sp. nov. 8 (L)	-	-	-	+	-	-	-	-	+	+
<i>Diplectronea</i> sp. nov. 10 (L)	-	-	-	-	+	-	+	-	-	-
<i>Diplectronea</i> sp. nov. PT-999	-	-	-	-	-	+	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-1000	-	-	+	-	+	+	+	-	-	-
<i>Diplectronea</i> sp. nov. PT-1002	-	-	-	-	-	+	-	+	+	-
<i>Diplectronea</i> sp. nov. PT-1003	-	-	+	-	-	-	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-1012	-	-	-	-	-	+	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-1016	-	+	-	-	-	-	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-1031	-	-	-	-	-	+	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-1040	-	-	+	-	+	-	+	-	-	-
<i>Diplectronea</i> sp. nov. PT-2007	-	-	+	-	-	-	-	-	-	-
<i>Diplectronea</i> sp. nov. PT-2042	+	-	-	-	-	-	-	-	-	-
<i>Macrostemum saundersii</i> (McLachlan)	+	-	-	-	-	-	-	-	-	-
<i>Smicrophylax</i> sp. 5 (L)	+	-	+	-	-	-	-	+	+	-
<i>Smicrophylax</i> sp. 6 (L)	-	-	+	-	-	-	-	-	-	-
<i>Smicrophylax ulmeri</i> (Banks)	-	-	+	+	+	+	+	+	+	+
<i>Smicrophylax</i> spp. indet.	+	+	+	+	+	-	+	+	+	-

Table 1 (cont.). Trichoptera species checklist for the Queensland Wet Tropics, with presence or absence within each investigation area noted. (Note: L - denotes taxa recorded as larvae only.)

INVESTIGATION AREAS	1	2	3	4	5	6	7	8	9	10
Hydroptilidae										
<i>Acanthotrichia bilamina</i> Wells	-	-	-	-	-	-	-	-	+	+
<i>Acritoptila capistra</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Acritoptila pearsoni</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Acritoptila</i> sp. indet.	-	-	-	-	-	-	-	-	+	-
<i>Chrysotrichia australis</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Gnathotrichia australiensis</i> Wells	-	-	-	-	-	-	-	-	-	+
<i>Hellyethira cornuta</i> Wells	+	+	-	-	+	-	+	-	+	+
<i>Hellyethira cubitans</i> Wells	-	-	-	-	+	-	-	-	-	+
<i>Hellyethira eskensis</i> (Mosely)	-	-	-	-	+	-	+	-	-	+
<i>Hellyethira imparalobata</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Hellyethira quadrata</i> Wells	-	-	-	-	-	-	-	-	+	+
<i>Hellyethira simplex</i> Mosely	-	-	-	-	+	-	+	-	+	+
<i>Hellyethira</i> sp. nov.	-	-	-	-	-	-	-	-	-	+
<i>Hellyethira</i> sp. nov. A	-	-	-	-	-	-	-	-	-	+
<i>Hellyethira spinosa</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Hellyethira vernoni</i> Wells	-	-	-	-	+	-	-	+	+	+
<i>Hellyethira</i> spp. indet.	+	-	-	-	+	-	+	-	+	+
<i>Hydroptila incertula</i> Mosely	-	-	-	-	-	-	-	-	+	+
<i>Hydroptila losida</i> Mosely	-	-	-	-	-	-	-	-	+	+
<i>Hydroptila obscura</i> Wells	-	-	-	-	+	-	+	+	+	+
<i>Hydroptila scamandra</i> Neboiss	-	-	-	-	-	-	+	+	+	-
<i>Hydroptila</i> spp. indet.	+	-	-	-	-	-	-	-	-	+
<i>Maydenoptila kurandica</i> Wells	-	-	-	+	-	-	-	-	-	-
<i>Mulgravia coronata</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Orphnino-trichia silicis</i> Wells	-	-	-	-	-	-	+	-	-	-
<i>Orthotrichia bensoni</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Orthotrichia bullata</i> Wells	-	-	-	-	+	-	-	-	-	+
<i>Orthotrichia conferta</i> Wells	-	-	-	-	+	+	-	-	+	-
<i>Orthotrichia constricta</i> Wells	-	-	-	-	-	-	-	-	+	-
<i>Orthotrichia divaricata</i> Wells	-	-	-	-	+	-	-	-	+	-
<i>Orthotrichia morula</i> Wells	-	-	-	-	+	-	+	-	-	-
<i>Orthotrichia</i> sp. nov. A (<i>aberrans</i> group)	-	-	-	-	-	-	-	-	-	+
<i>Orthotrichia turrita</i> Wells	-	-	-	-	-	-	-	-	-	+
<i>Orthotrichia velata</i> Wells	-	-	-	-	-	-	-	-	+	+
<i>Orthotrichia</i> spp. indet.	+	-	-	+	-	-	+	-	+	+
<i>Oxyethira bogambara</i> Schmid	-	-	-	-	-	-	-	-	+	-
<i>Oxyethira columba</i> (Neboiss)	-	-	-	-	-	-	-	-	+	+
<i>Oxyethira complicata</i> Wells	-	-	-	-	-	-	-	-	+	+
<i>Oxyethira</i> spp. indet.	+	-	-	-	-	-	-	+	+	+
<i>Oxyethira triangulata</i> Wells	-	-	-	-	+	-	+	+	+	+
<i>Stenoxyethira plumosa</i> Wells	-	-	-	-	+	-	-	-	-	-

Table 1 (cont.). Trichoptera species checklist for the Queensland Wet Tropics, with presence or absence within each investigation area noted. (Note: L - denotes taxa recorded as larvae only.)

INVESTIGATION AREAS	1	2	3	4	5	6	7	8	9	10
Hydroptilidae (cont.)										
<i>Stenoxyethira</i> sp. indet.	-	-	-	-	-	-	-	+	-	-
<i>Tricholeiochiton fidelis</i> Wells	-	-	-	-	-	-	-	-	-	+
<i>Tricholeiochiton</i> sp. indet.	-	-	-	-	-	-	-	-	-	+
<i>Xuthotrichia ochracea</i> Mosely	-	-	-	-	-	-	-	+	-	-
Leptoceridae										
Gen. indet. sp.	-	-	+	-	+	-	+	-	-	-
<i>Lectrides varians</i> Mosely	-	-	+	-	-	-	-	+	-	+
<i>Leptocerus assimulans</i> (Ulmer)	+	-	+	-	-	-	-	-	-	-
<i>Leptorussa darlingtoni</i> (Banks)	-	-	+	-	-	-	-	-	-	-
<i>Notalina</i> spp. indet.	+	+	+	-	+	-	+	+	+	+
<i>Notoperata</i> sp. nov. PT-2014	+	+	+	-	-	-	+	+	+	+
<i>Oecetis australis</i> (Banks)	-	-	-	+	+	-	+	-	-	+
<i>Oecetis burtoni</i> Neboiss	+	-	+	-	-	-	-	-	-	+
<i>Oecetis complexa</i> Kimmins	-	-	-	-	+	-	+	-	-	+
<i>Oecetis epekeina</i> Neboiss	-	-	-	-	-	-	+	-	-	-
<i>Oecetis inscripta</i> Kimmins	-	-	-	-	+	-	-	-	-	-
<i>Oecetis laustra</i> Mosely	-	-	+	-	+	-	+	-	+	+
<i>Oecetis multipunctata</i> Ulmer	-	-	+	+	+	-	+	+	-	-
<i>Oecetis oresbiosa</i> Neboiss	+	-	-	+	+	-	-	-	+	-
<i>Oecetis piptona</i> Neboiss	-	-	-	-	+	-	+	-	-	-
<i>Oecetis</i> spp. indet.	+	+	+	+	+	+	+	+	+	+
<i>Setodes bracteatus</i> Neboiss	-	+	-	+	+	-	+	+	+	+
<i>Symphitoneuria exigua</i> (McLachlan)	-	-	+	-	-	-	+	-	+	-
<i>Triaenodes</i> sp.	-	-	-	-	-	-	-	-	-	+
<i>Triaenodes</i> sp. nov. PT-659	-	-	-	-	+	-	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-756	-	-	-	-	-	-	-	-	-	+
<i>Triaenodes</i> sp. nov. PT-757	-	-	-	-	-	-	+	-	-	-
<i>Triaenodes</i> sp. nov. PT-759	-	-	-	-	+	-	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-761	-	-	-	-	-	-	+	-	-	-
<i>Triaenodes</i> sp. nov. PT-763	+	-	+	-	-	-	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-764	+	-	+	-	+	+	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-767	+	-	-	-	+	+	-	-	+	+
<i>Triaenodes</i> sp. nov. PT-783	+	-	-	+	+	-	+	-	-	+
<i>Triaenodes</i> sp. nov. PT-800	-	-	+	-	-	-	-	-	-	+
<i>Triaenodes</i> sp. nov. PT-1090	-	-	-	-	-	+	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-1091	-	-	-	-	-	+	-	-	-	-
<i>Triaenodes</i> sp. nov. PT-1116	-	-	-	-	-	-	-	-	+	+
<i>Triaenodes</i> sp. nov. PT-1117	-	-	-	-	-	-	-	+	-	+
<i>Triaenodes</i> sp. nov. PT-1122	-	-	-	-	-	-	+	-	-	-
<i>Triaenodes</i> sp. nov. PT-2021	-	-	-	-	-	-	-	-	-	+
<i>Triaenodes</i> sp. nov. PT-2040	-	-	-	-	-	-	+	-	-	-

Table 1 (cont.). Trichoptera species checklist for the Queensland Wet Tropics, with presence or absence within each investigation area noted. (Note: L - denotes taxa recorded as larvae only.)

INVESTIGATION AREAS	1	2	3	4	5	6	7	8	9	10
Leptoceridae (cont.).										
<i>Triaenodes</i> spp. indet.	+	-	+	-	+	-	+	-	+	-
<i>Triaenodes volda</i> Mosely	-	+	+	-	+	-	+	-	+	+
<i>Triplectides australicus</i> Banks	-	-	-	-	+	-	-	-	-	-
<i>Triplectides australis</i> Navas	-	-	-	-	-	-	+	-	+	-
<i>Triplectides ciuskus</i> Mosely	-	-	-	-	+	-	+	-	-	+
<i>Triplectides dolabratusa</i> Morse & Neboiss	-	-	+	-	-	+	-	+	+	-
<i>Triplectides elongatus</i> Banks	-	-	+	-	+	-	-	-	-	-
<i>Triplectides gonetalus</i> Morse & Neboiss	-	+	+	-	+	-	+	+	+	+
<i>Triplectides hamatus</i> Morse & Neboiss	-	-	-	-	-	-	-	-	-	+
<i>Triplectides helvolus</i> Morse & Neboiss	-	-	-	-	-	-	+	-	-	-
<i>Triplectides liratellus</i> Morse & Neboiss	-	-	-	-	+	-	-	-	-	-
<i>Triplectides liratus</i> Morse & Neboiss	+	+	-	-	-	+	+	-	+	+
<i>Triplectides parvus</i> (Banks)	+	-	-	+	+	-	+	-	+	+
<i>Triplectides prolatus</i> Morse & Neboiss	-	-	+	+	+	-	+	-	-	-
<i>Triplectides rossi</i> Morse & Neboiss	-	-	-	-	-	+	+	-	+	-
<i>Triplectides similis</i> Mosely	-	-	-	-	-	-	+	-	-	-
<i>Triplectides</i> spp. indet.	+	+	+	+	+	+	+	+	+	+
<i>Triplectides tambina</i> Mosely	-	-	-	-	-	-	-	-	-	+
<i>Triplexa</i> sp. nov. PT-1762	-	-	-	+	-	-	-	-	+	-
<i>Westriplectes angelae</i> Neboiss	-	-	-	-	+	-	-	-	-	-
Odontoceridae										
<i>Barynema</i> sp. nov. PT-1176	+	-	-	-	-	-	-	-	-	-
<i>Barynema</i> sp. nov. PT-1405	-	-	+	-	+	-	+	+	+	-
<i>Barynema</i> sp. nov. PT-2028	-	-	-	-	-	-	+	-	-	-
<i>Marilia bola</i> Mosely	+	-	+	-	-	-	+	-	+	-
<i>Marilia fusca</i> Kimmins (L)	-	-	-	-	-	-	+	-	-	-
<i>Marilia</i> spp. indet.	+	+	+	+	+	+	+	+	+	+
Philopotamidae										
<i>Chimarra australica</i> (Ulmer)	-	-	+	+	+	+	+	-	+	-
<i>Chimarra monticola</i> Kimmins	+	-	+	-	+	-	+	-	+	-
<i>Chimarra</i> sp. nov. 6 (L)	-	-	-	-	+	-	-	-	-	-
<i>Chimarra</i> sp. nov. 7 (L)	+	-	+	-	+	-	+	+	+	-
<i>Chimarra</i> sp. nov. 11 (L)	-	-	-	-	-	-	-	-	+	-
<i>Chimarra</i> sp. nov. CT-221	-	+	-	-	+	+	-	+	-	+
<i>Chimarra</i> sp. nov. CT-223	+	+	-	-	+	-	+	+	+	-
<i>Chimarra</i> sp. nov. CT-225	-	-	+	-	+	-	+	-	+	-
<i>Chimarra</i> sp. nov. CT-226	-	-	-	-	-	-	-	-	-	+
<i>Chimarra</i> sp. nov. CT-227	-	-	-	-	-	+	-	-	-	-
<i>Chimarra</i> sp. nov. CT-228	-	-	-	-	-	+	-	-	-	-
<i>Chimarra</i> spp. indet.	+	-	+	-	+	+	-	+	+	+
<i>Chimarra uranka</i> Mosely	+	+	+	+	+	-	+	+	+	+

Table 1 (cont.). Trichoptera species checklist for the Queensland Wet Tropics, with presence or absence within each investigation area noted. (Note: L - denotes taxa recorded as larvae only.)

INVESTIGATION AREAS	1	2	3	4	5	6	7	8	9	10
Philopotamidae (cont.)										
Gen. nov. sp. nov. PT-1640	-	-	-	-	+	-	-	-	+	-
<i>Hydrobiosella</i> sp. nov. 15 (L)	-	-	+	+	+	-	+	-	+	+
<i>Hydrobiosella</i> sp. nov. PT-1037	-	-	-	-	-	+	-	-	-	-
<i>Hydrobiosella</i> sp. nov. PT-1038	-	-	-	-	-	+	-	-	-	-
<i>Hydrobiosella</i> sp. nov. PT-1039	-	-	-	-	-	+	-	-	-	-
<i>Hydrobiosella</i> sp. nov. PT-1768	-	-	-	+	+	-	+	+	-	+
<i>Hydrobiosella</i> sp. nov. PT-2029	-	-	-	-	-	-	-	-	-	+
<i>Hydrobiosella</i> spp. indet.	-	-	-	-	-	+	+	-	+	+
Philorheithridae										
<i>Aphilorheithrus</i> sp. nov. PT-2038	-	-	+	-	-	-	-	+	+	+
Gen. Nov. P sp. nov. PT-1707	-	-	-	-	-	+	-	-	-	-
Gen. Nov. Q sp. nov. PT-1837	-	-	+	-	+	-	-	+	+	+
Polycentropodidae										
Gen. G sp. 1 (L)	-	-	-	-	-	-	+	-	-	-
<i>Paranyctiophylax</i> sp. nov. 5 (L)	-	-	-	-	-	-	-	-	+	-
<i>Paranyctiophylax</i> sp. nov. 7 (L)	-	-	-	-	-	-	-	-	+	-
<i>Paranyctiophylax</i> sp. nov. PT-1589	-	-	-	-	-	-	-	-	+	+
<i>Paranyctiophylax</i> sp. nov. PT-1625	-	-	-	-	-	-	-	-	+	-
<i>Paranyctiophylax</i> sp. nov. PT-1977	+	-	+	-	+	-	+	-	-	-
<i>Paranyctiophylax</i> sp. nov. PT-1979	+	-	+	-	+	-	+	-	+	-
<i>Plectrocnemia</i> sp. nov. PT-1817	+	-	+	-	+	-	+	-	+	+
<i>Plectrocnemia</i> sp. nov. PT-1822	-	-	+	-	-	+	+	-	+	+
<i>Plectrocnemia</i> sp. nov. PT-1976	-	-	-	-	-	-	-	-	+	-
<i>Plectrocnemia</i> spp. indet.	+	-	+	-	-	-	+	-	+	+
<i>Polyplectropus</i> sp. 2 (L)	-	-	+	-	-	-	+	-	+	-
<i>Polyplectropus</i> sp. 3 (L)	-	-	-	-	-	-	-	-	+	+
<i>Polyplectropus</i> sp. nov. PT-1821	-	-	+	+	+	+	+	+	+	-
Psychomyiidae										
<i>Tinodes radona</i> Neboiss	+	-	-	-	-	-	-	-	-	-
<i>Zelandoptila yuccabina</i> Neboiss	-	-	-	-	-	-	-	-	+	-
Stenopsychidae										
<i>Stenopsychodes mjobergi</i> Ulmer	+	-	+	+	+	+	+	+	-	+
<i>Stenopsychodes</i> sp. nov. A	-	-	-	-	-	+	-	-	+	-
<i>Stenopsychodes</i> sp. nov. B	-	-	-	-	-	+	-	-	+	-
<i>Stenopsychodes</i> sp. nov. C	-	-	+	-	+	+	+	+	+	-
<i>Stenopsychodes</i> sp. nov. D	-	-	-	-	-	+	-	-	-	-
<i>Stenopsychodes</i> sp. nov. E	-	-	-	-	-	+	-	-	-	-
<i>Stenopsychodes</i> sp. nov. F	-	-	-	-	-	+	-	-	-	-
<i>Stenopsychodes</i> spp. indet.	+	+	+	-	+	+	+	+	+	+
Tasimiidae										
<i>Tasiagma</i> sp.	-	-	-	-	-	-	+	+	-	-

A taxonomic summary of the QWT Trichoptera fauna is presented in Table 2. In total 217 species have been recognised, of which 95 species and four genera are undescribed. The highest number of species recorded at an individual site was 78 at Yuccabine Creek, while the 10 most speciose sites yielded an average of 41.8 species.

Table 2. Taxonomic evaluation of the Queensland Wet Tropics Area Trichopteran fauna.

Total number of species:	217
Taxa identified from adult material:	202
Taxa identified from immature material:	79
Immature taxa clearly different from adult taxa:	15
Immature taxa associated with adults:	30
Immature taxa not associated with adults:	49
New species recognised:	95
New genera recognised:	4
New species based on adult material only:	83
New species based on immature material only:	12

The PATN analyses were performed separately on the adult and immature data (for full details see Walker *et al.* 1993). As expected, the larger adult data set provided better resolution between the investigation areas, although results from immatures supported those obtained for adults. The analyses separated the investigation areas into three groups, each with a distinctive faunal composition. Investigation areas 9 and 10 were grouped, investigation area 6 was unique and there were no significant faunal differences between the remaining seven investigation areas.

Discussion

Comparison of the trichopteran faunas of Australia, the Queensland Wet Tropics and the Tasmanian World Heritage Area (Table 3) indicates: higher species richness in the QWT, with 36.7% of the known Australian fauna, compared to 22.7% occurring in the TWHA; QWT has a higher maximum number of species occurring at an individual site and within a single Investigation Area than has the TWHA; the QWT has a similar number of families and genera as the TWHA, though the family composition differs between the two areas. Families present in the QWT and not found in the TWHA are Antipodoeciidae, Dipseudopsidae (formerly Hyalopsychidae: Wells and Cartwright, 1993a), Odontoceridae, Psychomyiidae, Calamoceratidae and Stenopsychidae. Families present in the TWHA and not found in the QWT are Kokiriidae, Limnephilidae, Oeconesidae and Plectrotarsidae.

Table 3. Comparison of the trichopteran faunas of Australia (AUS), the Queensland Wet Tropics (QWT) and the Tasmanian World Heritage Area (TWHA) (Note: In calculating the numbers of species, immature and 'spp. indet.' taxa were excluded if it was likely that they were conspecific with listed adult species; undescribed species and genera, recorded from the QWT, have been included in the total number of known Australian taxa. Sources of species composition: AUS-Neboiss, 1991, 1992; Wells, 1990; TWHA- Neboiss, Jackson & Walker, 1989).

	AUS	QWT	TWHA
Number of known families:	26	21	19
Number of known genera:	106	67	62
Number of known species:	590	217	134
Highest number of species within an Investigation Area:	-	100	91
Highest number of species at a single site:	-	78	45
Number of species in the 10 most specious Australian families:			
Hydroptilidae	121	39	10
Leptoceridae	102	51	28
Ecnomidae	67	24	5
Hydrobiosidae	58	7	26
Hydropsychidae	46	26	5
Philopotamidae	32	16	7
Conoesucidae	23	2	15
Calocidae	19	8	5
Philorheithridae	16	3	8
Calamoceratidae	10	5	0

The species richness within investigation areas of the QWT is remarkably high, with seven investigation areas yielding 50 or more species. Species richness observed at many sites was greater than recorded in southern Australia from the Tasmanian World Heritage Area (Neboiss *et al.* 1989). The average number of species for the ten most diverse sites in the two World Heritage Areas was 41.8 species for the QWT and 37.0 species for the TWHA. The highest number of species recorded at an individual site within the QWT was at Yuccabine Creek (78 species). This site exceeded the richest TWHA site, which yielded 45 species (Franklin River, Roaring Creek Junction) and also the 44 species recorded from the O'Shannassy River in Victoria (Dean and Cartwright 1987) and the 47 species recorded from Gunshot Creek, Cape York Peninsula (Wells and Cartwright 1993b). It should be noted that the species lists for Yuccabine Creek (Benson and Pearson 1988) and O'Shannassy River were the result of extensive collecting programs; lower species numbers at other sites may merely reflect a lesser

collecting effort. There is therefore evidence that Trichoptera species richness is probably greater in the QWT than in other areas of Australia.

While it is difficult to quantify the definition of a rare species, a qualitative assessment of the conservation status of taxa within the QWT has been attempted. For the purposes of this study, we have developed the following definitions:

- a species is defined as "rare" when known from less than four specimens;
- a species is defined as "localised" when known only from a single site;
- a species is defined as "vulnerable" when known from a single site and less than four specimens.

Using the above definitions, the conservation status of the Trichoptera fauna of the QWT is as follows:

- 52.1% (113 species) of the QWT Trichoptera fauna is "rare";
- 30.0% (65 species) of the QWT Trichoptera fauna is "localised";
- 22.1% (48 species) of the QWT Trichoptera fauna is "vulnerable";
- 77.1% (37 species) of the "vulnerable" fauna is presumed to be endemic to the QWT based on current knowledge.

Of the 113 species defined as rare, 54 species have been recorded from one or more of the three sites, Yuccabine Creek, Bellenden Ker Range and Birthday Creek Falls. Furthermore, these three sites have yielded 27 of 65 species known from single sites only and 18 of 48 species defined as vulnerable. On the basis of current information, these sites are worthy of special conservation consideration.

Although the Australian Trichoptera fauna is reasonably well known, it is significant that 43.8% of the species recorded from the QWTA are new to science. This demonstrates the value of involving taxonomic experts at an early stage of any faunal survey. Table 2 also highlights the disparity between knowledge of the adult and immature stages. While 202 of 217 recognised species (93.1%) were based on adult specimens, only 15 of 217 recognised species (6.9%) were based solely on immature specimens. Furthermore, of the 202 species recognised from adult material, only 30 species (14.9%) were associated with larvae. In fact, the larvae of more than half the total known species from the QWTA have not yet been collected, let alone associated with adults.

Acknowledgments

The authors wish to acknowledge the following persons or institutions for their assistance. We are grateful to the Queensland Museum, Australian Museum and the Australian National Insect Collection, CSIRO for the loan and use of material from their respective collections. We sincerely thank: Dr Alice Wells for identification of Hydroptilidae and *Oecetis* specimens as well as critical comments on the manuscript, Ms Toulia Marra for database input and Ms Heather Martin and Ms Catriona McPhee for comments on earlier

drafts of the manuscript. Finally, we thank the Wet Tropics Management Agency for their financial assistance which allowed this study to be completed.

References

- BELBIN, L. 1988. *PATN, pattern analysis package*. CSIRO, Division of Wildlife and Rangelands Research, Canberra.
- BENSON, L. and PEARSON 1988. Diversity and seasonality of adult Trichoptera captured in a light-trap at Yuccabine Creek, a tropical Australian rainforest stream. *Australian Journal of Ecology* **13**: 337-344.
- DEAN, J. and CARTWRIGHT, D. 1987. Trichoptera of a Victorian forest stream: species composition and life histories. *Australian Journal of Marine and Freshwater Research* **38**: 845-860.
- NEBOISS, A. 1991. Trichoptera (caddis-flies, caddises). Pp 787-816 in CSIRO (ed.). *The insects of Australia*. Melbourne University Press.
- NEBOISS, A. 1992. Illustrated keys of the families and genera of Australian Trichoptera. 1. Adults. *Australian Society for Limnology Special Publication* No. 9.
- NEBOISS, A., JACKSON, J. and WALKER, K. 1989. Caddis-flies (Insecta: Trichoptera) of the World Heritage Area in Tasmania - species composition and distribution. *Occasional Papers of the Museum of Victoria* **4**: 1-41.
- WALKER, K.L., NEBOISS, J., DEAN, J. and CARTWRIGHT, D. 1993. A preliminary investigation of the caddis-flies (Trichoptera: Insecta) of the Queensland Wet Tropics World Heritage Area. Museum of Victoria.
- WELLS, A. 1990 New species and a new genus of micro-caddisfly from northern Australia, including the first Australian record of the tribe Stactobiini (Trichoptera: Hydroptilidae). *Transactions of the Royal Society of South Australia* **114**: 107-128.
- WELLS, A. and CARTWRIGHT, D. 1993a. Females and immatures of the Australian caddisfly *Hyalopsyche disjuncta* Neboiss (Trichoptera), and a new family placement. *Transactions of the Royal Society of South Australia* **117**: 97-104.
- WELLS, A. and CARTWRIGHT, D. 1993b. Trichoptera, Ephemeroptera, Plecoptera and Odonata of the Jardine River area, Cape York Peninsula, Northern Queensland, pp. 221-230 in Cape York Peninsula Scientific Expedition 1992, The Royal Geographical Society of Queensland Inc.



Walker, Ken et al. 1995. "A preliminary investigation of the caddis-flies (Insecta: Trichoptera) of the Queensland Wet Tropics." *The Australian Entomologist* 22(1), 19–31.

View This Item Online: <https://www.biodiversitylibrary.org/item/307390>

Permalink: <https://www.biodiversitylibrary.org/partpdf/344248>

Holding Institution

Entomological Society of Queensland

Sponsored by

Atlas of Living Australia

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: Entomological Society of Queensland

License: <http://creativecommons.org/licenses/by-nc-sa/4.0/>

Rights: <http://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.