

rays of the sun played upon the purplish feathers and the beautiful sapphire-blue eyes flashed. The bird kept spreading its tail, drooping its wings, and mimicking all birds common to the district—Lyre-Bird included. Then suddenly it would break into the purring note, and a “green” bird with a strawberry in its bill would join it. Upon the male bird taking the gift the other would fly off. Suddenly several green-plumed birds made their appearance, and one, settling close to me, gave a warning note, when they all disappeared in the scrub.

Satin Bower-Birds are very fond of fruit, early morning and evening being their favourite times for raiding orchards. The owner of the orchard mentioned informed me that the birds were a great nuisance, but it was a very rare thing to see a black-plumed bird out feeding in the open with the “green” ones, and that the “green” birds kept carrying fruit into the scrub. Other orchardists have told me the same thing.

I have heard it stated that the male Satin Bower-Bird does not live long after it attains the dark plumage. I have kept a caged bird five years after the change. It then accidentally gained its liberty, and, after staying about the house for a few weeks, disappeared.

### List of Birds Observed on Parry's Creek, North-West Australia.

BY GREGORY M. MATHEWS, F.L.S., &c.

FROM notes supplied to me by Mr. Rogers I make out the following list, showing the birds' movements during February, March, and April, 1909.

It will be noticed that some of the birds mentioned in my two lists\* of the birds of that district for 1908 have left the locality.

The numbers are from my “Handlist.” Those marked thus \* are rare.

No.					Feb.	Mar.	Ap.
13	Synœcus cervinus	..	..	..			
15	Turnix maculosa	..	..	..	“	“	“
20	„ pyrrhothorax	..	..	..	“	“	“
21	„ velox	..	..	..	“	“	“
33	Geopelia humeralis	..	..	..	“	“	“
34	„ placida	..	..	..	“	“	“
35	„ cuneata	..	..	..	“	“	“
37	Phaps chalcoptera	..	..	..	“	“	“
40	Petrophassa albipennis	..	..	..		1 bird in	Ap.
45	Lophophaps plumifera	..	..	..	Feb.	Mar.	Ap.
56	Poliolimnas cinereus	..	..	..	“	“	—
59	Microtribonyx ventralis	..	..	..	“	“	Ap.
62	Porphyrio melanotus	..	..	..	“	“	—
65	Podicipes novæ-hollandiæ	..	..	..	“	“	“
119	Hydrochelidon hybrida	..	..	..	“	“	“

\* Ante, in vol. ix., p. 1 and p. 53 respectively.

				Feb.	—	—
				"	Mar.	Ap.
137	Larus novæ-hollandiæ	..	..	..	—	—
146	Erythrogonyx cinctus	..	..	..	—	—
148	Lobivanellus miles	..	..	..	—	—
158	Ægialitis melanops	..	..	..	—	—
161	Himantopus leucocephalus	..	..	..	—	—
166	Mesoscolopax minutus	..	..	..	—	—
168	Limosa melanuroides	..	..	..	—	—
173	Tringoides hypoleucus	..	..	..	Mar.	—
175	Glottis nebularius	..	..	..	—	Ap.
176	Rhyacophilus glareola	..	..	..	—	—
179	Pisobia ruficollis	..	..	..	—	—
181	Heteropygia aurita	..	..	..	—	Ap.
185	Gallinago australis	..	..	..	—	—
187	Hydralector gallinaceous	..	..	..	—	—
188	Stiltia isabella	..	..	..	—	Ap.
189	Glareola orientalis	..	..	..	—	Ap.
190	Burhinus grallarius	..	..	..	—	—
192	Eupodotis australis	..	..	..	—	—
193	Antigone australasiana	..	..	..	—	—
194	Ibis molucca	..	..	..	—	—
195	Carpibis spinicollis	..	..	..	—	—
196	Plegadis falcinellus	..	..	..	—	—
197	Platalea regia	..	..	..	—	—
198	Platibis flavipes	..	..	..	—	—
199	Xenorhynchus asiaticus	..	..	..	—	Ap.
202	Mesophoyx plumifera	..	..	..	—	—
203	Herodias timoriensis	..	..	..	—	—
204	Notophoyx novæ-hollandiæ	..	..	..	—	—
205	„ pacifica	..	..	..	—	—
206	„ flavirostris	..	..	..	—	—
210	Nycticorax caledonicus	..	..	..	—	—
214	Dupetor gouldi	..	..	..	—	—
217	Anseranas semipalmata	..	..	..	Mar.	Ap.
218	Nettopus pulchellus	..	..	..	—	—
222	Dendrocygna arcuata	..	..	..	—	—
223	„ eytoni	..	..	..	—	—
224	Tadorna rufitergum	..	..	..	—	—
226	Anas superciliosa	..	..	..	—	—
228	Nettiump gibberifrons	..	..	..	—	—
234	Aythya australis	..	..	..	—	—
238	Phalacrocorax sulcirostris	..	..	..	—	—
241	„ melanoleucus	..	..	..	—	—
242	Plotus novæ-hollandiæ	..	..	..	—	—
252	Pelecanus conspicillatus	..	..	..	—	—
253	Circus assimilis	..	..	..	—	—
254	„ gouldi	..	..	..	—	—
256	Astur novæ-hollandiæ	..	..	..	—	—
258	„ fasciatus	..	..	..	—	—
262	Uroaëtus audax	..	..	..	—	—
265	Haliaëtus leucogaster	..	..	..	—	—
266	Haliastur girrenera	..	..	..	I bird in Feb.	—
267	„ sphenurus	..	..	..	Feb.	Mar. Ap.
268	Milvus affinis	..	Rare in Feb.	Common in Mar. and Ap.	—	—
271	Elanus axillaris.	Only 1 seen in each month	—	—	Feb.	Mar. Ap.
277	Falco lunulatus	..	..	..	*	—

278	Hieracidea berigora	..	..	Feb.	Mar.	Ap.
279	" orientalis	..	..	"	"	"
280	Cerchneis cenchroides	..	..	—	"	"
285	Ninox ocellata	..	..	Feb.	„	—
289	" occidentalis	..	..	"	"	Ap.
303	Trichoglossus rubritorques	..	..	"	"	"
306	Ptilosclera versicolor	..	..	"	"	—
316	Calyptorhynchus macrorhynchus	..	..	"	"	Ap.
320	Cacatua galerita	..	..	"	"	"
322	" gymnopis	..	..	"	"	"
324	" roseicapilla	..	..	"	"*	"*
327	Calopsittacus novæ-hollandiæ	..	..	—	"*	"*
331A	Ptistes coccineopterus	..	..	Feb.	„	"
372	Melopsittacus undulatus	..	..	—	"*	"
377	Podargus phalænoides	..	..	—	1 bird in Ap.	
381	Eurystomus pacificus	..	..	Feb.*	Mar.*	"
383	Alcyone pulchra	..	..	—	—	"
388	Dacelo cervina	..	..	Feb.	Mar.	"
390	Halcyon pyrrhopygius	..	..	"	"	"
391	" sanctus	..	..	"	"	"
396	Merops ornatus	..	..	"	"	"
398	Eurostopus argus	..	..	"*	"*	"
403	Cypselus pacificus	..	..	—	—	"
404	Cuculus saturatus	..	..	Feb.*	Mar.*	"
405	" inornatus	..	..	"	"	"
406	Cacomantis rufulus	..	..	"	—	—
409	Mesocalius palliolatus	..	..	"	Mar.	—
410	Chalcococcyx basalis	..	..	"	"	—
415	Eudynamis cyanocephala	..	..	"	"	—
417	Centropus phasianus	..	..	"	"	Ap.
432	Petrochelidon ariel	..	..	"	"	"
434	Microeca assimilis	..	..	"	"	"
450	Smicrornis flavesiensis	..	..	"	"	"
452	Gerygone cinerascens	..	..	"	"	"
469	Pœciliodryas cerviniventris	..	..	"	"	"
486	Rhipidura isura	..	..	"	"	"
487	" tricolor	..	..	"	"	"
489	Myiagra concinna	..	..	"	"	"
494	Sisura nana	..	..	"	"	"
504	Coracina robusta	..	..	"	"*	"
506	" hypoleuca	..	..	"*	"*	"
510	Lalage tricolor	..	..	"	"	"
532	Pomatostomus rubeculus	..	..	"	"	"
538	Cincloramphus cruralis	..	..	"*	"*	"
539	" rufescens	..	..	"	"	"
547	Ephthianura tricolor	..	..	—	"	"
552	Cisticola exilis	..	..	Feb.	"	"
603	Malurus assimilis	..	..	"*	"*	"
608	" cruentatus	..	..	"	"	"
624	Artamus leucogaster	..	..	"	"	—
629	" cinereus	..	..	"	"	Ap.
635	" minor	..	..	"*	"*	"
638	Collyriocichla brunnea	..	..	"	"	"
646	Grallina picata	..	..	"*	"*	"
655	Cracticus picatus	..	..	"*	"*	"

672	Pachycephala falcata	..	..	..	Feb.	Mar.	Ap.
699	Neositta leucoptera	..	..	..	"	"	"
701	Climacteris melaṇura	..	..	—	—	—	*
722	Dicæum hirundinaceum	..	..	..	Feb.	Mar.	Ap.*
728	Pardalotus rubricatus	..	..	..	"	"	*
730	" uropygialis	..	..	..	"	"	*
734	Melithreptus albicularis	..	..	..	"*	"*	*
738	" lætior	..	..	..	"*	"*	*
758	Glycyphila fasciata	..	..	—		very rare	
762	Conopophila rufigularis	..	..	..	Feb.	Mar.	Ap.
765	Stigmatops ocularis	..	..	..	"	"	*
772	Ptilotis sonora	..	..	..	"	"	*
788	" flavescens	..	..	—	"	"	*
795	" unicolor	..	..	..	Feb.	"	"
815	Entomyza albipennis	..	..	..	"	"	*
816	Tropidorhynchus argenticeps	..	..	—	—	—	*
819	Philemon citreigularis	..	..	..	Feb.	Mar.	,
822	Anthus australis	..	..	—	—	—	*
825	Mirafra milligani	..	..	..	Feb.	Mar.	,
831	Tæniopygia castanotis	..	..	..	"	"	"
833	Stictoptera annulosa	..	..	..	"	"	*
835	Munia flavigrymnæ	..	..	—	—	—	*
836	" pectoralis	..	..	..	Feb.*	Mar.*	,
839	Bathilda ruficauda	..	..	..	"*	—	,
841	Poephila acuticauda	..	..	..	"	Mar.	,
844	" personata	..	..	..	"	"	"
846	" gouldiae	..	..	..	"	"	"
849	Neochmia phaeton	..	..	..	"	"	,
850	Mimeta sagittata	..	..	..	"	I only	—
851	" flavicinctus	..	..	..	"*	Mar.*	Ap.*
863	Chlamydodera nuchalis	..	..	..	"	"	"
872	Corvus coronoides	..	..	..	"	"	"

## Birds about Drouin, Gippsland.

BY ISAAC BATEY.

WEDGE-TAILED EAGLE (*Uroaetus audax*).—Very rarely noted.

GOSHAWK (*Astur approximans*).—One seen.

BROWN HAWK (*Hieracidea orientalis*).—A single bird.

NANKEEN KESTREL (*Cerchneis cenchroides*).—Common at times, and supposed to breed hereabouts.

HARRIER (*Circus gouldi*).—Annual visitant, and breeds.

BLACK-SHOULDERED KITE (*Elanus axillaris*).—Once some appeared, and were quite new to residents in my vicinage.

BLACK-CHEEKED FALCON (*Falco melanogenys*).—One only.

BOOBOK OWL (*Ninox boobook*).—Seen, often heard ; habitat seems fixed. Call sounds like " Moopoop."

MASKED OWL (*Strix novæ-hollandiæ*).—By Mr. A. J. Campbell's description ("Nests and Eggs," p. 50, No. 39), there seems to be no mistake. It has large facial disc margined with purplish colour, while back outside of wings are richly marbled with tan, grey, and white.



BHL

# Biodiversity Heritage Library

Mathews, Gregory Macalister. 1910. "List of Birds Observed on Parry's Creek, North-West Australia." *The Emu : official organ of the Australasian Ornithologists' Union* 9(4), 238-241.  
<https://doi.org/10.1080/01584197.1910.11798219>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/55266>

**DOI:** <https://doi.org/10.1080/01584197.1910.11798219>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/379684>

## Holding Institution

Smithsonian Libraries and Archives

## Sponsored by

Smithsonian

## Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.