1. Turdinulus Brevicaudatus (Blyth); Büttikofer, Notes Leyd. Mus. xvii. p. 75 (1895).

Specimens in Brit. Mus. from:-

Mount Mooleyit, 5000 ft. W. Limborg. Tweeddale Coll.

Mount Mooleyit, 5000 ft. W. Davison. Hume Coll.

Thoungyah. J. Darling. Hume Coll.

Mount Byingyi, 5500 ft., Shan States. E. W. Oates, Esq. [P.].

2. Turdinulus leucostictus (Sharpe); Büttikofer, op. cit. p. 75 (1895).

Specimens in Brit. Mus. from:-

Perak, Malay Peninsula. L. Wray, Esq. [P.]. Types of the species.

Coll.

3. Turdinulus striatus (Walden); Büttikofer, op. cit. p. 75 (1895).

Turdinus williamsoni, Godwin-Austen, J. As. Soc. Beng. xlvi. pt. ii. p. 44 (1877).

Specimens in Brit. Mus. from:-

Khasi Hills.

Tweeddale Coll. Type of the species.

Garo Hills.

Tenga Pani, near Sadiya.

Noa Dihing, " Chakha, Manipur Hills.

Konchungbuna, Manipur

Hills.

Type of Turdinus Godwin-Austen williamsoni, Godw .- Aust.

4. Turdinulus Roberti (Godwin-Austen & Walden).

Pnoëpyga roberti, Godwin-Austen & Walden, Ibis, 1875, p. 252; Godwin-Austen, J. As. Soc. Beng. xlv. pt. ii. p. 195 (1876); Hume, Str. F. iv. p. 218 (1876).

Turdinulus murinus, Oates, Birds Brit. Burmah, i. p. 62 (1883) [part; description of type only].

Specimens in Brit. Mus. from:-

Chakha, Manipur Hills.

Types of the species.

Asalu, Manipur Hills.

Eastern Naga.

Noa Dihing, near Sadiya.

Manbúm, near Sadiya.

Godwin-Austen Coll.

Subsp. a. Turdinulus guttaticollis, Grant, Ibis, 1895, p. 432.

Specimens in Brit. Mus. from:-

Miri Hills. Godwin-Austen Coll. Mishmi Hills.

Type of the species.

5. Turdinulus exsul, Sharpe; Büttikofer, Notes Leyd. Mus. xvii. p. 76 (1895).

Turdinulus roberti, Hume & Davison (nec Godw.-Aust. & Wald.), Str. F. vi. p. 234 (1878); Sharpe, Notes Leyd. vi. p. 173 (1884); Oates, Faun. Brit. Ind., Birds, i. p. 176 (1889).

Turdinulus murinus, Hume (nec Blyth), Str. F. ix. p. 115 (1880); Oates, Birds Brit. Burmah, i. p. 62 (1883) [except description of type].

Specimens in Brit. Mus. from:-

Thoungyah.
Mt. Mooleyit.
W. Davison.
W. Limborg. Tweeddale Coll.

Cheeks and sides of the throat washed with rust - colour: underparts less clearly suffused with tawny buff.

Klang, Selangore. W. Davison. Hume Coll.

Cheeks and sides of the throat with scarcely a trace of rust-colour (as in Bornean examples); underparts mostly whitish, partially suffused and edged with pale buff and reddish brown.

Kina Balu. A. H. Everett, Esq. [C.]. Mt. Dulit. C. Hose, Esq. [C.]. Mt. Penrisen. C. Hose, Esq. [C.].

Cheeks and sides of the throat mostly white, with only a slight trace of rustcolour; underparts rather more richly suffused with tawny buff.

By some ornithologists the Tenasserim birds may be considered worthy of specific distinction, but the differences between them and the Bornean specimens are so slight that I prefer to include them under the heading of *T. exsul*.

With regard to the specimen from Klang, I think it quite possible that it may represent a distinct species, but without more material I cannot give a decided opinion. Mr. E. W. Oates was evidently also at one time inclined to consider the Klang specimen distinct, for it bears a MS. name in his handwriting, which, however, he never published.



Ogilvie-Grant, W. R. 1896. "On a new Species of Tit Babbler from the Naga Manipur Hills." *Ibis* 2(1), 61–62.

https://doi.org/10.1111/j.1474-919x.1896.tb06975.x.

View This Item Online: https://www.biodiversitylibrary.org/item/55020

DOI: https://doi.org/10.1111/j.1474-919x.1896.tb06975.x

Permalink: https://www.biodiversitylibrary.org/partpdf/382334

Holding Institution

Smithsonian Libraries and Archives

Sponsored by

Smithsonian

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.