

Initial Report on Discovery of Ordovician Scolecodonts from Eastern Australia.

TERRY FUREY-GREIG

Department of Environmental Sciences, University of Technology Sydney,
P.O. Box 123, Broadway 2007, Australia

FUREY-GREIG, T. (1999). Initial Report on Discovery of Ordovician scolecodonts from eastern Australia. *Proceedings of the Linnean Society of New South Wales* **121**, 85–88.

Scolecodonts recovered from conodont residues of a Late Ordovician (Eastonian) limestone block contained in the Wisemans Arm Formation include a range of elements and represent the first documented Ordovician scolecodont fauna from Australia.

Manuscript received 31 May 1999, accepted for publication 22 September 1999.

KEYWORDS: conodont, scolecodont.

INTRODUCTION

A clast of limestone from volcanoclastic conglomerate in the Wisemans Arm Formation (Leitch and Cawood 1980) at Roscommon, east of Manilla (Fig. 1) has yielded a small fauna of scolecodonts as a by-product of conodont extraction. The conodont fauna is Late Eastonian (Ea3) (Furey-Greig 1999), and correlates with faunas documented from Ea3 horizons in the Lachlan Fold Belt (eg. Trotter and Webby 1995; Zhen et al. 1999). Little has been published on Palaeozoic polychaete microfossils from Australia. The present report provides documentation of a scolecodont fauna from a residue of known age.

The olistostromal Wisemans Arm Formation (Leitch and Cawood 1980) is a fault-bounded unit within the structurally complex Tablelands Complex of the New England Fold Belt (Fig. 1), that crops out between Nundle and Warialda. The Wisemans Arm Formation is characterised by volcanoclastic sandstone and siltstone, rare radiolarian mudstone, and conglomerates and olistostromes containing masses of mafic and intermediate volcanics, limestones of Late Ordovician and Early Silurian ages (Furey-Greig 1999; in press) and chert.

Standard acetic acid dissolution of limestones at the Macquarie University Centre for Ecostratigraphy and Palaeobiogeography, and separation of residues using the method of Anderson et al. (1995), yielded scolecodonts and conodonts. Illustration was entirely digital using the method described by Furey-Greig (1999). Some vacuum-induced damage was sustained by the specimens, hence no attempt was made to re-position the fragile microfossils in case of further damage. Terminology used herein is based on that of Kielan-Jaworowska (1966).

FOSSIL POLYCHAETE TAXONOMY

Historically, the common occurrence of scolecodonts (the acid-resistant jaw elements of polychaete annelids) as disarticulated, dispersed elements in Palaeozoic acid-insoluble residues led to development of a complex form taxonomy along similar lines




Figure 1. A. Major structural elements of southern New England Fold Belt; B. Location of study area; C. Geological map of Roscommon olistostrome. Grid co-ordinates in Australian Metric Grid System.

to the early study of conodonts. This led to problems in reconciling that system with documented natural assemblages (eg., Jansonius and Craig 1974). The complication of early, premature attempts at reconciling the two systems, and subsequent advances in this area, were discussed in detail by Eriksson and Bergman (1998). Several multi-element scolecodont studies have provided a basis for improving the taxonomic situation (eg, Kielan-Jaworowska 1966; Jansonius and Craig 1974; Bergman 1989). It has been




Figure 2. Scale bars items 2–8 = 100 microns, item 1 = 1mm. 1. Right MI element, lateral view. 2. ?broken MII element, lateral view. 3, 6. M4 elements, lateral views. 4. ?Basal plate, lateral view. 5. unassigned element, dorsal view. 7. M2 element, lateral and dorsal views. 8. broken element.

shown in these and other reports that only a small number of scolecodont elements, the MI and MII, are genus or species-diagnostic (see Szaniawski 1996 for a detailed review).

The stratigraphic value of Palaeozoic scolecodonts is hampered by their strong relationship with facies (Szaniawski 1996) and by the lack of stratigraphic information in many publications (Eriksson and Bergman 1998). One exception is Hints' (1998) report on scolecodonts from the Caradoc of Estonia and the St. Petersburg region. It is hoped that this preliminary report on Late Ordovician scolecodonts from Australia, well constrained as to age, will lead to further documentation of such faunas, for scolecodonts are known to occur in other eastern Australian Late Ordovician conodont residues (Percival 1999; Zhen, pers. comm.).

DISCUSSION

The fauna from Roscommon numbers forty elements and includes only one broken dextral Mandible 1, generally accepted as being the genus-diagnostic element. This is insufficient for precise taxonomic assignment and the material is here left in open nomenclature. Nevertheless, some comparison with documented Ordovician material is appropriate. The M1 element (Fig. 2, item 1) features a pronounced falx and edentulate falcate arch and is comparable to jaws of Atraktoprionidae illustrated by Kielan-Jaworowska (1966, plates XXX1–XXX1V), and by Hints (1998, Fig. 16). The other elements may belong to one or more species and are illustrated for future comparison.

ACKNOWLEDGEMENTS

I thank my supervisor, Evan Leitch, for much support and material assistance during this research. Ian Percival and Theresa Winchester-Seeto are thanked for reviewing the manuscript. Theresa Winchester-Seeto was generous with some rare literature on fossil polychaetes and discussion. Ruth Mawson and John Talent are thanked for underwriting the acid processing of limestones at Macquarie University. My research benefitted from financial support from ARC Large Grant A39601646 (E.C. Leitch) and from a Doctoral Research Scholarship provided by the Faculty of Science, University of Technology Sydney. Noni Callan drafted the figure. Ambrose and Patricia Blanche, owners of Roscommon, are thanked for permitting access to their property.

REFERENCES

- Anderson, M.A., Dargan, G., Brock, G.A., Talent, J.A., and Mawson, R. (1995). Maximising efficiency of conodont separations using sodium polytungstate. *Courier Forschungsinstitut Senckenberg* **182**, 515–522.
- Bergman, C.F. (1989). Silurian paulinitid polychaetes from Gotland. *Fossils and Strata* **25**, 128p.
- Eriksson, M. and Bergman, C.F. (1998). Scolecodont systematics exemplified by the polychaete *Hadoprion cervicornis* (Hinde, 1879). *Journal of Paleontology* **72**(3), 477–485.
- Furey-Greig, T.M. (1999). Late Ordovician conodonts from the olistostromal Wisemans Arm Formation, New England region, Australia. *Geologische Bundesanstalt (Wien), Abhandlungen* **54**. "North Gondwana: Terranes, Stratigraphy and Biota" [IGCP 421], 303–321.
- Furey-Greig, T.M. (in press). Late Ordovician and Silurian conodonts from the "Uralba Beds", east of Manilla, northern New South Wales. *Alcheringa* **24**.
- Hints, O. (1998). Late Viruan (Caradoc) polychaete jaws from North Estonia and the St. Petersburg region. *Acta Palaeontologica Polonica*, **43**(3) 471–516.
- Jansonius, J. and Craig, J.H. (1974). Some scolecodonts in organic association from Devonian strata of western Canada. *Geoscience and Man* **IX**, 15–26.
- Kielan-Jaworowska, Z. (1966). Polychaete jaw apparatuses from the Ordovician and Silurian of Poland and a comparison with modern forms. *Palaeontologia Polonica* **16**, 152p., 56 plates.
- Leitch, E.C. and Cawood, P. (1980). Olistoliths and debris flows at ancient consuming plate margins: an eastern Australian example. *Sedimentary Geology* **25**, 5–22.
- Percival, I.G. (1999). Late Ordovician biostratigraphy of the northern Rockley-Gulgong Volcanic Belt. *Quarterly Notes of the Geological Survey of New South Wales* **108**, 1–8.
- Szaniawski, H. (1996). Scolecodonts; in: Jansonius, J. and McGregor, D.C. (eds), *Palynology: principles and applications*, American Association of Stratigraphic Palynologists Foundation **1**, 337–354.
- Trotter, J.A. and Webby, B.D. (1995). Upper Ordovician conodonts from the Malongulli Formation, Cliefden Caves area, central New South Wales. *AGSO Journal of Australian Geology and Geophysics* **15**, 475–499.
- Zhen, Y.Y., Webby, B.D. and Barnes, C.R. (1999). Upper Ordovician conodonts from the Bowan Park Succession, central New South Wales, Australia. *Geobios* **32**, 73–104.



Furey-Greig, Terry. 1999. "Initial report on discovery of Ordovician scolecodonts from eastern Australia." *Proceedings of the Linnean Society of New South Wales* 121, 85–88.

View This Item Online: <https://www.biodiversitylibrary.org/item/108642>

Permalink: <https://www.biodiversitylibrary.org/partpdf/47536>

Holding Institution

MBLWHOI Library

Sponsored by

Boston Library Consortium Member Libraries

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.