

## MISCELLANEOUS.

*Descriptions of a new Genus and two new Species of African Freshwater Crabs.* By MARY J. RATHBUN.

THE crabs described below were found in mud under boards and timbers by Mr. J. H. Camp at Stanley Pool, Congo, West Central Africa.

## Family Thelphusidæ.

*Parathelphusa Campi*, sp. n.

Carapace subquadrilateral, conspicuously punctate. Depression between the gastric and cardiac regions deep. Protogastric lobes prominent, separated by a median groove which extends backward from the frontal margin. The postfrontal crest begins behind the base of the eyestalk and is continued to the lateral margin; it is finely denticulate and is almost straight, sloping backward and outward. Front a little more than one third the width of the carapace, deflexed, divided by a very shallow sinus into two lobes, with a raised margin, which is continued to the postfrontal crest. Superior orbital margin sinuous, advanced in its middle portion. Postorbital tooth acute, prominent. Lateral branchial spines three. In one of the two specimens there is a spinule between the first and second spines and a short fourth spine on the left side. Between the first spine and the postfrontal crest there are two or three spinules, forming a short ridge in the same line with the postfrontal crest. From the last spine a raised line extends backward upon the carapace, and is followed by several broken parallel lines. The inferior margin of the orbit is rather deeply rounded below the outer angle; from that point the margin is directed inward and forward; the inner angle is obtuse; the margin is set with a row of bead-like granules. The terminal segment of the abdomen of the female is subtriangular and the length is nearly half the width. The merus of the maxillipeds is very transverse, the antero-external angle rounded, the antero-internal angle not deeply cut.

Chelipeds of female unequal; merus rugose above, inner margin tuberculous, with a sharp spine just below the margin; carpus with two spines on the inner margin. Hand slightly inflated; upper margin straight, lower slightly convex. Fingers irregularly toothed and almost touching. Ambulatory legs rather slender, flattened; propodal joints indistinctly spinulose on the margins.

*Measurements* (No. 18065, two females).—Length of larger specimen 21.5 millim., width 29.5; length of smaller specimen 19, width 27.

This species in its three lateral teeth resembles *P. pæcilei*, A. Milne-Edwards, but differs from that species in its narrower and more quadrate carapace and interrupted postfrontal crest.

## ERIMETOPUS, gen. nov.

Carapace arcuate anteriorly, quadrate posteriorly, moderately convex. Front advanced beyond the antennular cavities, composed of two distinct rounded lobes. Orbits narrow; eyestalks tapering



to the extremity. Postfrontal crest short and inconspicuous or wanting. Lateral margins spinous. The merus of the maxillipeds is transverse, the antero-external angle rounded, the palpus articulating at the internal angle, which is very slightly notched. Chelipeds with a row of spines on the anterior margin of the carpus. Ambulatory legs with margins spinous.

*Erimetopus spinosus*, sp. n.

Carapace about four fifths as long as broad, convex longitudinally, postero-lateral margins long, sloping slightly inward and backward, antero-lateral margins arcuate. The cardiac region and the posterior portion of the gastric region are outlined by shallow depressions. Front about one third the width of the carapace, advanced, two-lobed, lobes separated by a broad V-shaped sinus. Margin of front and orbits granulous. Orbits well-defined, the outer angle a sharp incurved spine. There are two protogastric lobes, little elevated and often not discernible, except by two transverse lines of a lighter colour. A very shallow median groove extends backward from the frontal margin and forks directly behind the protogastric lobes. The postfrontal crest, when present, is short, arcuate, tuberculous, and indistinct; it begins back of the cornea, and for a short distance is nearly straight, directed outward and slightly forward, then curves almost parallel to the antero-lateral margin. In most specimens, however, the crest is obsolete, indicated only by the smoothly rounded elevation behind the orbit. Antero-lateral margin with a row of from five to eight spines next the orbit, of which the orbital spine is the largest. The spines are irregular in size and position. On the anterior branchial region are five or more marginal spines separated by a space from the hepatic spines; the anterior is by far the larger and is sometimes bifid. The others decrease in length posteriorly. The suborbital margin is granulous except at the notch beneath the postorbital spine. The abdomen of the female covers the sternum.

Chelipeds in the female unequal. The margins of the merus are spinulous, the upper margin with a sharp spine near the carpus, the inner surface tuberculous at its base, the lower surface with a transverse tuberculous ridge at its distal end. The carpus has two strong spines on its inner margin and a row of about seven smaller spines on the anterior margin, two of which are close to the condyle of the manus and are separated by a wide interval from the remaining spines. Sometimes one of the spines is bifid, and occasionally additional spinules occur on the upper surface behind the marginal spines. The manus is slightly swollen, with a convex lower margin and almost straight upper margin, which sometimes in the smaller cheliped has a small sharp spine at the distal end; in a few specimens there are one or two spines at the proximal end. Fingers irregularly dentate on their prehensile edges and slightly gaping. The ambulatory legs are rather broad, flattened; meral joints with two spines at the distal end; carpal and propodal joints strongly spined above; carpal joints with distal spines in the first pair and often in the second and third pairs; propodal joints with a few spines below; dactyli with four rows of spines.



*Measurements* (No. 18066, female). — Length 30.5 millim.; reatest width 38; width between postorbital spines 19.

The male is unknown.

This species can be distinguished from other Thelphusidæ by its produced round-lobed front, narrow orbits, and numerous spines.—*Proc. Nat. Mus.* vol. xvii. no. 980, pp. 25–27. (Communicated by the Author.)

*The Dipterous Parasites (Sarcophagidæ) of Locusts.—Apteny and Parasitic Sterilization.* By M. J. KÜNCKEL D'HERCULAIS.

Among Diptera the countless species comprised in the family Muscidæ furnish their contingent of enemies of the Locust, some of which attack their victims in the perfect stage, while others prey upon the eggs which are deposited in the ground; not content with playing a beneficent rôle in contributing to a large degree to arrest the multiplication of the Acridians, they present biological peculiarities and possess physiological attributes which are worthy of being recorded.

We shall devote our attention in the first instance to the Muscidæ which prey upon Acridians.

At the time of the great invasion of Northern Africa by migratory locusts in 1866 it was found at different points in Algeria (military subdivisions Aumale and Médéah) that Muscid larvæ were present in a certain number of these insects\*. In 1889 the considerable invasion of *Stauronotus maroccanus* gave an opportunity of making fresh observations in the Department of Constantine, in civil as well as in military territory; I myself found around Constantine, which was besieged by the Acridians, that a quantity of *Stauronotus*, as well as of other species, were affected; but it was not until the year 1890, at Teniet-el-Haad (Department of Algiers), that I was able to carry out methodical investigations. Just as in the previous year, I observed everywhere where the bands of locusts, escaping from destruction, had acquired their wings, that numbers of individuals crawled along among the herbage without having been able to follow their companions, the flights of which traversed the air. I had some sacks full of them collected: a few days afterwards some hundreds of larvæ were crawling at the bottom of these sacks. If the return of 1889† showed that 65 per cent. of the locusts lagging behind were infected with parasites, that which I made in 1890 gives the number thus affected as 75 per cent., and a post-mortem examination revealed the fact that each *Stauronotus* contained one, two, or three Muscid larvæ.

These larvæ hid themselves immediately in earth which was supplied to them in order to undergo the transformation into the

\* Letter of the General of Division commanding the province of Algiers (General de Wimpffen) to the Marshal the Governor of Algeria (Marshal MacMahon), 25th July and 5th August, 1866: registered nos. 2541 and 2776.

† Return drawn up at my initiative by the exertions of M. L. Tardieu, administrator of the mixed commune of R'hiras (Department of Constantine). Observations of M. Chartrousse, deputy-administrator.



Rathbun, Mary Jane. 1894. "Miscellaneous." *The Annals and magazine of natural history; zoology, botany, and geology* 14, 72–74.

<https://doi.org/10.1080/00222939408677768>.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/53354>

**DOI:** <https://doi.org/10.1080/00222939408677768>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/51488>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Smithsonian

**Copyright & Reuse**

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.