THE ENCARSIA CUBENSIS SPECIES-GROUP (HYMENOPTERA: APHELINIDAE)

GREGORY A. EVANS AND ANDREW POLASZEK

(GAE) Entomology and Nematology Department, University of Florida, PO Box 110620, Gainesville, FL 32611-0620, U.S.A. (e-mail: gae@gnv.ifas.ufl.edu); (AP) International Institute of Entomology, % The Natural History Museum, Cromwell Road, London SW7 5BD, U.K.

Abstract.—Taxonomic, host, and geographic range information for the six species comprising the *Encarsia cubensis* species-group is provided. Three new species, *E. aleurothrixi, E. hansoni*, and *E. hamoni*, are described from Brazil, Costa Rica, and Florida (USA), respectively. A lectotype is designated for *E. nigricephala* Dozier.

Resumen.—Se informa sobre la taxonomía, los hospederos y la distribución geográfica de seis especies que pertenecen al grupo de la especie *Encarsia cubensis*. Se describe tres nuevas especies, *E. aleurothrixi, E. hansoni* y *E. hamoni* recolectadas en Brasil, Costa Rica y Florida (EEUU), respectivamente. Se designa el lectotipo para *E. nigricephala* Dozier.

Key Words: Encarsia, whiteflies, parasitoids, biological control, taxonomy, Nearctic, Neotropical, Bemisia, Aleurothrixus.

As here defined, the Encarsia cubensis species-group consists of six species, all of which parasitize whiteflies in the New World, except for one species (E. nigricephala Dozier) which apparently was accidently introduced beyond its natural range into Réunion (Mascarene Islands, Indian Ocean). Various economically important whitefly species are attacked by members of this group. Encarsia nigricephala is one of the most common parasitoids of the Bemisia tabaci species-complex (Aleyrodidae) throughout the New World. Encarsia quaintancei Howard is a major parasitoid of the banded-winged whitefly, Trialeurodes abutiloneus (Haldeman), and also attacks the Bemisia tabaci species-complex.

Gahan (1931) first recognized these species as comprising a distinct group in the genus *Encarsia* and characterized it as species having only four tarsomeres on the middle tarsus and the presence of an area around the stigmal vein devoid of setae. DeBach and Rose (1981) erected the *Encarsia cubensis* species-group. *Encarsia cubensis* and *E. quaintancei* had previously been included in the 'formosa-group' (= lu-teola-group) by Viggiani and Mazzone (1979). Polaszek et al. (1992) provided taxonomic, host and distribution information for *E. nigricephala* and *E. quaintancei*. Schauff et al. (1996) updated host and distribution records for these two species, and provided information on *E. cubensis*.

The *Encarsia cubensis* species-group can be defined using a combination of characters, none of which alone, however, distinguishes this group from other speciesgroups of *Encarsia*. For example, species belonging to the *parvella*- and *citrina*-

Species Group	Tarsal Formula	Fore Wing Asetose Area	Fore Wing Disc Width	Gastral Tergite	Host	Male Antenna
cubensis	5-4-5	yes	normal-broad	smooth	whitefly	F2 modified
luteola	5-4-5	no	normal-broad	smooth	whitefly	normal
singularis	5-4-5	no	narrow	sculptured	diaspine scale	F1 modified
citrina	5-5-5	yes	narrow-normal	sculptured	diaspine scale	normal
parvella/ pergandiella	5-5-5	yes	narrow-broad	smooth	whitefly	normal

Table 1. Characteristics of the *Encarsia cubensis* species group compared with those of similar *Encarsia* species groups.

groups also possess the asetose area around the stigmal vein of the fore wing. The 4segmented mid tarsus is also found in the *luteola-* and *singularis* groups, as well in some unplaced species (e.g. *E. africana* (Hill) and *E. mohyuddini* Shafee and Rizvi). Reduction in the number of segments of the mid tarsus appears to have occurred independently in different lineages within *Encarsia*. Certainly within the Aphelinidae and the Chalcidoidea as a whole, reduction in number of tarsal segments by fusion of the two distal segments is common.

An ovoid sensory/glandular structure is present on F2 antennal segment of the males of the three species in this group whose males are known, and is present on both the F1 and F2 segment in the male of E. hansoni. Potentially, this structure is a unique synapomorphy for the cubensis species group; however, males of the singularis-group, a small group of armored scale parasitoids from the Oriental region, have a similar structure on the F1 antennal segment. Pedata et al. (1995) first demonstrated that these structures, often referred to as 'sensorial complexes' (Viggiani and Mazzone 1982, Polaszek et al. 1992), are at least partially glandular. Viggiani (1996) demonstrated that in E. nigricephala (and presumably in E. quaintancei and E. hamoni) the structure is entirely glandular. The form of these structures agrees at least partially with the 'release and spread' function attributed to similar antennal glands by Isidoro et al. (1996).

Within Encarsia, members of the E. cub-

ensis-group show greatest affinity with members of the *parvella*-group, with which they share the asetose area of the fore wing, and from within which they may be derived. Characteristics of the *E. cubensis*group are compared with those of related *Encarsia* species groups (Table 1).

The following six species are placed in the *E. cubensis* group: *E. aleurothrixi* Evans and Polaszek, n. sp., *E. cubensis* Gahan, *E. hamoni* Evans and Polaszek, n. sp., *E. hansoni* Evans and Polaszek, n. sp., *E. nigricephala* Dozier and *E. quaintancei* Howard.

Morphological terminology follows that of Hayat (1989). The scutellar setae are designated Sc1 (anterior pair) and Sc2 (posterior pair) as used in Evans et al. (1995). The length of the fore wing disc (Fig. 18, fd) is measured from the midpoint of an imaginary line drawn between the apex of the stigmal vein and the distal end of the frenal fold of the fore wing, to the fore wing apex. The width of the fore wing (Fig. 18, fww) refers to the maximum width of the fore wing; the length of the marginal fringe, refers to the length of the longest seta of the marginal fringe. Habitus drawings for each species show the gaster divided medially, with the dorsum on the left side and the venter on the right side. The tibia and tarsus of the middle leg for the female of each species is drawn on its left side. An asterisk is placed before new host and/or distribution records.

Holotypes of all members of the *E. cub*ensis species group are deposited in the National Museum of Natural History, Smithsonian Institution (USNM) in Washington, D.C., U.S.A. Paratypes of *E. aleurothrixi* are deposited in The Natural History Museum, London, U.K. (BMNH), Florida State Collection of Arthropods, Gainesville, Florida, U.S.A (FSCA) and in the G. A. Evans, personal collection (GAE). Additional specimens of *E. cubensis* and *E. nigricephala* are deposited in the Centre de Cooperation Internationale en Recherche Agronomique pour le Developpement, Montpellier, France (CIRAD).

ENCARSIA CUBENSIS SPECIES-GROUP

Diagnosis.—Tarsal formula 5-4-5; fore wing moderate to broad in width with large asetose area beneath the stigmal vein, longest seta of marginal fringe not longer than maximum width of fore wing, distance between placoid sensilla on scutellum more than $2.5 \times$ the diameter of one sensillum, base of gastral tergite I smooth.

Additional characteristics observed in species of this group include: antennal club 3-segmented, F1 antennal segment usually short, quadrate or nearly so and shorter than pedicel, ovipositor short to moderate in length, subequal to length of tibia II and not exceptionally extruded, mesoscutum with 2 to 3 pairs of setae, males with antennal segments F5 and F6 fused, F2 (rarely F1) with an ovoid sensory/glandular structure covered by a plate (Figs. 11, 15).

KEY TO ENCARSIA CUBENSIS SPECIES-GROUP

2

3

4

1	Mesoscutum with 2 pairs of setae, gaster com-
	pletely yellow, or dark brown with central por-
	tion of tergites I and II yellow, (Figs. 4, 6, 8,
	10
-	Mesoscutum with 3 pairs of setae, gaster com-
	pietery dark brown. (Figs. 2, 14)
2.	Gaster completely yellow, fore wing almond-
	shaped, pointed at apex (Figs. 19, 21); males
	known
-	Gaster dark brown with central portion of ter-
	gites I and II yellow, fore wing round at apex
	(Figs. 18, 20); males unknown
3.	Scutellum dark brown, F1 oblong and quad-
	rate, F6 elongate, 3.2× as long as wide (Fig.
	7), 2 pairs of setae between cerci of tergite VI.

(Fig. 8). Male with ovoid glandular structure on F1 and F2, F3 slightly longer than F2 ...

Scutellum yellow, F1 cylindrical, 1.5–1.7× as long as wide, F6 not exceptionally elongate, 2.4× as long as wide (Fig. 9), 1 pair of setae between cerci of tergite VI. (Fig. 10). Male (Fig. 12) with ovoid glandular structure on F2, F3 shorter than F2 segment (Fig. 11)

- 4. Valvula III elongate (Fig. 4), 0.6× as long as ovi-
- positor, F2 longer than F3 (Fig. 3) cubensis – Valvula III $0.4 \times$ as long as ovipositor (Fig. 6),
- F2 shorter than F3 (Fig. 5) hamoni
- 5. Axillae and posterior half of mesoscutum yellow, second pair of mesoscutal setae proximal to anterior pair of setae (Fig. 2), valvula III with spatulate apical setae, F1 quadrate, $0.5 \times$ as long as F2 (Fig. 1); male unknown.

Encarsia aleurothrixi Evans and Polaszek, new species (Figs. 1, 2a, 2b, 17)

Diagnosis.-The female of E. aleurothrixi can be distinguished from other Encarsia cubensis group species by the unusual spatulate setae located on the apex of valvula III (may be difficult to discern unless viewed with phase-contrast microscopy), and by the anterolateral position of the second pair of mesoscutal setae. This species is most similar to E. quaintancei in coloration and number of mesoscutal setae, but can be distinguished from the latter species which has the second pair of mesoscutal setae located approximately halfway between the anterior and posterior pair of setae, the F1 antennal segment more elongate, the axillae dark brown, and slender apical setae on valvula III.

Description.—Female holotype. Coloration (Fig. 2): Head dark brown, eyes red, antenna pale, anterior half of mesoscutum, metanotum and gaster, dark brown; rest of



Figs. 1–8. 1–2 Encarsia aleurothrixi 1, Antenna, female. 2a, Habitus, female. 2b, Valvula III, apex. 3–4, E. cubensis. 3, Antenna, female. 4, Habitus, female. 5–6, E. hamoni. 5, Antenna, female. 6, Habitus, female. 7–8, E. hansoni. 7a, Antenna, female. 7b, Antenna, male. 8, Habitus, female.

mesosoma and legs, yellow; coxa III and valvula III infuscate; wings hyaline. Morphology: Antennal (Fig. 1) radicle (R), scape (S), pedicel (P), 3 funicle segments (F1-F3) and 3 club segments (F4-F6) with the following ratios of length to width: R: 3.6, S:4.5, P:1.4, F1:1.0, F2:1.7, F3:1.7, F4: 1.5, F5:1.5 and F6:1.8; relative lengths of segments R-F6 to length of F1: R:1.8, S: 4.5, P:1.8, F1:1.0, F2:1.9, F3:2.1, F4:2.1, F5:2.1, and F6:2.4; F1-F6 with the following number of linear sensilla: F1:0, F2:2, F3:2, F4:2, F5:3, F6:3. Mesoscutum with narrow, hexagonal sculpture, and 3 pairs of slender setae arranged as in Fig. 1; each side lobe with 2 setae; each axilla with 1 short seta, scutellar seta Sc1 0.7× as long as Sc2; distance between scutellar sensilla $5 \times$ the width of one sensillum; endophragma short, reaching halfway to posterior base of gastral tergite I; tibial spur of middle leg $0.7 \times$ corresponding basitarsus. Fore wing (Fig. 17) broad, disc $0.9 \times$ as long as wide, marginal fringe $0.3 \times$ width of wing; discal setae uniformily distributed except wide bare area along the distal margin of the fore wing and area surrounding stigmal vein; 6-7 costal, 2-3 basal group, and 2 submarginal vein setae; marginal vein with 5-6 long and stout setae along its anterior margin, 2 large setae at its base, and 7-8 short setae along its interior. Gastral tergites I-V dorsum with reticulate lateral margins, tergites V and VI with stipules, centrally; tergites I-VII with 0, 1, 1, 1, 2, 2, 2 pairs of setae, respectively. Venter with 2 pairs of setae at sternite II, III and IV, respectively; ovipositor arising at anterior base of tergite IV, as long as tibia of middle leg; valvula III slender, $0.5 \times$ ovipositor, with 2–3 pairs of distally-expanded, apical setae.

Male.—Unknown.

Distribution.-Brazil.

Hosts.—Aleurothrixus aepim (Goldi), Aleurothrixus floccosus (Maskell).

Material examined.—Brazil, Aguas de São Paolo, 24.i.1988, ex. Aleurothrixus aepim on Manihot esculentum, F.D. Bennett, (Holotype $\[mathbb{]$ and 2 $\[mathbb{]$ paratypes, USNM; 5 $\[mathbb{]$ paratypes, BMNH; 8 $\[mathbb{]$ paratypes, FSCA; 8 $\[mathbb{]$ paratypes, GAE); Brazil, Piracicaba, 27.v.1991, ex. Aleurothrixus floccosus on Citrus sp., F. D. Bennett, (12 $\[mathbb{]$ paratypes, GAE).

Etymology.—The species name is derived from its host, *Aleurothrixus*.

Encarsia cubensis Gahan (Figs. 3, 4, 18)

Encarsia cubensis Gahan 1931:121. Trichoporus cubensis: Dozier 1933:92.

Types.—Three syntype females reared from *Aleurothrixus howardii* (= *A. floccosus*), Santiago de las Vegas, Cuba, received from S. C. Bruner, in USNM [examined].

Diagnosis.—The female of *E. cubensis* can be distinguished from other *E. cubensis* group species (except *E. hamoni*) by having 2 pairs of mesoscutal setae and dark brown gaster with central portion of tergites I and II, yellow. *E. cubensis* is most similar to *E. hamoni* in coloration and number of mesoscutal setae and can be distinguished from that species by having the F2 longer than F3 antennal segment, the F1 antennal segment very short, $0.5 \times$ as long as F2, and valvula III $0.6 \times$ as long as the ovipositor.

Description.—Female syntype. Coloration (Fig. 4): Head, pronotum, mesoscutum (except lateral margins), axillae (except small, basal triangular area), lateral margins of metanotum and gasterl tergites I and II, and tergites III–VI, dark brown; lateral margins of mesoscutum, side lobes, scutellum, central area of tergites I and II, tergite VII and legs, yellowish; F6 antennal segment infuscate; wings hyaline. Morphology: Antennal segments (Fig. 3) with the following length to width ratios: R:3.0, S:5.0, P:1.8, F1:1.0, F2:1.8, F3:1.6, F4:1.8, F5:1.8, and F6:2.3; relative lengths of segments R–F6 to length of F1: R:1.7, S:4.7, P:2.0, F1:1.0, F2:2.0, F3:1.8, F4:2.0, F5:2.2 and F6:2.5; linear sensilla present on flagellar segments F2-F6. Mesoscutum with broad, hexagonal sculpture and 2 pairs of short setae; each axilla with 1 short seta, scutellar seta Sc1 $0.6 \times$ as long as Sc2; distance between scutellar sensilla $5.2 \times$ width of one sensillum; endophragma short, not reaching tergite II; tibial spur of middle leg $0.8 \times$ length of corresponding basitarsus. Fore wing (Fig. 18) broad, disc $1.1 \times$ as long as wide, marginal fringe $0.4 \times$ width of fore wing; discal setae uniformily distributed except a wide asetose band along the distal margin of the fore wing and asetose area surrounding stigmal vein; 6-7 costal, 2 basal group, and 2 submarginal vein setae; marginal vein with 5-6 setae along its anterior margin, 2 setae at its base, and 7-8 short setae along its interior. Gastral tergites I-V dorsum with imbricate lateral margins, stipules present on tergites V-VI, centrally; tergites I-VII with 0, 1, 1, 1, 2, 2, 2 pairs of setae, respectively. Venter with a pair of setae at sternites I and II; ovipositor arising at level of tergite III, as long as tibia of middle leg; valvula III, $0.6 \times$ as long as ovipositor.

Male.—Unknown.

Hosts.—Aleurothrixus floccosus (Maskell), *Aleurotrachelus trachoides (Back).

Distribution.—Brazil; Cuba; *Dominican Republic; *Guadeloupe; Haiti; Puerto Rico; USA: Florida.

Specimens examined.—In addition to syntypes, Dominican Republic, Charco Azul, 16.i.1995, ex. ?Aleurothrixus sp. on Manihot esculentum, C. A. Serra (1 \Im , GAE); Guadeloupe: Vernon, 21.vi.1990, ex. *Aleurotrachelus trachoides, J. Etienne, (1 \Im , CIRAD); Dom. Dudos, 1.ii.1990, J. Etienne (1 \Im , CIRAD); Puerto Rico: Avecibo, 19.x.1990, Aleurothrixus floccosus on Tabebuia glomerata, F. D. Bennett (2 \Im , GAE); USA: Florida, Miami, 22.iii.1992, ex. whitefly on Cocos nucifera, F. D. Bennett (1 \Im , GAE).

Encarsia hamoni Evans and Polaszek, new species (Figs. 5, 6, 20)

Diagnosis.—The female of *E. hamoni* can be distinguished from the other *Encarsia cubensis* group species (except *E. cubensis*) by its dark brown gaster with yellow central area of tergites I and II, 2 pairs of mesoscutal setae, and rounded apex of the fore wing. *E. hamoni* is most similar to *E. cubensis* and can be distinguished from the latter species by having valvula III shorter, $0.4 \times$ as long as the ovipositor; F2 shorter than F3; and axillae completely dark brown.

Description.-Female holotype. Coloration (Fig. 6): Head, pronotum, mesoscutum (except lateral margins), axillae, lateral margins of metanotum and gastral tergites I and II, and tergites III-VI, dark brown; lateral margins of mesoscutum, side lobes, scutellum, central area of tergites I and II, tergite VII and legs, yellow; wings hyaline. Morphology: Antennal segments (Fig. 5) with the following ratios of length to width: R:1.9, S:4.5, P:1.9, F1:1.0, F2:1.6, F3:1.9, F4:1.5, F5:1.5 and F6:1.8; relative lengths of segments R-F6 to length of F1: R:1.9, S:4.5, P:1.9, F1:1.0, F2:1.6, F3:1.9, F4:1.9, F5:2.0, and F6:2.4; F1-F6 with the following number of linear sensilla: F1:0, F2:2, F3:2, F4:2, F5:3, F6:3. Mesoscutum with broad, hexagonal sculpture and 2 pairs of slender setae; each side lobe with 2 setae; each axilla with 1 seta, scutellar setae Sc1 0.7× as long as Sc2; distance between scutellar sensilla $5 \times$ the width of one sensillum; endophragma short, reaching almost to posterior base of gastral tergite I; tibial spur of middle leg 0.7× corresponding basitarsus. Fore wing (Fig. 20) broad, disc $0.9 \times$ as long as wide, marginal fringe $0.3 \times$ width of wing; discal setae uniformily distributed except wide bare area along the distal margin of the fore wing and the area surrounding stigmal vein; 6-7 costal, 1 basal group, and 2 submarginal vein setae; marginal vein with 5-6 long and stout setae along its anterior margin, 2 large setae at its base, and

7–8 short setae along its interior. Gastral tergites I–V dorsum with reticulate lateral margins, tergites V and VI with stipules, centrally; tergites I–VII with 0, 1, 1, 1, 2, 2, 2 pairs of setae, respectively. Venter with 2 pairs of setae at sternite II, III and IV, respectively; ovipositor arising at anterior margin of tergite IV, as long as tibia of middle leg; valvula III slender, $0.4 \times$ ovipositor.

Male.—Unknown.

Distribution.—U.S.A.: Florida, Georgia; Mexico.

Hosts.—*Tetraleurodes ursorum* (Cockerell); *Bemisia tabaci* species-complex.

Material examined.—U.S.A: Florida, Davie, 24.ii.1994, ex. *Tetraleurodes ursorum* on *Annona glabra*, A. B. Hamon, (Holotype $\[mathscale]$ and 3 $\[mathscale]$ paratypes, USNM; 2 $\[mathscale]$ paratypes, BMNH; 1 $\[mathscale]$ paratype, GAE). Additional specimens: USA: Georgia, Tifton, x.1992, *Bemisia tabaci* species-complex on *Gossypium hirsutum*, J. Chamberlain, (1 $\[mathscale]$, GAE); Mexico: Sinaloa, Guasave, 12.x.1990, B. Alvarado, (2 $\[mathscale]$, GAE).

Etymology.—This species is named in honor of Avas Hamon for his many years of assistance in the identification of whitefly and scale insect hosts.

Encarsia hansoni Evans and Polaszek, new species

(Figs. 7a, 7b, 8, 19)

Diagnosis.—The female of *E. hansoni* can be distinguished from the other *E. cub*ensis group species by having the scutellum completely dark brown; the F1 antennal segment short and oblong; and the F6 antennal segment very elongate, $3.4 \times$ as long as wide.

Description.—Female holotype. Coloration (Fig. 8): Head dark brown, eyes red, antennae pale; mesoscutum (except for lateral margins), axillae and scutellum, dark brown; metanotum, gaster, and legs, yellowish; wings hyaline. Morphology: Antennal segments (Fig. 7a) with the following ratios of length to width: R:4.0, S:5.0, P: 1.3, F1:0.8 (ventral margin) and 1.0 (dorsal margin), F2:1.8, F3:1.9, F4:1.8, F5:2.0 and F6:3.4; relative lengths of segments R-F6 to length of F1(ventral margin): R:2.0, S: 5.6, P:2.6, F1:1.0, F2:2.0, F3:2.1, F4:2.3, F5:2.5, and F6:3.9; F1-F6 with the following number of linear sensilla: F1:0, F2:1, F3:2, F4:3, F5:3, F6:3. Mesoscutum with broad, hexagonal sculpture and 2 pairs of slender setae; each side lobe with 2 setae; each axilla with 1 seta, scutellar setae Sc1 $0.9 \times$ as long as Sc2; distance between scutellar sensilla $5 \times$ the width of one sensillum; endophragma short, reaching anterior base of tergite II; tibial spur of middle leg $0.9 \times$ as long as corresponding basitarsus. Fore wing (Fig. 19) almond-shaped, disc $1.4 \times$ as long as wide, marginal fringe $0.5 \times$ width of wing; discal setae uniformily distributed except for wide asetose band along the distal margin and large, asetose area surrounding stigmal vein; 5 costal, 2 basal group, and 2 submarginal vein setae; marginal vein with 5 setae along its anterior margin, 2 large setae at its base, and 7-8 short setae along its interior. Gastral tergites II-V dorsum with imbricate lateral margins, tergite VI smooth, tergite VII rugose; tergites I-VII with 0, 1, 1, 1, 2, 3, 2 pairs of setae, respectively. Venter with a pair of setae at sternite I, II and III, respectively; ovipositor arising at level of tergite III, $1.1 \times$ as long as tibia of middle leg; valvula III slender, $0.4 \times$ length of ovipositor.

Male.—(Specimen collected in a pan trap near the location of the holotype female and assumed to be the male of E. hansoni.) Coloration: Body dark brown, lateral margins of mesoscutum, side lobes, base of axillae, and scutellum, yellow; gaster tergite VII light brown; legs and antenna, pale; wings hyaline. Antennal segments (Fig. 7b) with the following length to width ratios: R:2.5, S:3.7, P:1.1, F1:1.0, F2:1.1, F3:1.2, F4:1.6, F5 + F6 (fused):3.6; ratio of each segment to length of F1: R:1.0, S:2.1, P:1.0, F1:1.0, F2:0.9, F3:1.1, F4:1.4, F5 + F6 (fused):3.4; F2 with 2 ovoid sensory/glandular structures covered by plate having 2 holes; F1 with 1 ovoid sensory/glandular structure; proximal half of F3 surface with rugose area; mesosomal sculpturing and setation and fore wing similar to those of female.

Distribution.-Costa Rica.

Host.—Unknown.

Material examined.—Costa Rica, La Selva, 17.vii.1995, C. Godoy and P. Hanson, (Holotype \mathcal{P} , USNM); Costa Rica, 3 km south of Puerto Viejo, La Selva, i.1993, pan trap, P. Hanson, 1 \mathcal{E} , USNM).

Etymology.—This species in named in honor of Paul Hanson, University of Costa Rica.

Encarsia nigricephala Dozier (Figs. 9, 10, 11, 12, 21)

Encarsia nigricephala Dozier, 1937: 129.

Type.—Lectotype female and two paralectotype females (here designated) reared from *Bemisia* sp. on *Euphorbia hypericifolia*, Feb. 26, 1936, Mayaguez, Puerto Rico, in USNM. [examined]

Diagnosis.—The female of *E. nigrice-phala* can be distinguished from other *E. cubensis* group species by its dark brown head and anterior third to two thirds of mesoscutum, contrasting with the pale color of the remainder of the body. Males of *E. ni-gricephala* can be distinguished from *E. quaintancei* males by having 2 pairs of setae on the mesoscutum, the F2 longer than F3 antennal segment, and the hind tibia yellow. Males of *E. nigricephala* lack the ovoid sensory/glandular structure on the F1 antennal segment which is present in males of *E. hansoni.*

Description.—Female holotype. *Coloration*: (Fig. 10) Body yellowish with head, pronotum, and anterior third to two thirds of mesoscutum, dark brown; antenna and legs pale, F6 slightly infuscate; wings hyaline. Morphology: Antennal segments (Fig 9) with the following ratios of length to width: R:3.4, S:5.6, P:1.5, F1:2.0, F2:2.4, F3:2.0, F4:1.8, F5:1.7 and F6:2.3; relative lengths of segments R–F6 to length of F1: R:1.1, S:2.8, P:1.1, F1:1.0, F2:1.2, F3:1.1, F4:1.1, F5:1.2, and F6:1.5; linear sensilla present on F2–F6. Mesoscutum with me-



Figs. 9–16. 9–12, *Encarsia nigricephala*. 9, Antenna, female. 10, Habitus, feamle. 11, Antenna, male (dorsal/ventral). 12, Habitus, male. 13–16, *E. quaintancei*. 13, Antenna, female. 14, Habitus, female. 15, Antenna, male. 16, Habitus, male.

dium to broad hexagonal sculpture and 2 pairs of slender setae; each side lobe with 2 setae; each axilla with 1 seta, scutellar with Sc1 $0.6 \times$ as long as Sc2; distance between scutellar sensilla about $6 \times$ the width of one sensillum; endophragma short, reaching base of tergite II; tibial spur of middle leg 0.6-0.8× as long as corresponding basitarsus. Fore wing (Fig. 21) almondshaped, disc $1.3 \times$ as long as wide, marginal fringe 0.5× width of wing; discal setae uniformily distributed except wide asetose band along the distal margin and large, asetose area surrounding stigmal vein; 4-5 costal, 1 basal group, and 2 submarginal vein setae; marginal vein with 5-6 setae along its anterior margin, 2 setae at its base, and 6-7 short setae along its interior. Gastral tergites I-VI dorsum with imbricate lateral margins, tergite VII rugose; tergites I-VII with 0, 1, 1, 1, 2, 2, 2 pairs of setae,

respectively. Venter with a pair of setae at sternite II and III, respectively; ovipositor arising at level of tergite III, $0.9 \times$ as long as tibia of middle leg; valvula III slender, $0.4-0.5 \times$ length of ovipositor.

Male.--(Description based upon specimen reared from the Bemisia tabaci species-complex, Puerto Rico, Mayaguez, 31.v.90, on Lantana camara, F.D. Bennett.) Coloration: Body dark brown, lateral margins of mesoscutum, side lobes, base of axillae, and scutellum, yellowish; legs and antenna, pale; wings hyaline. Antennal segments (Fig. 11) with the following length to width ratios: R: 3.0, S:4.1, P:1.3, F1:0.9, F2:1.6, F3:1.2, F4:1.3, F5 + F6 (fused):3.1; ratio of each segment to length of F1: R: 1.4, S:2.8, P:1.2, F1:1.0, F2:1.8, F3:1.2, F4: 1.3, F5 + F6 (fused):3.2; F2 longer than F3 and with 2 large, ovoid sensory/glandular structures covered by plate having 2 holes.

Ventral margins of funicle segments F2 and F3 longer than dorsal margins; mesosomal sculpturing and setation and fore wing similar to those of female.

Morphological variation.—We have examined hundreds of specimens of this species from a wide variety of hosts throughout the New World and have found very little morphological variation. The size of the dark brown area on the mesoscutum varies occasionally. Normally, this area covers approximately the anterior half of the mesoscutum; however, several specimens reared from *Bemisia argentifolia* in Guadeloupe and Colombia have the mesoscutum nearly entirely dark brown. The posterior margin of the scutellum is dark brown in a few specimens reared from *B. tabaci* speciescomplex in Georgia.

One male specimen, reared from an aleyrodid species on *Sonchus* sp. in Piracicaba, Brazil by F.D. Bennett, differs from *E. nigricephala* by having the F2 wider than long, subequal in length to the F3 antennal segment, and the last segment of the mid tarsus, dark brown. This specimen may represent a variation of *E. nigricephala* or possibly a distinct species.

Hosts.—*Aleurodicus dispersus Russell, *Aleurotrachelus atratus Hempel, Bemisia argentifolia Perring and Bellows, B. tabaci (Genn.), *Crenidorsum sp., *Dialeurodes kirkaldyi (Kotinsky), *Tetraleurodes acaciae (Quaintance), Trialeurodes abutiloneus (Haldeman), T. floridensis (Quaintance), T. vaporiorarum (Westwood).

Distribution.—Barbados; Brazil; Colombia; *Ecuador; Grenada; Guadeloupe; *Guatemala; Honduras; Jamaica; Mexico; Puerto Rico; USA: Florida, Georgia, *Maryland, *Mississippi, Texas; Venezuela.

Specimens examined representing new host or distribution records.—Ecuador: Queredo, *Bemisia tabaci* species-complex on *Chamaesyce hirta*, 14.vi.1994, P. Stansly; Guatemala: (sent to Gainesville, Florida quarantine), 21.x.1991, ex. *Bemisia tabaci* species-complex; U.S.A., Florida, Indian River Beach, 10.iv.1987, ex. *Crenidorsum* sp., on Coccoloba uvifera, D. Mooney; U.S.A., Maryland, Colesville, 8.xi.1993, ex. Bemisia argentifolia on Lycopersicon esculentum, G. A. Evans; U.S.A., Mississippi, 1996, Bemisia tabaci species-complex on Gossypium hirsutum, M. Smith; U.S.A., Florida, Homestead, 5.ii.1991, Aleurodicus dispersus on ornamental tree, F. D. Bennett; U.S.A., Miami, 22.iii.1990, Aleurotrachelus atratus on Cocos nucifera, F. D. Bennett; U.S.A., Florida, Homestead, Dialeurodes kirkaldvi on ornamental tree, F. D. Bennett; Puerto Rico: Rio Piedras, 18.xi.1989, Tetraleurodes acaciae on Jatropa gossypifolia, F. D. Bennett. All of the above specimens deposited in GAE collec-Reunion: Piton tion. 1 Ŷ, Saint Leu, 21.v.1990, ex. Trialeurodes vaporariorum, B. Reynaud (CIRAD).

Encarsia quaintancei Howard (Figs. 13, 14, 15, 16, 22)

- Encarsia quaintancei Howard 1907: 79. Prospaltella perspicuipennis Girault 1910: 234.
- *Encarsia perspicuipennis*: Viggiani 1986: 71, n. comb., lectotype designated.
- Prospaltella perspicuipennis: Polaszek et al. 1992: 387, synonymy.

Type.—Holotype female reared from *Aleyrodes* sp. on *Polygonum* sp., Washington, D.C., Bladensburg Road, 29.viii.1900, Theo. Pergande, in USNM [examined]. We consider the identification of the whitefly host of the holotype specimen as *Aleyrodes* sp. to be is erroneous and probably represents either *Trialeurodes abutilonea* or *T. vaporariorum* listed by Mound and Halsey (1978) as the whitefly species found on this host plant in the United States.

Diagnosis.—The female of *E. quaintancei* can be distinguished from other *E. cubensis* group species by having the scutellum bright yellow, contrasting with its dark brown head, mesoscutum and gaster, the F1 antennal segment cylindrical, and three pairs of mesoscutal setae. Males of this species can be distinguished from *E. nigrice-*



Figs. 17–22. Fore wings of Encarsia females. 17, E. aleurothrixi. 18, E. cubensis. 19, E. hansoni. 20, E. hamoni. 21, E. nigricephala. 22, E. quaintancei.

phala and *E. hansoni* males by having three pairs of mesoscutal setae, the F2 shorter than the F3 antennal segment, and the hind tibia brownish.

Description.-Holotype female. Coloration (Fig. 14): head, pronotum, axillae, metanotum and gaster, dark brown, scutellum bright lemon yellow, legs and antenna yellowish, F6 slightly infuscate; wings hyaline. Morphology: Antennal segments with the following ratios of length to width: R: 3.0, S:6.0, P:1.9, F1:1.3, F2:1.6, F3:2.2, F4: 1.9, F5:1.7 and F6:2.4; relative lengths of segments R-F6 to length of F1: R:1.5, S: 4.2, P:1.7, F1:1.0, F2:1.3, F3:1.8, F4:1.7, F5:1.7, and F6:2.3; F1-F6; linear sensilla present on F3-F6. Mesoscutum with narrow to medium hexagonal sculpture and 3 pairs of slender setae; each side lobe with 3 setae; each axilla with 1 short seta, scutellar setae Sc1 $0.7 \times$ as long as Sc2; distance between scutellar sensilla $3\times$ the width of one sensillum; endophragma short, not reaching posterior margin of tergite I; tibial spur of middle leg $0.75 \times$ as long as corresponding basitarsus. Fore wing (Fig. 22) apex rounded, disc $1.3 \times$ as long as wide, marginal fringe $0.34 \times$ width of wing; discal setae uniformily distributed except wide asetose band along the distal margin and large and large asetose area surrounding stigmal vein; 5-6 costal, 2-3 basal group, and 2 submarginal vein setae; marginal vein with 5 setae along its anterior margin, 2 setae at its base, and 6-7 short setae along its interior. Gastral tergites I-VI dorsum with imbricate lateral margins, tergite VII rugose; tergites I-VII with 0, 1, 1, 1, 2, 2, 2 pairs of setae, respectively. Venter with a pair of setae at sternite II and III,

respectively; ovipositor arising at level of tergite IV, as long as tibia of mid leg; valvula III slender, $0.5 \times$ length of ovipositor.

Male.-(Description based upon specimen reared from Bemisia tabaci speciescomplex on Emilia fosbergi, USA, Florida, Ft. Pierce, 28.x.1990, F. D. Bennett). Coloration: Body (Fig. 16) dark brown, lateral and posterior margins of mesoscutum, side lobes, and scutellum, yellowish; antenna and legs pale, coxa III and femur III, brownish; wings hyaline. Antennal segments (Fig. 15) with the following length to width ratios: R:2.5, S:3.8, P:1.1, F1:0.9, F2:1.0, F3:1.8, F4:1.8, F5 + F6 (fused):3.7; ratio of each segment to length of F1: R: 1.2, S:3.2, P:1.4, F1:1.0, F2:1.2, F3:1.9, F4: 1.9, F5 + F6 (fused):3.8. F2 shorter than F3 and with 2 large, ovoid sensory/glandular structures covered by plate having 2 holes, F3 surface with striate, trapezoidal area; mesosomal sculpturing and setation and fore wing similar to those of female.

Hosts.—Aleurothrixus floccosus; Bemisia argentifolia; B. tabaci; Trialeurodes abutiloneus; T. packardi (Morill); *T. vaporariorum; Trialeurodes sp.

Distribution.—*Brazil; El Salvador; Guadeloupe; Jamaica; Mexico; Puerto Rico; USA: District of Colombia, Florida, Illinois, Louisiana, Maryland, Mississippi; Venezuela.

Specimens examined representing new host or distribution records.—Brazil, Brasilia, iv.1992, *Trialeurodes vaporariorum*, R. Oliveira (1 \degree , 2 \circlearrowright , GAE).

Biology.—Dysart (1966) reported that this species completed its life cycle on *Trialeurodes abutiloneus* in 10 to 25 days, overwintered in the whitefly pupae and darkened the puparia of its whitefly host. However, Dysart's figure of the male antenna shows the F5 and F6 separated and the F2 segment without the ovoid sensory/glandular structure; this specimen was not *E. quaintancei* and probably represents the male of a species belonging to the *Encarsia luteola* species group.

WHITEFLY-PARASITOID HOST LIST

- Aleurodicus dispersus Russell.—E. nigricephala.
- Aleurothrixus aepim (Goldi).—E. aleurothrixi.
- Aleurothrixus floccosus (Maskell), woolly whitefly.—E. aleurothrixi, E. cubensis, E. quaintancei.
- Aleurotrachelus atratus Hempel.—E. nigricephala.
- Aleurotrachelus trachoides (Back).—E. cubensis.
- Bemisia argentifolia Perring and Bellows, silverleaf whitefly.—E. nigricephala, E. quaintancei.
- Bemisia tabaci (Gennadius), sweetpotato whitefly.—E. nigricephala, E. quaintancei.
- Bemisia tabaci species-complex.—E. hamoni.
- Crenidorsum sp.—E. nigricephala.
- Dialeurodes kirkaldyi (Kotinsky), Kirkaldy whitefly.—E. nigricephala, E. quaintancei.
- *Tetraleurodes acaciae* (Quaintance).—*E. nigricephala.*
- Tetraleurodes ursorum (Cockerell).—E. hamoni.
- Trialeurodes abutiloneus (Haldeman), bandedwinged whitefly.—E. nigricephala, E. quaintancei.
- Trialeurodes floridensis (Quaintance), avocado whitefly.—E. nigricephala.
- Trialeurodes packardi (Morill), strawberry whitefly.—E. quaintancei.
- Trialeurodes vaporiorarum (Westwood), greenhouse whitefly.—E. nigricephala, E. quaintancei.

ACKNOWLEDGMENTS

We thank F.D. Bennett, H. Browning, J. Etienne, D. Schuster, C. Serra, P. Stansly and many others who collected specimens used in this study; M. Schauff for the loan of type specimens of *E. quaintancei* and *E. nigricephala;* A. B. Hamon who identified many of the host whiteflies; and J. M. Heraty and an anonymous reviewer for their

helpful comments and suggestions for this manuscript. Funding for the senior author was provided by the USDA/APHIS/National Biological Control Institute Postdoctoral Fellowship in Systematics. The junior author thanks the Keeper of Entomology and the Trustees of The Natural History Museum, London, for access to the collections in their care. Florida Agricultural Experimental Station Journal Series No. R-05819.

LITERATURE CITED

- DeBach, P. and M. Rose. 1981. A new genus and species of Aphelinidae with some synonymies, a rediagnosis of *Aspidiotiphagus* and a key to pentamerous and heteronomous Prospaltellinae (Hymenoptera: Chalcidoidea: Aphelinidae). Proceedings of the Entomological Society of Washington 83(4): 658–679.
- Dozier, H. L. 1933. Miscellaneous notes and descriptions of chalcidoid parasites (Hymenoptera). Proceedings of the Entomological Society of Washington 35(6): 85–100.
 - —. 1937. Descriptions of miscellaneous chalcidoid parasites from Puerto Rico. Journal of Agriculture of the University of Puerto Rico 21: 121– 135.
- Dysart, R. J. 1966. Natural enemies of the bandedwinged whitefly, *Trialeurodes abutilonea* (Hemiptera: Aleyrodidae) Annals of the Entomological Society of America 59: 28–33.
- Evans, G. A., M. E. Schauff, M. L. Kok-Yocomi, and R. K. Yocomi. 1995. A new species of *Aphelinus* (Hymenoptera: Aphelinidae) that parasitizes the spirea aphid, *Aphis spiraecola* Patch (Homoptera: Aphididae). Proceedings of the Entomological Society of Washington 97(1): 17–21.
- Gahan, A. B. 1931. A new species of *Encarsia* from Cuba (Hymenoptera: Aphelininae). Proceedings of the Entomological Society of Washington 33: 121–122.
- Girault, A. 1910. Synonymic and descriptive notes on the chalcidoid family Mymaridae. Journal of the New York Entomological Society 18: 233–259.
- Hayat, M. 1989. A revision of the species of *Encarsia* Foerster (Hymenoptera: Aphelinidae) from India and adjacent countries. Oriental Insects 23: 1–131.

- Howard, L. O. 1907. New genera and species of Aphelinidae with a revised table of genera. United States Department of Agriculture Technical Series 12: 69–88.
- Isidoro, N., F. Bin, S. Colazza, and S. B. Vinson. 1996. Morphology of antennal gustatory sensilla and glands in some parasitoid Hymenoptera with hypothesis on their role in sex and host recognition. Journal of Hymenoptera Research 5: 206–239.
- Pedata, P. A., N. Isidoro, and G. Viggiani. 1995. Evidence of male sex-glands on the antennae of *Encarsia asterobemisiae* Viggiani and Mazzone (Hymenoptera: Aphelinidae). Bollettino del Laboratorio di Entomologia agraria 'Filippo Silvestri' Portici 50: 271–280.
- Polaszek, A., G. A. Evans, and F. D. Bennett. 1992. *Encarsia parasitoids* of *Bemisia tabaci* (Hymenoptera: Aphelinidae, Homoptera: Aleyrodidae): a preliminary guide to identification. Bulletin of Entomological Research 82: 375–392.
- Schauff, M. E., G. A. Evans, and J. M. Heraty. 1996. A pictorial guide to the species of *Encarsia* (Hymenoptera: Aphelinidae) parasitic on whiteflies (Homoptera: Aleyrodidae) in North America. Proceedings of the Entomological Society of Washington 98(1): 1–35.
- Viggiani, G. 1986. Notes on some species of Coccophagus Westwood, Coccophagoides Girault, Encarsia Foerster and Encarsiella Hayat (Hymenoptera: Aphelinidae), mainly from the Nearctic and Neotropical regions. Bollettino del Laboratoria di Entomologia agraria 'Filippo Silvestri' Portici 43: 59–78.
- —. 1996. Descriptive notes on the males of *Encarsia nigricephala* Dozier and *Encarsia porteri* (Mercet) (Hymenoptera: Aphelinidae). Bollettino del Laboratoria di Entomologia agraria 'Filippo Silvestri' Portici 51: 101–106.
- Viggiani, G. and P. Mazzone. 1979. Contributi alla conoscenza morfo-biologica delle specie del complesso *Encarsia* Foerster-*Prospaltella* Ashmead (Hym. Aphelinidae). Bollettino del Laboratoria di Entomologia agraria 'Filippo Silvestri' Portici 35: 42–50.
 - . 1982. Antennal sensilla of some *Encarsia* Foerster (Hymenoptera: Aphelinidae), with particular reference to sensorial complexes of the male. Bollettino del Laboratoria di Entomologia agraria 'Filippo Silvestri' Portici 39: 19–26.



Evans, Gregory A. and Polaszek, Andrew. 1998. "THE ENCARSIA CUBENSIS SPECIES-GROUP (HYMENOPTERA: APHELINIDAE)." *Proceedings of the Entomological Society of Washington* 100, 222–233.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/54709</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/54491</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Smithsonian

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: Entomological Society of Washington License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.