A. mexicanus Flint, which is known from the same physiographic region some 560 miles to the southwest in El Salto, Mexico (the previous most northern generic record). The Austrotinodes larval description by Flint (1973) was based on Chilean-group species; however, our specimen, which evidently belongs to the Central American-Mexican-group species, differs significantly from the latter by the absence of teeth on the anal claws and the presence of a small but distinct tooth on the dorso-medial surface of the propleural/trochantoral process. Other features of the larva appear to agree with the general facies presented by Flint. The habitat cited above is a small, cold rapid stream along the roadside in the central "Hills" region of Texas.

R. D. Waltz and W. P. McCafferty, *Purdue University, Department of Ento*mology, West Lafayette, Indiana 47907. (Purdue Agricultural Experiment Station Journal Series No. 9191.)

> PROC. ENTOMOL. SOC. WASH. 85(1), 1983, p. 182

NOTE

On the Homonymy of *Euschistus luridus* Dallas, 1851 (Hemiptera: Pentatomidae)

In his list of Hemiptera in the British Museum, Dallas (1851. List of the Specimens of Hemipterous Insects in the Collection of the British Museum, Part 1) described two stink bugs with the specific epithet *luridus*, a North American species, *Euschistus luridus* (p. 207) and a South American species, *Agroecus luridus* (p. 200). In a recent revision of the subgenus *Lycipta* Stål of *Euschistus*, Rolston (1982. Proc. Entomol. Soc. Wash. 84: 281–296) transferred *A. luridus* into *Euschistus*, thus inadvertently creating a secondary homonym. This situation is especially confusing since both "*Euschistus luridus*" have the same author and date. Since page priority is not binding, the opportunity is now taken to make the least disruptive nomenclatural change.

Euschistus aceratos Berg (1894, An. Mus. Nac. Hist. Nat. Montev. 1: 13–17), replacement name for *Euschistus luridus* (Dallas) (1851: 200), a secondary homonym of *Euschistus luridus* Dallas (1851: 207).

The South American species has appeared seldom in the literature. It's only known synonym is chosen as a replacement name. The North American form is considered to be a subspecies of the dusky stink bug. It is retained in its trinomial, *Euschistus tristigmus luridus* Dallas. McPherson (1976. Ann. Entomol. Soc. Am. 69: 691–694; 1982. The Pentatomoidea (Hemiptera) of Northeastern North America with Emphasis on the Fauna of Illinois. South. Ill. Univ. Press, Carbondale) discusses the status of these subspecies. Rule 57e of the *International Code of Zoological Nomenclature* 1961) giving precedence to species names over subspecies names does not apply in this instance since both names were originally proposed as full species.

Donald B. Thomas, Jr., 4119 Baldwin No. 3, Lincoln, Nebraska 68504.



Thomas, Donald B. 1983. "On The Homonymy Of Euschistus luridus Dallas, 1851 (Hemiptera, Pentatomidae)." *Proceedings of the Entomological Society of Washington* 85, 182–182.

View This Item Online: <u>https://www.biodiversitylibrary.org/item/54778</u> Permalink: <u>https://www.biodiversitylibrary.org/partpdf/55010</u>

Holding Institution Smithsonian Libraries and Archives

Sponsored by Smithsonian

Copyright & Reuse Copyright Status: In copyright. Digitized with the permission of the rights holder. Rights Holder: Entomological Society of Washington License: <u>http://creativecommons.org/licenses/by-nc-sa/3.0/</u> Rights: <u>https://biodiversitylibrary.org/permissions</u>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.