

INSECT ASSOCIATES OF *GINKGO BILOBA*<sup>1</sup>A.G. Wheeler, Jr.<sup>2</sup>

**ABSTRACT:** Biological notes are given for the insect species collected on *Ginkgo biloba* trees in nurseries and ornamental plantings in Pennsylvania and the Charlotte, North Carolina area during 1971-72. The few plant-feeding insects found breeding on ginkgo are all known as polyphagous species; the predator fauna was also poorly developed. As an introduced species without any close relatives in North America, it is not surprising that ginkgo has failed to acquire an extensive fauna. However, even in its native habitat of eastern Asia, ginkgo has no known insect specific to it. Apparently no intimate herbivore-host relationship has evolved. This relative immunity to insect injury perhaps has aided the survival of this ancient tree species.

**DESCRIPTORS:** *Ginkgo biloba*, insect associates, polyphagous species, depauperate fauna, Pennsylvania, North Carolina.

Coadaptation between insects and their plant hosts has resulted in intimate plant-insect relationships and complex interactions (Ehrlich and Raven, 1965; de Wilde and Schoonhoven, 1969; Whittaker and Feeny, 1971; Root, 1973). It is common knowledge that some trees harbor large numbers of insect species, and others comparatively few (Southwood, 1960a); most vascular plants have developed a characteristic fauna and are subject to insect attack. However, *Ginkgo biloba* L., a deciduous gymnosperm and sole survivor of the family Ginkgoaceae, has a depauperate fauna and often is cited as being unusually free from insect injury (Herrick, 1935). In the words of Li (1956), ginkgo "defies the scourge of all pests." In Japan its leaves have been used as an insecticide (Franklin, 1959), but Hartzell and Wilcoxon (1941) and Heal et al. (1950) did not find evidence for insecticidal activity. Recently, Major (1967) reported that the production of 2-hexenal by damaged leaves may provide some resistance to insect attack.

Ginkgo in its native habitat of eastern Asia (Franklin, 1959) is fed on by 2 species of scale insects, a tiger moth, a silkworm, and a looper (Hase, 1955). This tree was introduced as an ornamental into the Philadelphia area during the late 18th century (Franklin, 1959), and in North America certain polyphagous insects have become occasional pests: grape mealybug, *Pseudococcus maritimus* (Ehrhorn); peach lecanium, *Lecanium persicae* (F.); white-marked tussock moth, *Hemerocampa leucostigma* (J.E. Smith); omnivorous looper, *Sabulodes caberata* (Guenée); and fruittree leafroller, *Archips argyrospilus* (Walker) (Herrick, 1935; Pirone, 1970). In addition, Dekle

---

<sup>1</sup> Accepted for publication: December 3, 1974.

<sup>2</sup> Bureau of Plant Industry, Pennsylvania Department of Agriculture, Harrisburg, PA., 17120



(1965) listed ginkgo as a host of several scale insects, and recently the zebra caterpillar, *Ceramica picta* (Harris), was reported to defoliate small trees in a nursery (Coop. Econ. Ins. Rept., 1973). Major and Tietz (1962) noted that Japanese beetles, *Popillia japonica* Newman, fed slightly on leaves of ginkgo when trees were in direct sunlight.

This paper summarizes observations of insects associated with *Ginkgo biloba* made while studying species on other ornamentals. Predators collected as adults only and other species thought to be of accidental occurrence have been omitted. Most of the observations were made on trees and seedlings in nurseries and landscape plantings in Pennsylvania during 1971-72. A few records from 1971 in the Charlotte, North Carolina area are included. All localities are in Pennsylvania except those labeled "Charlotte" or "Monroe, N.C."

## ORTHOPTERA

### Gryllidae

*Oecanthus nigricornis* Walker – Last-instar nymph 19 Aug., Fairview; last instar 25 Aug., Manchester.

## DERMAPTERA

### Forficulidae

*Forficula auricularia* L. – Adult under bark 4 Aug., Harrisburg.

## PSOCOPTERA

### Caeciliidae

*Caecilius* sp. – Nymph on dead, fungus-covered leaves 11 Aug., Charlotte, N.C.

*Graphopsocus cruciatus* (L.) – 1 ♀ taken with above.

### Ectopsocidae

*Ectopsocus meridionalis* Ribaga – 8 ♀♀, 6 nymphs on dead fungus-covered leaves 5 Aug., Manchester; 1 ♀ 12 Aug., Charlotte, N.C.

*Ectopsocopsis cryptomeriae* (Enderlein) – 1 ♀, Manchester and 1 ♂, Charlotte, N.C., same data as *Ectopsocus meridionalis*.

### Psocidae

*Blaste* sp. prob. *quieta* (Hagen) – 1 ♂ 5 Aug., Manchester.

*Metylophorus novaescotiae* (Walker) – 1 ♀ 11 Aug., Charlotte, N.C.

## HEMIPTERA – HETEROPTERA

### Pentatomidae

*Podisus maculiventris* (Say) – Eggs on leaf 22 June, Johnstown; eggs, and adult found feeding on meadow spittlebug, *Phalaenus spumarius* (L.), 5 Aug., Manchester; adult and nymph 18 Aug., Butler.



**Miridae**

*Lygus lineolaris* (Palisot de Beauvois) – 2 adults, 2 nymphs on seedlings 8 June, near Indiana; probably accidental on ginkgo.

**Anthocoridae**

*Xylocoris* sp. – Fifth-instar nymph under bark 11 Aug., Charlotte, N.C.

**HEMIPTERA – HOMOPTERA****Diaspididae**

*Lepidosaphes ulmi* (L.) – Several ♂♂ and ♀♀ 16 June, Harrisburg; eulophid wasp *Aphytis mytilaspidis* (Le Baron) reared from scales collected at same locality on 6 and 16 July. Oystershell scale is known from more than 100 tree species (Baker, 1972), but apparently has not been reported from ginkgo.

**Pseudococcidae**

*Pseudococcus maritimus* (Ehrhorn) – Nymphs in leaf axils 16 June, 26 Aug., Harrisburg. Grape mealybug is also known from more than 100 hosts (Neiswander 1949) and has been reported from ginkgo (Herrick, 1935).

**Flatidae**

*Metcalfa pruinosa* (Say) – Nymph 15 July, Manchester; adult, several cast nymphal skins 18 Aug., Butler.

**Cicadellidae**

*Graphocephala coccinea* (Forster) – 2 adults 17 Aug., Stroudsburg; adult 18 Aug., Butler; adult 20 Aug., Meadville; 7 adults 25 Aug., Manchester.

**NEUROPTERA****Chrysopidae**

*Chrysopa oculata* Say – Eggs and adult 5 Aug., Manchester.

**COLEOPTERA****Melandryidae**

*Symphora flavicollis* (Haldeman) – pupa in dead twig 13 May, Stroudsburg.

**Anthicidae**

*Anthicus scabriceps* LeConte – 2 adults in dead stem 19 Aug., Fairview.

*A. virginiae* (Casey) – 2 adults in dead stem of seedling 24 Sept., Indiana.

**Cryptophagidae**

*Toramus pulchellus* (LeConte) – Adult in dead stem of seedling 24 Sept., Indiana.

**Coccinellidae**

*Coccinella transversoguttata* Faldermann – 2 adults on seedlings 8 June, Indiana; pupa 19 Aug., Fairview.

*Coleomegilla maculata* (De Geer) – same data as *C. transversoguttata*.

*Cycloneda munda* (Say) – Adult (parasitized by *Perilitus coccinellae*) and pupa 18 Aug., Butler.



- Hippodamia convergens* Guérin – Adult 5 Aug., Manchester; pupa 19 Aug., Fairview.  
*Olla abdominalis* (Say) – 2 adults, 1 pupa 17 June, Charlotte, N.C.; 2 adults 13 Aug., Monroe, N.C.  
*Psyllobora vigintimaculata* (Say) – Pupa 12 Aug., Charlotte, N.C.

#### Phalacridae

- Acylomus ergoti* Casey – Adults on dead, fungus-covered leaves 5 Aug., Manchester.

#### Orthoperidae

- Orthoperus* sp. prob. *glaber* (Le Conte) – Adults on dead leaves 5 Aug., Manchester; larvae on dead, fungus-covered leaves 12-13 Aug., Charlotte, N.C.; adult 25 Aug., Manchester.

#### Lathridiidae

- Melanophthalma distinguenda* (Comolli) – Adults, larvae on dead, fungus-covered leaves 5 Aug., Manchester; adult 18 Aug., Butler; larvae on fungus-covered leaves 12 Aug., Charlotte, N.C.  
*M. pumila* (Le Conte) – Larvae on fungus-covered leaves 12 Aug., Charlotte, N.C.

#### Mycetophagidae

- Litargus nebulosus* Le Conte – Adult in crevice of bark 4 Aug., Harrisburg.

### LEPIDOPTERA

#### Arctiidae

- Spilosoma virginica* (F.) – Several egg masses, groups of larvae and feeding damage on leaves 19 Aug., Fairview.

#### Noctuidae

- Agrotis ipsilon* (Hufnagel) – Egg mass on leaf 28 July, Lancaster Co.; reared to maturity on ginkgo in laboratory, but in nature larvae probably would have dropped to ground to feed.  
*Ceramica picta* (Harris) – Egg mass on seedling 8 June, Indiana; reared to maturity on ginkgo in laboratory; reported from ginkgo (Coop. Econ. Ins. Rept., 1973).  
*Peridroma saucia* (Hübner) – Egg mass 8 July, State College; reared to maturity on ginkgo in laboratory.  
*Prodenia ornithogalli* Guenée – 4 larvae, feeding damage 12-13 Aug., Charlotte, N.C.

#### Lasiocampidae

- Malacosoma americanum* (F.) – Larva 27 May, State College; 2 cocoons 8 June, numerous cocoons 23 June, Indiana; 2 egg masses 22 June, Johnstown. Occurrence on ginkgo may be accidental.

#### Geometridae

- Anavitrinella pampinaria* (Guenée) – Larva feeding on leaves 13 Aug., Charlotte, N.C.  
*Biston cognataria* (Guenée) – Larva 17 June, Charlotte, N.C.; larva 18 Aug., Butler.

#### Lymantriidae

- Hemerocampa leucostigma* (J.E. Smith) – Early-stage larva 22 June, Johnstown; 1 early- and 1 late-stage larva 12-13 Aug., Charlotte, N.C.; several larvae 18 Aug., Butler;



egg masses, several groups of early-stage larvae and feeding damage 19 Aug., Fairview. The whitemarked tussock moth, a general feeder on deciduous and coniferous trees (Baker, 1972), is known to feed on ginkgo (Pirone, 1970).

### Psychidae

*Thyridopteryx ephemeraeformis* (Haworth) — 6 larvae in small bags on undersides of leaves 16-20 June, Harrisburg; 3 larvae, feeding damage 22 June, Johnstown; larva 25 June, nr. Carlisle; larvae common on small trees in nursery 15 July, Manchester; larva 3 Sept., Ephrata; 3 larvae, damage 30 Sept., nr. Carlisle. Although the bagworm is known from a wide range of hosts (Davis, 1964), it apparently has not been reported to feed on ginkgo. Larvae usually fed on the lower surfaces, leaving unsightly blotched areas.

### Lyonetiidae

*Bucculatrix* sp. — Pupae on leaves 17 June, Charlotte, N.C.; pupa 12 Aug., emerged 15 Aug., Monroe, N.C.; ca. 40 pupae 17 Aug., Stroudsburg; pupa 18 Aug., Butler. Cocoons were sometimes abundant, but the relationship to ginkgo was not determined. It is possible that nearby deciduous trees had served as hosts for this *Bucculatrix* and that merely pupation had occurred on ginkgo.

### Limacodidae

*Sibine stimulea* (Clemens) — 4-5 larvae, feeding damage 12-13 Aug., Charlotte, N.C.

### Tortricidae

*Archips argyrospilus* (Walker) — Larvae averaged 2/tree on 16 of 20 trees (8'-10') sampled at random in street planting 27 May, State College; 2 pupae and numerous pupal cases on same trees 8 July; larva rolling leaves of seedlings in nursery 8 June, Indiana. The fruittree leafroller feeds on a wide variety of hosts (Chapman and Lienk, 1971) and has been recorded from ginkgo (Tulecke and Colavito, 1966).

*A. griseus* (Robinson) — Larva 29 May, State College.

*Argyrotaenia velutinana* (Walker) — Larva on seedling 8 June, Indiana.

*Choristoneura rosaceana* (Harris) — Larva on seedling 8 June, Indiana; pupa 14 June, Montoursville; larva 18 Aug., Butler. The obliquebanded leafroller is a general feeder, although the number of its primary hosts may be limited (Chapman and Lienk, 1971).

*Platynota flavedana* Clemens — Larva on seedling 8 June, Indiana; larva 19 Aug., Fairview.

*P. idaeusalis* (Walker) — Larva 15 July, Manchester.

*Sparganothis sulfureana* (Clemens) — Larva 18 Aug., Butler. Howard (1896) reported this species from ginkgo at Washington, D.C.

### Tineidae

? Genus poss. *Opogona* sp. — 4 larvae under bark 22 Sept., Fairview.

## DIPTERA

### Phoridae

*Phalacrotophora epeirae* (Brues) — Puparia on leaf 5 Aug., Manchester; a parasite of spider eggs (Brues, 1902).

### Lonchaeidae

*Lonchaea* sp. prob. *polita* Say — Teneral adult at base of dead seedling and numerous larvae under bark of dead seedlings 8 June, Indiana.



**Chloropidae**

? *Oscinella* sp. — 2nd or early 3rd-stage larvae inside dead twigs of living trees 14 April, State College; reared in laboratory with adults emerging during 1st week of May; empty puparia of probably this species in dead twigs 11 May, Stroudsburg and 31 Aug., State College.

**Sarcophagidae**

*Sarcophaga houghi* Aldrich — Numerous larvae in old cocoons of *Malacosoma americanum* collected 23 June, Indiana.

**HYMENOPTERA****Braconidae**

*Perilitus coccinellae* (Schrank) — Emerged from coccinellid *Cycloneda munda* collected 18 Aug., Butler.

**Ichneumonidae**

*Itopectis conquisitor* (Say) — Emerged from tortricid larva collected 18 Aug., Butler.  
*Zatypota cingulata* Townes — Reared from cocoon collected 12 July, Harrisburg.

**Eulophidae**

*Aphytis mytilaspidis* (Le Baron) — Ex. oystershell scale, *L. ulmi*, collected 6 and 16 July, Harrisburg.

**Scelionidae**

*Telenomus* sp. — Ex. eggs of lepidopteran collected 18 Aug., Butler.

**Torymidae**

*Monodontomerus minor* (Ratzeburg) — Ex. ? bagworm, *T. ephemeraeformis*, collected 18 Aug., Butler. Known as a parasite of bagworm (Burks, 1967).

**DISCUSSION**

These observations on the insect associates of ginkgo support its placement among tree species most immune to insect attack (Felt, 1905; Herrick, 1935). Ginkgo does appear to be attacked rarely, and then almost exclusively by chewing insects. The only sucking species found breeding on ginkgo were oystershell scale and grape mealybug (the flatid *Metcalfa pruinosa* possibly completes its development on ginkgo). Certainly there are few other vascular plants that do not serve as host for some aphid, leafhopper, or plant bug.

Insects capable of feeding on ginkgo appear also to be generalists having a wide host plant range. Oystershell scale, grape mealybug, whitemarked tussock moth, bagworm, and fruittree leafroller all are polyphagous species.

As might be expected from the absence of a well-developed herbivore fauna, ginkgo harbors few predaceous species. Immature stages of predators



were found only rarely. Pupae of 5 coccinellid species were collected, but since larvae were never taken, development may have been completed on some other plant species.

Southwood (1960a) noted that tree species recently introduced into Britain tend to have fewer insect associates than native trees, especially those species that have been abundant throughout recent geological history. Southwood (1960b) further pointed out that introduced species that have no closely related species in the native fauna and that remain relatively rare, may have few, if any, insect associates. Since ginkgo is the sole survivor of the family Ginkgoaceae and order Ginkgoales, it is not surprising that this tree has not acquired an extensive fauna in North America. What is perhaps unusual is that no insect is known to be specific to ginkgo in its native habitat of eastern Asia. According to Hase (1955): "Wir kennen von Ginkgo keine einzige Insektenart, die daran monophag oder oligophag lebt. Auch in der Heimat in Ostasien wird Ginkgo nur von sehr polyphagen Insekten befallen." Thus no intimate herbivore-host relationship has evolved; no characteristic fauna organized around key chemicals, as in members of the Cruciferae, Umbelliferae, Solanaceae (Fraenkel, 1959), has developed.

Reasons for the survival of ginkgo are not clearly understood (Franklin, 1959), but to survive for a hundred million years, ginkgo "... must have come from a most tenacious stock ..." (Li, 1956). Perhaps its relative immunity to insect attack has aided survival. This resistance to insect injury should encourage the further use of ginkgo as an ornamental plant.

#### ACKNOWLEDGEMENTS

B.D. Burks, R.J. Gagne', J.M. Kingsolver, P.M. Marsh, C.W. Sabrosky, G.C. Steyskal, T.J. Spilman, D.M. Weisman, and W.W. Wirth of the Systematic Entomology Laboratory, USDA, Washington, D.C.; J.G. Franclemont, Cornell University; and E.L. Mockford, Illinois State University, kindly identified specimens. I thank Heinz Wolff, a Pennsylvania Department of Agriculture regional employee, for collecting the tineid larvae. I also appreciate the help of L.L. Pechuman, Cornell University, and my colleague at the Bureau of Plant Industry, T.J. Henry in improving the manuscript.

#### LITERATURE CITED

- Baker, W.L. 1972. Eastern forest insects. USDA For. Serv., Misc. Publ. No. 1175. 642 pp.
- Brues, C.T. 1902. Notes on the larvae of some Texan Diptera. *Psyche* 9: 351-354.
- Burks, B.D. 1967. Superfamily Chalcidoidea, pp. 213-282. In Krombein, K.V. and B.D. Burks. Hymenoptera of America North of Mexico, Synoptic Catalog. 2nd Supplement. U.S. Dept. Agr., Agr. Monogr. 2.
- Chapman, P.J. and S.E. Lienk. 1971. Tortricid fauna of apple in New York. N.Y. Agric. Exp. Sta. Special Publ. 122 pp.



- Cooperative Economic Insect Report. 1973. Plant Protection Division, USDA, ARS 23 (45-48): 757.
- Davis, D.R. 1964. Bagworm moths of the Western Hemisphere (Lepidoptera: Psychidae). U.S. Nat. Mus. Bull. 244: 1-233.
- Dekle, G.W. 1965. Florida armored scale insects. Arthropods of Florida and Neighboring Land Areas. 3:1-265.
- Ehrlich, P.R. and P.H. Raven. 1965. Butterflies and plants: A study in coevolution. Evolution 18: 586-608.
- Felt, E.P. 1905. Insects affecting park and woodland trees. N.Y. State Mus. Mem. 8, Vol. 1, 459 pp.
- Fraenkel, G.S. 1959. The raison d' Etre of secondary plant substances. Science 129: 1466-1470.
- Franklin, A.H. 1959. *Ginkgo biloba* L.: Historical summary and bibliography. Virginia J. Sci. 10 (3): 131-176.
- Hartzell, A. and F. Wilcoxon. 1941. A survey of plant products for insecticidal properties. Boyce Thompson Inst. Contr. 12: 127-141.
- Hase, A. 1955. Über Insektenschäden an *Ginkgo biloba* L. (Kl. Ginkgoinae). Meded. LandbHooges. OpzoekStns Gent 20: 331-336.
- Heal, R.E., E.F. Rogers, R.T. Wallace, and O. Starnes. 1950. A survey of plants for insecticidal activity. Lloydia 13: 89-162.
- Herrick, G.W. 1935. Insect enemies of shade-trees. Comstock Publishing Co., Ithaca, N.Y. 417 pp.
- Howard, L.O. 1896. The shade-tree insect problem in the eastern United States. pp. 361-384. In Yearbook U.S. Dept. Agric., 1895.
- Li, H.L. 1956. A horticultural and botanical history of ginkgo. Morris Arboretum Bull. 7(1): 3-12.
- Major, R.T. 1967. The ginkgo, the most ancient living tree. Science (Washington, D.C.) 157: 1270-73.
- Major, R.T. and H.J. Tietz. 1962. Modification of the resistance of *Ginkgo biloba* leaves to attack by Japanese beetles. J. Econ. Ent. 55: 272.
- Neiswander, R.B. 1949. The grape mealybug on *Taxus* in Ohio. J. Econ. Ent. 42: 41-44.
- Pirone, P.P. 1970. Diseases and pests of ornamental plants. 4th ed. Ronald Press, New York. 546 pp.
- Root, R.B. 1973. Organization of a plant-arthropod association in simple and diverse habitats: The fauna of collards (*Brassica oleracea*). Ecol. Monogr. 43: 95-124.
- Southwood, T.R.E. 1960 a. The number of species of insect associated with various trees. J. Anim. Ecol. 30: 1-8.
- Southwood, T.R.E. 1960 b. The abundance of the Hawaiian trees and the number of their associated insect species. Proc. Haw. Ent. Soc. 17: 299-303.
- Tulecke, W. and L. Colavito. 1966. Fruit-tree leaf rollers on plant tissue cultures and on leaves of *Ginkgo biloba*. J. Econ. Ent. 59: 1277-1278.
- Whittaker, R.H. and P.P. Feeny. 1971. Allelochemicals: Chemical interactions between species. Science 171: 757-770.
- de Wilde, J. and L.M. Schoonhoven. ed. 1969. Insect and host plant. Proc. 2nd Int. Symp. held at Wageningen, Netherlands, Ent. Exp. Appl. 12: 471-810.





Wheeler, A. G. 1975. "Insect Associates Of Ginkgo biloba." *Entomological news* 86, 37-44.

**View This Item Online:** <https://www.biodiversitylibrary.org/item/55165>

**Permalink:** <https://www.biodiversitylibrary.org/partpdf/57219>

**Holding Institution**

Smithsonian Libraries and Archives

**Sponsored by**

Smithsonian

**Copyright & Reuse**

Copyright Status: In copyright. Digitized with the permission of the rights holder.

Rights Holder: American Entomological Society

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.