

LVII.—Description of a new *Dyscophid* Frog from Yunnan.

By G. A. BOULENGER, F.R.S.

(Published by permission of the Trustees of the British Museum.)

THE family *Dyscophidæ*, with Madagascar as its headquarters, was known to be scantily represented in Burma and Siam (*Calluella*, Stol.), and Borneo (*Colpoglossus*, Blgr., *Calliglutus*, Barbour & Noble). Yunnan may now be added to the range of its distribution in Asia.

Calluella yunnanensis, sp. n.

Habit rather stout, similar to *Microhyla rubra*, Jerd. Head much broader than long; snout rounded, not projecting beyond the mouth, as long as the eye; canthus rostralis feeble; loreal region very oblique; nostril equally distant from the eye and from the tip of the snout; interorbital space as broad as the upper eyelid. Pupil round. Fingers with obtuse tips, first shorter than second, half as long as third; subarticular tubercles moderately large, feebly prominent; three carpal tubercles, median smallest. The tibio-tarsal articulation reaches the shoulder or the temple; heels feebly overlapping when the limbs are folded at right angles to the body; tibia shorter than the foot, $2\frac{1}{3}$ to $2\frac{1}{2}$ times in length from snout to vent; toes moderately long, with slightly swollen tips, the web reaching the tips, but deeply emarginate; subarticular tubercles moderately large, feebly prominent; two metatarsal tubercles, inner oval, somewhat compressed and very prominent, $\frac{3}{4}$ the length of the inner toe, outer smaller, round and flat. Skin smooth, except for some feebly raised glandular ridges which correspond with the outlines of the markings. Pale greyish or pinkish brown above, with a symmetrical vase-shaped brown marking, edged with darker and lighter, from between the eyes to the sacral region; a dark lateral band from the eye to the groin, bordered above by a series of more or less confluent black spots, which extends forwards to the canthus rostralis; limbs with dark cross-bars, one or two on the tibia; lower parts white, mottled with brown, especially on the throat.

From snout to vent 29 mm.

Two specimens from Yunnan Fou, from the collection of Mr. J. Graham.



Boulenger, George Albert. 1919. "LVII.—Description of a new Dyscophid frog from Yunnan." *The Annals and magazine of natural history; zoology, botany, and geology* 3, 548–548. <https://doi.org/10.1080/00222931908673856>.

View This Item Online: <https://www.biodiversitylibrary.org/item/61791>

DOI: <https://doi.org/10.1080/00222931908673856>

Permalink: <https://www.biodiversitylibrary.org/partpdf/58280>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.