

Fig Wasp (Hymenoptera: Chalcidoidea: Agaonidae, Pteromalidae, Eurytomidae and Ormyridae) and *Ficus* (Moraceae) Species Richness and Biogeography of Monts Doudou in Southwestern Gabon

Simon van Noort

Natural History Division, South African Museum (Iziko Museums of Cape Town),
P.O. Box 61, Cape Town, 8000, South Africa
email: svannoort@iziko.org.za

Thirty species of Chalcidoidea were reared from figs of three *Ficus* species (*F. sansibarica macrosperma*, *F. louisii*, and *F. elasticoides*) that were sampled on Monts Doudou, of which 28 are new records for the country, more than doubling the number of species previously known from Gabon, all of which were recorded from the vicinity of Makokou. Even though the majority of the fig wasp species sampled on Monts Doudou are currently only known from Gabon, their host fig tree distribution suggests that they will be more widespread through the Congo basin or even further afield. Of the 105 Afrotropical fig tree species, 47 are recorded from, or predicted to occur, in Gabon, illustrating the rich representation of *Ficus* species present in the country. These fig tree species are listed, together with the known fig wasps associated with each host tree. Based on current data, a conservative species richness of 225 fig wasps is associated with *Ficus* species in Gabon. The expected species richness, however, is likely to be closer to 400–500 species. Twenty to thirty-five fig tree species are predicted to occur on Monts Doudou, which from a *Ficus* perspective does not have sufficient elevation to support elements of Pleistocene refugia. The *Ficus* species predicted to occur on Monts Doudou are expected to have an associated fig wasp faunal richness of around 200–350 species.

RÉSUMÉ

Trente espèces de Chalcidoidea vivant sur trois espèces de figuiers (*Ficus sansibarica macrosperma*, *F. louisii* et *F. elasticoides*) ont été inventoriées au Mont Doudou; 28 de ces espèces sont nouvellement découvertes doublant le nombre d'espèces connues au Gabon. Ces espèces ont seulement été récoltées aux alentours de Makokou. Même si, aujourd'hui, la majorité des guêpes de figuiers récoltées au Mont Doudou sont seulement connues du Gabon, la large distribution des figuiers suggère que ces guêpes pourraient également être présentes jusqu'au bassin du Congo, voire plus loin.

Sur les 105 figuiers afrotropicaux, 47 sont connus ou supposés être présents au Gabon, illustrant une riche représentation d'espèces de figuiers dans le pays. Ces espèces de figuiers sont ici listées avec les guêpes qui leur sont associées. À partir des données actuelles, la diversité minimale de guêpes associées aux figuiers du Gabon devrait être de 225 espèces, mais le nombre pourrait atteindre 400 à 500 espèces. Vingt à trente-cinq espèces de figuiers devraient se trouver au Mont Doudou qui, par la perspective des *Ficus*, n'a pas une élévation suffisante pour supporter des éléments de refuges du Pléistocène. Les

espèces de *Ficus* qui sont supposées se trouver au Mont Doudou devraient présenter environ 200 à 350 espèces de guêpes de figuiers associées.

INTRODUCTION

The relationship between pollinating fig wasps (Chalcidoidea, Agaonidae) and their host fig trees (*Ficus*, Moraceae) is a classic example of an obligate mutualism, where neither partner can reproduce without the other, the wasp providing a pollination service and the fig tree in turn providing a breeding site for the pollinating wasp's progeny (Galil 1977; Janzen 1979). Besides the pollinator, each fig tree species has a diverse assemblage of non-pollinating fig wasps associated with it. The non-pollinating wasps are either phytophagous, galling the ovules as do the pollinators, or parasitoids of the gall formers (Compton and van Noort 1992; West and Herre 1994; West et al. 1996; Kerdelhué and Rasplus 1996a, 1996b). The mutualism between pollinating fig wasps and fig trees is usually a one-to-one relationship (Ramirez 1970; Wiebes 1979a; Wiebes and Compton 1990; Rasplus 1996) with each fig wasp species associated with a particular host fig tree species, of which there are approximately 750 worldwide (Berg 1989). There are, however, a few exceptions to this rule. A number of cases have been documented where two pollinator species are associated with a single host species (Compton et al. 1991; Michaloud et al. 1985; Michaloud et al. 1996; Berg and Wiebes 1992). Non-pollinating wasps appear to be almost as host-specific as the pollinators (Ulenberg 1985; van Noort 1994c).

In contrast to Bouček's (1988) classification, in which fig wasps were united under the single family Agaonidae, recent molecular investigations have shown that the Agaonidae is paraphyletic. Three of the constituent subfamilies of non-pollinating fig wasps (Sycoecinae, Otitesellinae, and Sycoryctinae) were reassigned to the Pteromalidae, leaving the pollinating fig wasps in the Agaonidae (Rasplus et al. 1998). The true taxonomic affinities of the Sycophaginae and Epichrysomallinae remain undecided, but they do not belong in the Agaonidae. In addition, there are representatives of the Eurytomidae and Ormyridae that are also associated with figs in Africa.

Fig wasps are circumtropical in distribution, with about 230 described species of an estimated 700–1000 species in the Afrotropical region (van Noort and Rasplus 1997; van Noort unpublished). Fig wasps are exclusively associated with figs and since *Ficus* species richness is centered in the tropics, Gabon has a rich representation of species, with a total of 47 fig tree species recorded from, or predicted to, occur in the country (Berg et al. 1984). This high diversity is in line with other tropical African countries: Cameroon (58 species) (Berg et al. 1985), Tanzania (39), Kenya (34) and Uganda (43) (Berg and Hijman 1989). Given the pan-tropical distribution of *Ficus*, it is predictable that temperate countries such as South Africa support a lower diversity (22 species) (Berg 1990).

Local *Ficus* species richness is extremely high in Gabon. In the Lopé Reserve in central Gabon, which lies approximately 250 km northwest of Monts Doudou, at least 18 species of *Ficus* are present (White and Abernethy 1997). Thirty-four species have been recorded from the vicinity of Makokou in northeastern Gabon (Berg et al. 1984), with 28 hemi-epiphytic *Ficus* species recorded around the Institut de Recherches en Ecologie Tropicale station at M'Passa, situated 10 km south of Makokou (Nieder et al. 2001). Very few assessments of local fig wasp species richness have been conducted in Africa. During an ecological inventory survey of the fauna and flora occurring in Mkomazi Game Reserve in northeastern Tanzania (Coe et al. 1999), an assessment of fig tree and fig wasp species richness of this semi-arid east African savanna region was carried out (van Noort and

Compton 1999). Eighty-five species of fig wasp, of an expected species richness of 170, were reared from nine host fig tree species. For comparative purposes, van Noort and Compton (1999) demarcated an area in Kwazulu-Natal (South Africa) that was similar in size to Mkomazi Game Reserve, where 90 fig wasp species of a potential 117 species were recorded from 12 host fig tree species.

Research on fig trees and fig wasps in Gabon has centered on ecological and evolutionary investigations. These were conducted on *Ficus* species and their associated fig wasp assemblages in the vicinity of the Institut de Recherches en Ecologie Tropicale station at M'Passa, situated 10 km south of Makokou in northeastern Gabon (Gautier-Hion and Michaloud 1989; Michaloud 1982; 1988; Michaloud et al. 1985; Michaloud et al. 1996; Nieder et al. 2001). A number of fig wasp species have been described from collections made by Georges Michaloud and others in the vicinity of Makokou (Wiebes 1979b, 1979c, 1986, 1988, 1989; Michaloud et al. 1985; van Noort 1993a, 1993b, 1994a, 1994b, 1994c). Prior to the collections made on Monts Doudou only 26 described species of fig wasp had been recorded from Gabon (Berg and Wiebes 1992; van Noort, unpublished, see also web site at: <http://www.museums.org.za/sam/collect/life/ento/simon/figwasp.htm>), which is simply a function of a lack of collecting fig wasps in the region. Within Gabon all these species are only known from the vicinity of Makokou and for a number of them Makokou is the type (and currently only known) locality.

In this paper the sampled species richness of fig wasps on Monts Doudou is reported on and a summary of expected *Ficus* and fig wasp species richness is provided for Gabon and Monts Doudou. The biogeographical affinities of Gabon's fig wasps and fig trees are discussed.

MATERIALS AND METHODS

Study site

Monts Doudou is situated in Province Ogoové-Maritime in southwestern Gabon and straddles two reserves: Réserve de la Moukalaba-Dougoua and Réserve des Monts Doudou. Fig wasps were collected at three localities: 2°17.00'S, 10°29.83'E (110 m); 2°12.36'S, 10°25.11'E (110 m); 2°13.63'S, 10°23.67'E (660 m). The habitat is defined as coastal lowland rain forest (White 1983).

Sampling methods

Fig wasps were collected on an opportunistic basis. The majority of fig trees that were located were stranglers in other tree species, and the fig crops were located in the forest canopy 30–50 m above the ground. Accessing the forest canopy could not be accommodated in the logistics of the survey and sampling of fig crops relied on locating figs that had been knocked to the forest floor. This was facilitated by the activities of primates feeding on ripe figs and by adverse weather conditions. One of the collections was made possible through a severe thunderstorm that knocked many figs to the forest floor. Figs were placed in a plastic jar with a gauze lid. Fig wasps that emerged were aspirated and preserved in 96% ethanol. After a couple of days the figs were split open and any fig wasps remaining within the fig cavity were extracted and preserved.

Identification and analyses

Specimens were identified to family and subfamily, and where possible, to species. However, the majority of the species are undescribed and hence were sorted to

TABLE 1. Fig wasps reared from the *Ficus* species that were sampled on Monts Doudou. Family placement of the Epichrysomallinae is currently unresolved.

<i>Ficus</i> sp.	Agaoonidae	Pteromalidae Sycocinae	Pteromalidae Otitellinae	Pteromalidae Sycoryctinae	Family ? Epichrysomallinae	Eurytomidae	Ormyridae
<i>F. sansibarica</i> <i>macrosperma</i>	<i>Courtella armata</i>	<i>Seres a. armipes</i>	<i>Otitella</i> sp. 1	<i>Sycoscapteridea</i> sp. <i>Sycoryctes</i> sp. 1 <i>Sycoryctes</i> sp. 2 <i>Sycoryctes</i> sp. 3 <i>Watshamiella</i> sp. 1 <i>Watshamiella</i> sp. 2	<i>Camarothonax</i> sp. 1	<i>Sycophila</i> sp. 1 <i>Sycophila</i> sp. 2 <i>Sycophila</i> sp. 3	
<i>F. louisii</i>	<i>Paragaon josephi</i>	<i>Crossogaster</i> sp.	<i>Otitella</i> sp. 2 <i>Otitella</i> sp. 3 <i>Otitella</i> sp. 4 <i>Philosycus</i> sp.	<i>Sycoryctes</i> sp. 4 <i>Sycoryctes</i> sp. 5 <i>Sycoryctes</i> sp. 6 <i>Sycoryctes</i> sp. 7 <i>Watshamiella</i> sp. 3	<i>Camarothonax</i> sp. 2 <i>Camarothonax</i> sp. 3 <i>Camarothonax</i> sp. 4	<i>Sycophila</i> sp. 4	<i>Ormyrus</i> sp.
<i>F. elasticoides</i>	<i>Elisabethiella articulata</i>						

morpho-species. Owing to the limited number of samples it was not possible to carry out species richness analyses or to assess the effect of elevation on patterns of species richness. The collected material is deposited in the South African Museum, Cape Town. Representative specimens will be returned to Libreville, Gabon.

RESULTS

Thirty fig wasp species, representing 12 genera and seven higher taxa at family or subfamily level, were reared from three species of fig tree: *Ficus sansibarica macrosperma* (Mildbr. and Burret) C. C. Berg; *Ficus louisii* Bout. and Leon; and *Ficus elasticoides* De Wild. The fig wasp faunal assemblages reared from these three host fig tree species are listed in Table 1. The forty-seven *Ficus* species that occur, or are predicted to occur, in Gabon, currently have a predicted associated fig wasp faunal assemblage of 225 species (Appendix A).

DISCUSSION

Species richness

Monts Doudou is only the second locality in Gabon that has been sampled for fig wasps. This survey added a further 28 species to the previously recorded total of 26 described species from the vicinity of Makokou in northeastern Gabon. Based on this total of 54 species in conjunction with the fig wasp species richness that has been recorded elsewhere in Africa from those *Ficus* species that occur in Gabon, at least 225 species of fig wasp are predicted to occur in this central African country (Appendix A). Seven *Ficus* species that occur, or are predicted to occur, in Gabon have never had their associated fig wasp faunas reared, and as such their pollinators and non-pollinating fig wasp faunas are unknown (Appendix A). In addition, the majority of the 47 *Ficus* species that occur in Gabon have not had their full assemblage of fig wasps reared from them any-

where within their distributional range in Africa. As such the estimate of 225 fig wasp species associated with Gabon's fig species is extremely conservative. As with many invertebrate taxa, the lack of alpha-level taxonomic and biogeographical knowledge of fig wasps is a function of insufficient sampling. The paucity of distributional information in Gabon is indicative of most of the rest of Africa, where there are still vast areas where no collecting of fig wasps has been undertaken. Collecting of fig wasps is hindered by the need to rear specimens from their host figs, a procedure which is constrained by the ecology of the mutualism. Fig crops are produced randomly throughout the year and individual trees produce crops at different times to each other, both essential traits to ensure the continued cycling of the mutualism (Bronstein, 1992). Because of this, most of the fig trees that are located during field surveys either have no figs or have figs at the wrong stage of development for rearing of fig wasps. On average only one out of every 30 trees has a fig crop at the right stage of development (pers. observ.). Furthermore, not every fig wasp species associated with a particular fig tree species is reared from every sample of figs. There are two reasons for this. First, not every fig crop borne by the tree has all the possible fig wasp species present, due to either the cycling of the mutualism, where the full fig wasp species complement does not locate every potential fig crop, or alternatively some fig wasp species may be absent from the local geographical area. Second, it is not practical to sample every fig in a particular crop. Since not all the fig wasp species associated with a particular fig crop will be breeding in every fig, rare species may be missed. To fully sample the fig wasp species assemblage associated with a fig tree species may require up to 23 collections from different trees of that species in a particular geographical area (Compton and Hawkins 1992; Compton et al. 1994; West et al. 1996). This is illustrated by an example from the New World where a single crop of *F. aurea* in Florida (USA) produced four species of fig wasp, a sample of 12 crops over time from the same tree produced seven species, and a sample of 60 crops from 23 different trees produced a total of nine species (Bronstein and Hossaert-McKey, 1996). Thus, the single samples collected from Monts Doudou *Ficus* are insufficient to assess fig wasp species richness associated with these host species.

The total of three *Ficus* species recorded for Monts Doudou is likely to be a small fraction of the total number of species occurring there. Based on assessments of local *Ficus* species richness in other areas of Gabon, such as a richness of at least 18 species of *Ficus* in the Lopé Reserve (White and Abernethy 1997) and 34 species within the vicinity of Makokou (Berg et al. 1984), around twenty to thirty-five fig tree species are predicted to occur on Monts Doudou, with an estimated associated fig wasp faunal richness of roughly 200–350 species. The estimation of fig wasp richness is based on known species richness of fig wasp assemblages associated with well-sampled fig tree species and the actual richness may be as much as three times greater, since up to 31 species of fig wasp have been recorded as being associated with a single *Ficus* species (van Noort and Compton 1999; J. Y. Rasplus pers. comm.). This, however, is tempered by the fact that the degree of host-specificity has not been determined for all of the non-pollinating groups of fig wasps, although the groups that have been revised so far show a high degree of specificity (Ulenberg 1985; van Noort 1994c). As such, a hypothesis that 10 wasp species are specific to each fig tree species may be a more realistic interpretation, albeit an extremely conservative one.

Biogeography

Ficus endemism is low in Gabon with no species restricted to the country. However, three species are sub-endemic with a distribution encompassing Gabon and Cameroon: *F.*

abscondita C. C. Berg, *F. subsagittifolia* C. C. Berg, and *F. cyathistipula pringsheimiana* (Braun and K. Shum.) C. C. Berg (Berg et al. 1984). The nominate subspecies of *F. cyathistipula* Warberg has a wider distribution from Ivory Coast to Angola, N. Zambia, Kenya and Tanzania, but has a different species of pollinating wasp and a different assemblage of non-pollinating wasps, suggesting that these two subspecific taxa are distinct at species level. A further six Gabon species are restricted to the central African region. Eighteen of the Gabon species (plus a further two species that are predicted to occur in Gabon) are distributed through central and west Africa. The remaining Gabon species enjoy a more widespread distribution extending into east and southern Africa. Given this fairly widespread distribution of the *Ficus* species that occur in Gabon and the fact that all three of the sub-endemic Gabon species are lowland forest taxa, Monts Doudou is unlikely to have any endemic species. From a biogeographical perspective Monts Doudou does not have a high enough altitude to support (sub)montane *Ficus* species, and although the mountain has been postulated to be a Pleistocene refuge based on the presence of a number of *Begonia* species that occur towards the peak (Sosef 1994), the presence of *F. louisii*, a lowland forest species, on the peak (660 m) of Monts Doudou suggests that this elevation is within the climatic tolerance of widespread lowland *Ficus* species. Species with distributions conforming to Pleistocene refugia, such as *Ficus oreodryadum* Mildbr., which is predicted to occur in Gabon and is distributed from Uganda, Rwanda, Burundi to Cameroon, SE Nigeria and Fernando Po, are restricted to montane and submontane forest between 1300–2500 m (Berg et al. 1984), and the only likely locality that this species may be present in Gabon is on Mont Cristal. Given the obligate mutualism between fig wasps and fig trees, the above discussion concerning endemism and Pleistocene refugia will hold for fig wasps as well. However, this assumption needs to be treated with caution as not all fig wasp species are distributed throughout the range of their host fig tree species (Compton et al. 1994).

Biogeographical affinities of Monts Doudou fig wasps could only be ascertained for the four described species that were collected. *Courtella armata*, the pollinator of *Ficus sansibarica*, has been reared from both subspecies of this host, the nominate subspecies and *F. sansibarica macrosperma*, in Ivory Coast, Nigeria, Cameroon, Zambia, Zimbabwe and South Africa (Wiebes 1986; Berg and Wiebes 1992; van Noort, unpub.). This is the first published record of this species from Gabon. *Ficus sansibarica* is a widespread Afrotropical species distributed from eastern South Africa and northern Angola north to Kenya and Uganda and west to Guinea-Bissau (Berg and Wiebes 1992), and the pollinator would be expected to exhibit the same distributional pattern. *Seres armipes*, a non-pollinating sycoecine fig wasp reared from *F. s. macrosperma* on Monts Doudou, was described from specimens reared from *Ficus ovata*, which is pollinated by *Courtella hamifera* Kieffer, and has been recorded from Ivory Coast, Cameroon and Uganda associated with *Ficus ovata*. *Seres armipes* has previously been recorded from *Ficus sansibarica macrosperma* in Ivory Coast and Zambia (van Noort 1993a), but this is the first record of this species from Gabon. *Paragaon josephi*, the pollinator of *Ficus louisii*, is only known from Gabon having been described from specimens reared from *F. louisii* at Makokou (Wiebes 1986). However, the species is likely to be more widespread as *F. louisii* is a central and west African species recorded from Democratic Republic of the Congo, Gabon, Cameroon and Guinea (Berg and Wiebes 1992). DNA has been successfully extracted from the samples of *Paragaon josephi* Wiebes collected on Monts Doudou as part of an investigation into the phylogeny and evolution of fig-pollinating wasps at generic level (J. M. Cook, V. Bull, and C. Lopez Vaamonde in prep.). The *Crossogaster* species that was reared

from this fig tree is undescribed. This sycoecine genus was recently revised (van Noort 1994a) and the species associated with *F. louisii* is distinct from the 16 described *Crossogaster* species. *Elisabethiella articulata*, the pollinator of *F. elasticoides*, has previously been reared from this host species in Gabon (Makokou) (Wiebes 1986), and has also been collected in a Berlese trap (Equatorial Guinea) and at light (Ivory Coast) (Berg and Wiebes 1992). *Ficus elasticoides* is a lowland rain forest species distributed from Angola to Cameroon and is also present in Ivory Coast (Berg and Wiebes 1992). The remainder of the fig wasps that were reared from the 3 sampled fig tree species are likely to be undescribed species, but this is difficult to ascertain without the revision of the groups in question having been completed (van Noort, in prep.).

In summary, the sampling of fig wasps during the survey of Monts Doudou was done on an opportunistic basis. Consequently, it is likely that many further species of *Ficus* still wait to be recorded, each with its own host-specific fig wasp fauna. In addition, the fig wasp species richness recorded from the three sampled fig tree species on Monts Doudou is currently an under-representation, given the limited sampling effort. For both these reasons, the results are a gross underestimation of fig wasp species richness in this locality. Nevertheless, indications are that Gabon supports a rich *Ficus* and fig wasp fauna, a factor that will likely hold for Monts Doudou as well.

ACKNOWLEDGMENTS

I would like to thank all of the staff, in particular Olivier Langrand, Prosper Obame Ondo, Andre Kamdem Toham, the late Faustin Oyono, and Jean-Jacques Tanga of WWF-Gabon in Libreville for logistical support and assistance in the field. Collecting and export permits were kindly granted by Emile Mamfoumbi Kombila, Directeur de la Faune et de la Chasse, Libreville. Thanks to Georges Michaloud and a referee for improving the manuscript. Fieldwork was funded by the World Bank through WWF-US, and the processing and identification of the samples by the National Research Foundation (South Africa).

APPENDIX A

Checklist of fig tree species recorded from Gabon or predicted to occur there (after Berg et al. 1984; nomenclature updated), and including a list of the predicted associated fig wasp fauna for each *Ficus* species based on collections made elsewhere in Africa. The fig wasp species that have been recorded (both previously and during this survey) from Gabon are indicated in bold text. Habitat associations of fig tree species are after Berg and Wiebes (1992). Distributions of the fig tree species are summarised as Southern Africa (S); East Africa (E); Central Africa (C) and West Africa (W).

FICUS OF GABON

SYCIDIUM (subgenus)

SYCIDIUM (section)

1. *F. exasperata* Vahl (S, E, C, W AFRICA) Forest up to 1800 m
Kradibia gestroi afrum (Wiebes)
Sycoscapteridea longipalpus (Joseph)
Philotrypesis quadrisetosa (Westwood)
Sycoryctes sp.

- 2a. *F. asperifolia* Miq. (E, C, W AFRICA) Wet and dry forest along streams
Kradibia gestroi afrum (Wiebes)

- 2b. *F. asperifolia* Miq. ("variety" *urceolaris*)
Kradibia hilli Wiebes

SYCOMORUS (subgenus)

3. *F. mucoso* Ficalho (E, C, W AFRICA) Forest

Ceratosolen arabicus Mayr

Ceratosolen galili Wiebes

Apocrypta crypta Ulenberg

Apocryptophagus sp. A

Apocryptophagus sp. B

4. *F. sur* Forsk. (S, E, C, W AFRICA) Woodland and forest

Ceratosolen silvestrianus Grandi

Ceratosolen flabellatus Grandi

Sycophaga cyclostigma Wiebes

Sycophaga silvestrii Grandi

Apocryptophagus sp. A

Apocryptophagus sp. B

Apocryptophagus sp. C

Apocryptophagus sp. D

Apocrypta guineensis Grandi

Apocrypta sp. B

Sycoscapter nigrum Risbec

Sycoscapter sp. B

Watshamiella sp. A

Watshamiella sp. B

Watshamiella sp. C

Camarothorax sp.

Sycophila sp.

Acophila sp.

5. *F. vogeliana* (Miq.) Miq. (E, C, W AFRICA) Riverine and swamp forest

Ceratosolen acutatus Grandi

Sycophaga sp.

Apocryptophagus sp. A

Apocryptophagus sp. B

Apocrypta sp.

Watshamiella sp.

PHARMACOSYCEA (subgenus)

OREOSYCEA (section)

6. *F. variifolia* Warb. (E, C, W AFRICA) Wet and dry forest up to 1300 m

Dolichoris flabellata Wiebes

7. *F. dicranostyla* Mildbr. (E, C, W AFRICA) Savanna woodland up to 1500 m

Dolichoris flabellata Wiebes

UROSTIGMA (subgenus)

UROSTIGMA (section)

8. *F. ingens* (Miq.) Miq. (S, E, C, W AFRICA) (Predicted to occur in Gabon) Woodland

Platyscapa soraria Wiebes

Otitesella rotunda van Noort

Otitesella longicauda van Noort

Sycoryctes sp. A

Sycoryctes sp. B

Sycoryctes sp. C

Philotrypesis sp.

GALOGLYCHIA (section)

GALOGLYCHIA (subsection)

9. *F. lutea* Vahl (S, E, C, W AFRICA) Forest up to 1800 m

Allotriozone heterandromorphum Grandi

Philocaenus silvestrii (Grandi)

Sycoryctes sp. A

Sycoryctes sp. B

Sycoscapter sp.

Watshamiella sp.

Philotrypesis selenetica Grandi

Philotrypesis sp.

Otitesella africana Grandi

Otitesella sp. B

Camarothorax sp. A

Camarothorax sp. B

Sycophila sp. A

Sycophila sp. B

Sycophila sp. C

Sycophila sp. D

Sycophila sp. E

PLATYPHYLLAE (subsection)

10. *F. recurvata* De Wildeman (C, W AFRICA) Forest at low altitudes

11. *F. jansii* Boutique (C AFRICA) (Predicted to occur in Gabon) Forest up to 1500 m

12. *F. trichopoda* Baker (S, E, C, W AFRICA)

Elisabethiella bergi Wiebes

Crossogaster robertsoni van Noort

Philocaenus hippopotomus van Noort

Sycoryctes sp. A

Sycoryctes sp. B

Sycoscapter sp. A

Sycoscapter sp. B

Philotrypesis sp.

Otitesella sp.
Watshamiella sp. A
Watshamiella sp. B
Camarothorax sp.

CHLAMYDODORAE (subsection)

13. *F. calyprata* Vahl (= *F. mallotoides*) (C, W AFRICA) Forest up to 700 m
Elisabethiella pectinata (Joseph)
14. *F. amadiensis* De Wild (E, C AFRICA) (Predicted to occur in Gabon) Woodland up to 2100 m
15. *F. craterostoma* Mildbr. and Burret (S, E, C, W AFRICA) Forest up to 2100 m
Alfonsiella michaloudi Wiebes
Philocaenus liodontus (Wiebes)
Philocaenus insolitus van Noort
Sycoryctes sp.
Otitesella sp.
16. *F. lingua lingua* De Wild. and T. Durand (E, C, W AFRICA) Forest up to 1200 m
Alfonsiella michaloudi Wiebes
Philocaenus liodontus (Wiebes)
17. *F. natalensis lepieurii* (Miq.) C. C. Berg (C, W AFRICA) Forest, woodland up to 1200 m
Alfonsiella fimbriata Waterston
Philocaenus liodontus (Wiebes)
18. *F. thonningii* Bl. (S, E, C, W AFRICA) Wet and dry forest up to 2000 m
Elisabethiella stuckenbergi Grandi
Alfonsiella brongersmai Wiebes
Alfonsiella longiscapa Joseph
Philocaenus barbarus (Grandi)
Crossogaster odorans Wiebes
Philotrypesis parca Wiebes
Sycoscapter cornutus Wiebes
Sycoryctes remus Wiebes
Sycoryctes hirtus Wiebes
Watshamiella alata Wiebes
Watshamiella sp. A
Watshamiella sp. B
Otitesella tsamvi Wiebes
Otitesella sp. B
Camarothorax brevimucro Bouček
Camarothorax equicollis Bouček
Camarothorax longimucro Bouček
Sycotetra serricornis Bouček
Eurytoma ficusgallae Bouček

Ficomila curtivena Bouček
Ficomila gambiensis (Risbec)
Syceurytoma ficus Bouček
Sycophila flaviclava Bouček
Sycophila kestraneura (Masi)
Sycophila modesta Bouček
Sycophila naso Bouček
Sycophila punctum Bouček
Sycophila sessilis Bouček
Ormyrus flavipes Bouček
Ormyrus subconicus Bouček
Ormyrus watshami Bouček

19. *F. kamerunensis* Mildbr. and Burret (C, W AFRICA) Forest at low altitudes
Alfonsiella fimbriata Waterston
Philocaenus liodontus (Wiebes)

CRASSICOSTAE (subsection)

20. *F. elasticoides* De Wild. (C, W AFRICA) Forest at low altitudes
Elisabethiella articulata (Joseph)
Otitesella sp. A
Philosycus sp.
Sycoryctes sp. A
Sycoryctes sp. B
Watshamiella sp.
Camarothorax sp.
21. *F. burretiana* Hutch. (C AFRICA) Forest at low altitudes
Elisabethiella longiscapa Wiebes
22. *F. oreodryadum* Mildbr. (C, W AFRICA) (Predicted to occur in Gabon) (Sub)montane forest 1300–2500 m
23. *F. pseudomangifera* Hutch. (C, W AFRICA) Forest up to 1200 m
Nigeriella letouzeyi Wiebes
24. *F. adolfi-friderici* Mildbr. (C, W AFRICA) Forest at low altitudes
25. *F. louisii* Boutique and J. Léonard (C, W AFRICA) Forest at low altitudes
Paragaon josephi Wiebes
Crossogaster sp.
Otitesella sp. A
Otitesella sp. B
Sycoryctes sp. A
Sycoryctes sp. B
Camarothorax sp. A
Camorothorax sp. B
Sycophila sp.

***Ormyrus* sp.**

26. *F. leonensis* Hutch. (C, W AFRICA) (Predicted to occur in Gabon) Forest

CYATHISTIPULAE (subsection)

27. *F. conraui* Warb. (C, W AFRICA) Forest

Agaon kiellandi Wiebes

28. *F. tessellata* Warb. (C, W AFRICA) Forest, often riverine, up to 1900 m

Agaon taiense Wiebes

Sycoecus bergi van Noort

- 29a. *F. ardisioides ardisioides* Warb. (C AFRICA) (Predicted to occur in Gabon) Forest at low altitudes

- 29b. *F. ardisioides camptoneura* (Mildbr.) C. C. Berg (C, W AFRICA) Forest at low altitudes

Agaon megalopon Wiebes

Philotrypesis sp.

30. *F. preussii* Warb. (C, W AFRICA) Forest up to 1200 m

Agaon sp.

31. *F. abscondita* C. C. Berg (C AFRICA) Forest at low altitudes

Agaon sp.

- 32a. *F. cyathistipula cyathistipula* Warb. (E, C, W AFRICA) Forest, often riverine up to 1800 m

***Agaon fasciatum* Waterston**

Sycoecus thaumastocnema Waterston

- 32b. *F. cyathistipula pringsheimiana* (Braun and K. Shum.) C. C. Berg (C AFRICA) Forest at low altitudes

***Agaon kiellandi* Wiebes**

Sycoecus oculabulbus van Noort

Sycoryctes sp.

33. *F. cyathistipuloides* De Wild. (C, W AFRICA) Forest at low altitudes

Agaon obtusum Wiebes

Sycoecus wiebesi van Noort

Watshamiella sp. A

Watshamiella sp. B

Watshamiella sp. C

Philotrypesis sp.

34. *F. densistipulata* De Wild. (C AFRICA) Forest up to 1250 m

***Agaon kiellandi* Wiebes**

35. *F. subcostata* De Wild. (C AFRICA) Forest at low altitudes

***Agaon acutatum* Wiebes**

36. *F. barteri* Sprague (C, W AFRICA) Forest up to 1200 m

***Agaon paradoxum* Dalman**

37. *F. subsagittifolia* C. C. Berg (C AFRICA) Forest at low altitudes

Agaon cicatriferens multum* Wiebes**Sycoecus wiebesi* van Noort**

38. *F. wildemaniana* De Wild. and T. Durand (C AFRICA) Forest up to 1100 m

Agaon gabonense* Wiebes*CAULOCARPAE (subsection)**

- 39a. *F. ottoniifolia ottoniifolia* (Miq.) Miq. (C, W AFRICA) Open forest up to 1600 m

Courtella camerunensis* (Wiebes)**Courtella gabonensis* Wiebes**

- 39b. *F. ottoniifolia lucanda* (Ficalho) C. C. Berg (C AFRICA) Wet to dry forest up to 1500 m

Courtella scobinifera (Waterston)*Philocaenus levis* (Waterston)

40. *F. tremula kimuenzensis* (Warb) C. C. Berg (C, W AFRICA) Forest at low altitudes

41. *F. artocarpoides* Warb. (C, W AFRICA) Forest up to 1200 m

Courtella penicula* (Wiebes)**Courtella hladikae* (Wiebes)*****Seres longicalcar* van Noort*****Crossogaster michaloudi* van Noort*****Philocaenus rasplusi* van Noort**

42. *F. polita polita* Vahl (S, E, C, W AFRICA) Forest up to 1200 m

Courtella bekiliensis (Risbec)*Seres solweziensis* van Noort*Sycoryctes* sp. A*Sycoryctes* sp. B*Sycoscapter* sp.*Sycoscapteridea* sp.*Watshamiella* sp.*Philotrypesis* sp.*Otitesella* sp.*Camarothorax* sp.*Sycophila* sp. A*Sycophila* sp. B

43. *F. sansibarica macrosperma* (Mildbr. and Burret) (C, W AFRICA) Forest and woodland up to 1200 m

Courtella armata* (Wiebes)**Seres armipes* Waterston**

Seres solweziensis van Noort

Otitesella sp.

Sycoryctes sp. A

Sycoryctes sp. B

Sycoryctes sp. C

Sycoscapteridea sp.

Sycoscapter sp.

Watshamiella sp. A

Watshamiella sp. B

Camarothorax sp.

Sycophila sp. A

Sycophila sp. B

Sycophila sp. C

44. *F. dryepondtiana* De Wild. (C AFRICA) Forest at low altitudes

Courtella sylviae Wiebes

45. *F. umbellata* Vahl (C, W AFRICA) Forest and woodland at low altitudes

Courtella medleri (Wiebes)

46. *F. bubu* Warb. (S, E, C, W AFRICA) Forest and woodland up to 1200 m

Courtella michaloudi (Wiebes)

Seres wardi van Noort

Sycoryctes sp. A

Sycoryctes sp. B

Sycoryctes sp. C

Sycoryctes sp. D

Sycoryctes sp. E

Sycoryctes sp. F

Sycoscapter sp.

Sycoscapteridea sp.

Philotrypesis sp.

Watshamiella sp. A

Watshamiella sp. B

Otitesella sp.

Philosycus sp.

Camarothorax sp. A

Camarothorax sp. B

Camarothorax sp. C

Camarothorax sp. D

nr. *Camarothorax* sp.

Ficomila sp.

Sycophila sp. A

Sycophila sp. B

Sycophila sp. C

Sycophila sp. D

Sycophila sp. E

Sycophila sp. F

Sycophila sp. G

Sycophila sp. H

Ormyrus sp.

47. *F. ovata* Vahl (S, E, C, W AFRICA) Forest, often riverine and woodland up to 2100 m
Courtella hamifera Kieffer
Seres armipes Waterston
Seres solweziensis van Noort
Crossogaster ovata van Noort
Sycoryctes sp. A
Sycoryctes sp. B
Watshamiella sp.
Philotrypesis africana Grandi
Philotrypesis sp. B
Philosycus monstruosus (Grandi)
Otitesella sp.

LITERATURE CITED

- BERG, C. C. 1989. Classification and distribution of *Ficus*. *Experientia* 45:605–611.
 ———. 1990. Annotated check-list of the *Ficus* species of the African floristic region, with special reference and a key to the taxa of southern Africa. *Kirkia* 13:253–291.
 BERG, C. C. AND M. E. E. HIJMAN. 1989. *Ficus*. Chapter 11. Pp. 43–86 in *Flora of Tropical East Africa*, R. M. Polhill, ed. A. A. Balkema, Rotterdam.
 BERG, C. C., M. E. E. HIJMAN AND J. C. A. WEERDENBURG. 1984. Moracées (incl. Cécropiacées). *Flore du Gabon* 26:1–276.
 ———. 1985. Moracées (incl. Cécropiacées). *Flore du Cameroun* 28:1–298.
 BERG, C. C. AND J. T. WIEBES. 1992. African Fig Trees and Fig Wasps. K. ned. Akad. Wet., Verhandelungen Afdeling Natuurkunde, Tweede Reeks, Deel 89, Amsterdam. 298 pp.
 BOUËEK, Z. 1988. Australasian Chalcidoidea (Hymenoptera). A biosystematic revision of genera of fourteen families with a reclassification of species. C.A.B. International, Wallingford. 832 pp.
 BRONSTEIN, J. L. 1992. Seed predators as mutualists: ecology and evolution of the fig/pollinator interaction. Pp. 1–44 in *Insect-Plant Interactions Vol IV*, E. Bernays, ed. CRC Press, London.
 BRONSTEIN, J. L. AND M. HOSSAERT-MCKEY. 1996. Variation in reproductive success within a sub-tropical fig/pollinator mutualism. *J. Biogeog.* 23:433–466.
 COE, M., N. MCWILLIAM, G. STONE, AND M. PACKER, eds. 1999. Mkomazi: the ecology, biodiversity and conservation of a Tanzanian savanna. Royal Geographical Society (with The Institute of British Geographers), London. 608 pp.
 COMPTON, S. G. AND B. A. HAWKINS. 1992. Determinants of species richness in southern African fig wasp assemblages. *Oecologia* 91:68–74.
 COMPTON, S. G., K. C. HOLTON, V. K. RASHBROOK, S. VAN NOORT, S. L. VINCENT, AND A. B. WARE. 1991. Studies of *Ceratosolen galili*, a non-pollinating agaonid fig wasp. *Biotropica* 23:188–194.
 COMPTON, S. G. AND S. VAN NOORT. 1992. Southern African fig wasps (Hymenoptera: Chalcidoidea): resource utilization and host relationships. *Proc. Kon. Ned. Akad. v. Wetensch.* 95:423–435.
 COMPTON, S. G., J. Y. RASPLUS, AND A. B. WARE. 1994. African fig wasp parasitoid communities. Pp. 343–368 in *Parasitoid Community Ecology*, B. A. Hawkins and W. Sheenan, eds. Oxford University Press, Oxford.
 GALIL, J. 1977. Fig Biology. *Endeavour* 1:52–56.
 GAUTIER-HION, A. AND G. MICHALOUD. 1989. Are figs always keystone resources for tropical frugivorous vertebrates? A test in Gabon. *Ecology* 70:1826–1833.
 JANZEN, D. H. 1979. How to be a fig. *A. Rev. Ecol. Syst.* 10:13–51.
 KERDELHUÉ, C. AND J. Y. RASPLUS. 1996a. Non-pollinating Afrotropical fig wasps affect the fig-pollinator mutualism in *Ficus* within the subgenus *Sycomorus*. *Oikos* 75:3–14.

- . 1996b. The evolution of dioecy among *Ficus* (Moraceae): an alternative hypothesis involving non-pollinating fig wasp pressure on the fig-pollinator mutualism. *Oikos* 77:163–166.
- MICHALOUD, G. 1982. *Figuiers tropicaux et pollinisation*. Film, réalisation A. R. Devez, production Service du Film de Recherche Scientifique, Paris.
- . 1988. *Aspects de la reproduction des figuiers monoïques en forêt équatoriale africaine*. Thèse d'Université. Université de Montpellier II, France. 256 pp.
- MICHALOUD, G., S. CARRIÈRE, AND M. KOBBI. 1996. Exceptions to the one:one relationship between African fig trees and their fig wasp pollinators: possible evolutionary scenarios. *J. Biogeog.* 23:513–520.
- MICHALOUD, G., S. MICHALOUD-PELLETIER, J. T. WIEBES, AND C. C. BERG. 1985. The co-occurrence of two pollinating species of fig wasp and one species of fig. *Proc. Kon. Ned. Akad. v. Wetensch.* 88:93–119.
- NIEDER, J., J. PROSPERI, AND G. MICHALOUD. 2001. Epiphytes and their contribution to canopy diversity. *Plant Ecology* 153:51–63.
- RAMIREZ, W. B. 1970. Host specificity of fig wasps (Agaonidae). *Evolution*, N.Y. 24:680–691.
- RASPLUS, J. Y. 1996. The one-to-one species specificity of the *Ficus*-Agaoninae mutualism: how causal? Pp. 639–649 in *The biodiversity of African plants*, L. J. G. van der Maesen, X. M. van den Burgt and J. M. van den Medenbrah de Rooy, eds. Kluwer Academic Publishers, Dordrecht.
- RASPLUS, J. Y., C. KERDELHUÉ, I. LE CLAINCHE, AND G. MONDOR. 1998. Molecular phylogeny of fig wasps. Agaonidae are not monophyletic. *C. R. Acad. Sci. Paris, Sciences de la vie* 321:517–527.
- SOSEF, M. S. M. 1994. Glacial rainforest refuges and begonias. *Wageningen Agric. Univ. Papers* 94:113–121.
- ULENBERG, S. A. 1985. The systematics of the fig wasp parasites of the genus *Apocrypta* Coquerel. *Verh. K. ned. Akad. Wet., Tweede Reeks* 83:1–176.
- VAN NOORT, S. 1993a. Systematics of the sycoecine fig wasps (Agaonidae, Chalcidoidea, Hymenoptera), I (*Seres*). *Proc. Kon. Ned. Akad. v. Wetensch.* 96:233–251.
- . 1993b. Systematics of the sycoecine fig wasps (Agaonidae, Chalcidoidea, Hymenoptera), II (*Sycoecus*). *Proc. Kon. Ned. Akad. v. Wetensch.* 96:449–475.
- . 1994a. Systematics of the sycoecine fig wasps (Agaonidae, Chalcidoidea, Hymenoptera), III (*Crossogaster*). *Proc. Kon. Ned. Akad. v. Wetensch.* 97:83–122.
- . 1994b. Systematics of the sycoecine fig wasps (Agaonidae, Chalcidoidea, Hymenoptera), IV (*Philocaenus*, in part). *Proc. Kon. Ned. Akad. v. Wetensch.* 97:311–339.
- . 1994c. Systematics of the sycoecine fig wasps (Agaonidae, Chalcidoidea, Hymenoptera), V (*Philocaenus* concluded, generic key, checklist). *Proc. Kon. Ned. Akad. v. Wetensch.* 97:341–375.
- VAN NOORT, S. AND S. G. COMPTON. 1999. Fig wasps (Agaonidae, Hymenoptera) and fig trees (Moraceae) of Mkomazi. Pp. 299–320 in *Mkomazi: the ecology, biodiversity and conservation of a Tanzanian Savanna*, M. J. Coe, N. C. McWilliam, G. N. Stone, and M. Packer, eds. Royal Geographical Society (with The Institute of British Geographers), London. 608 pp.
- VAN NOORT, S. AND J. Y. RASPLUS. 1997. Revision of the otiteselline fig wasps (Hymenoptera, Chalcidoidea, Agaonidae), I: the *Otitesella digitata* species-group of the Afrotropical region, with a key to Afrotropical species of *Otitesella* Westwood. *African Entomology* 5:125–147.
- WEST, S. A. AND E. A. HERRE. 1994. The ecology of the New World fig-parasitizing wasps *Idarnes* and implication for the evolution of the fig-pollinator mutualism. *Proc. R. Soc. Lond. B* 258:67–72.
- WEST, S. A., E. A. HERRE, D. M. WINDSOR, AND P. R. S. GREEN. 1996. The ecology and evolution of the New World non-pollinating fig wasp communities. *J. Biogeog.* 23:447–458.
- WHITE, F. 1983. *The vegetation of Africa. A descriptive memoir to accompany the Unesco/AETFAT/UNSO vegetation map of Africa*. Unesco, Paris.
- WHITE, L. AND K. ABERNETHY. 1997. *A Guide to the Vegetation of the Lopé Reserve*. ECOFAC, Gabon. 224 pp.
- WIEBES, J. T. 1979a. Coevolution of figs and their insect pollinators. *Ann. Rev. Ecol. Syst.* 10:1–12.
- . 1979b. The fig wasp genus *Dolichoris* Hill (Hymenoptera, Chalcidoidea, Agaonidae). *Proc. Kon. Ned. Akad. v. Wetensch.* 82:181–196.
- . 1979c. Fig wasps from Gabon: new species of *Agaon* (Agaonidae) and *Phagoblastus* (Torymidae) (Hymenoptera, Chalcidoidea). *Proc. Kon. Ned. Akad. v. Wetensch.* 82:391–400.
- . 1986. Agaonidae (Hymenoptera Chalcidoidea) and *Ficus* (Moraceae): fig wasps and their figs, I. *Proc. Kon. Ned. Akad. v. Wetensch.* 89:335–355.

- . 1988. Agaonidae (Hymenoptera Chalcidoidea) and *Ficus* (Moraceae): fig wasps and their figs, II (*Alfonsiella*). Proc. Kon. Ned. Akad. v. Wetensch. 91:429–436.
- . 1989. Agaonidae (Hymenoptera Chalcidoidea) and *Ficus* (Moraceae): fig wasps and their figs, V (*Agaon*). Proc. Kon. Ned. Akad. v. Wetensch. 92:395–407.
- WIEBES, J. T. AND S. G. COMPTON. 1990. Agaonidae (Hymenoptera Chalcidoidea) and *Ficus* (Moraceae): fig wasps and their figs, VI (Africa concluded). Proc. Kon. Ned. Akad. v. Wetensch. 93:203–222.



Van Noort, Simon. 2004. "Fig wasp (Hymenoptera: Chalcidoidea: Agaonidae, Pteromalidae, Eurytomidae and Ormyridae) and Ficus (Moraceae) species richness and biogeography of Monts Doudou in southwestern Gabon." *Memoirs of the California Academy of Sciences* 28, 217–233.

View This Item Online: <https://www.biodiversitylibrary.org/item/99512>

Permalink: <https://www.biodiversitylibrary.org/partpdf/70302>

Holding Institution

California Academy of Sciences

Sponsored by

Biodiversity Heritage Library

Copyright & Reuse

Copyright Status: In copyright. Digitized with the permission of the rights holder.

License: <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Rights: <https://biodiversitylibrary.org/permissions>

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.