IX. NATATORES.

178. PLOTUS LEVAILLANTI, Licht.

Plotus levaillanti, Heugl. Orn. N.O.-Afr. p. 1475; Pelz. Verh. Wien, xxxi. 1881, pp. 606, 618.

No. 844. Q. Lado, Sept. 29, 1883.

179. QUERQUEDULA HARTLAUBI, Cass.

Querquedula hartlaubi, Oust. Bull. N. Arch. Mus. Paris, 1879, pl. 6.

No. 4. J. Tingasi, July 20, 1883.

Iris yellowish brown; bill black, with a broad subterminal yellow band; legs brown.

3. Report on a Collection of Reptiles and Batrachians sent by Emin Pasha from Monbuttu, Upper Congo. By Dr. A. GÜNTHER, F.R.S.

[Received January 10, 1888.]

The specimens, 27 in number, belong to the following species:—

TORTOISES.

1. Pelomedusa subrufa, Lac.

LIZARDS.

- 2. Varanus niloticus, L.—Two young specimens.
- 3. Hemidactylus brookii, Gray.
- 4. Lygodactylus gutturalis, Bocage.
- 5. CHAMÆLEON SENEGALENSIS, var. LÆVIGATA, Gray.—Three specimens. Although I adopt Mr. Boulenger's view that C. lævigatus of Gray should not be specifically distinguished from C. senegalensis, still the former may be readily recognized by its conspicuously finer granulation of the skin, and especially the smaller size of the scutella of the side of the head. The type of C. lævigatus came from the Soudan; and the three specimens sent by Emin Pasha agree in the characters mentioned with this geographically not very remote form of Chamæleon.

SNAKES.

- 6. Typhlops eschrichtii, Schleg.—Three specimens.
- 7. Typhlops liberiensis, Hallow.—Two specimens.

- 8. NEUSTEROPHIS ATRATUS, Ptrs. (1877).—This example possesses one anteocular only. The species is new to the collection of the British Museum.
- 9. AHAETULIA EMINI, sp. n.—Ventral shields without keels, 151; anal bifid; upper labials 9, the 4th, 5th, and 6th entering the orbit; 1 anteocular, 2 postoculars; 6 of the lower labials are in contact with the chin-shields; loreal not twice as long as broad; temporal shields 1+2; scales smooth, in 15 rows. Head of moderate size, not elongate or depressed; body and tail moderately slender. Uniform green; skin between the scales black, each scale with a white spot on the basal half of its outer margin. The specimen measures 29 inches, the head being \(\frac{3}{4}\), and the tail 10 long.
 - 10. DASYPELTIS SCABRA, L.
 - 11. PSAMMOPHIS SIBILANS, L.—Three specimens.
- 12. ATRACTASPIS IRREGULARIS, Reinh.—This species has invariably the subcaudals divided into "scutella"—a term which evidently has been misunderstood by Peters, who persisted in applying it to undivided subcaudal shields, properly "scuta," Peters using the term of squamæ for the former. A renewed study of the species of this genus has also shown me that the doubts which Peters cast upon them can only have arisen from the incomplete materials at his disposal.
- 13. ATRACTASPIS ATERRIMA, Gthr.—The specimen is very much shrunk and possesses 299 ventral shields, the type having 274.

FROGS.

- 14. RANA OCCIPITALIS, Gthr.
- 15. RANA MASCARENIENSIS, D. B.—Two specimens.
- 16. Bufo regularis, Reuss.—Two specimens.
- 17. RAPPIA CINCTIVENTRIS, Cope.

Of these 17 Reptiles and Frogs, 9 are almost generally distributed over the African Region; of the remainder 7 are known from various parts of West Africa, whilst not a single species known to be peculiar to East Africa is included in this collection. Although it might thus appear that the Reptilian Fauna of the Upper Congo is rather West-African than East-African in its character, we must not lose sight of the fact that many species extend right across from the West to the East Coast, and that if in the end the Eastern and Western Reptilian Faunas should prove to be sufficiently distinct to call for the establishment of two or more separate zoo-geographical districts, our knowledge of the Reptiles of the central parts is at present much too fragmentary to assist in fixing the boundary line between such districts.



Günther, Albert C. L. G. 1888. "Report on a Collection of Reptiles and Batrachians sent by Emin Pasha from Monbuttu, Upper Congo." *Proceedings of the Zoological Society of London* 1888, 50–51.

https://doi.org/10.1111/j.1469-7998.1888.tb06676.x.

View This Item Online: https://www.biodiversitylibrary.org/item/99299

DOI: https://doi.org/10.1111/j.1469-7998.1888.tb06676.x

Permalink: https://www.biodiversitylibrary.org/partpdf/72935

Holding Institution

Natural History Museum Library, London

Sponsored by

Natural History Museum Library, London

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at https://www.biodiversitylibrary.org.